



Europäisches Patentamt
European Patent Office
Office européen des brevets



(11) Publication number:

0 508 393 A1

(12)

EUROPEAN PATENT APPLICATION

(21) Application number: 92106050.5

(51) Int. Cl.5: **C07D 401/14**, A61K 31/415,
C07D 401/10, C07D 401/06,
C07D 409/14

(22) Date of filing: 06.04.92

(30) Priority: 05.04.91 US 681011
01.04.92 PCT/US92/02439

(43) Date of publication of application:
14.10.92 Bulletin 92/42

(84) Designated Contracting States:
PT


(71) Applicant: G.D. Searle & Co.
P.O. Box 5110
Chicago Illinois 60680(US)

(72) Inventor: Reitz, David B
14814 Pleasant Ridge Court
USA-Chesterfield, Missouri 63017(US)
Inventor: Manning, Robert E.
1298 South Mason Road
USA-St. Louis, Missouri(US)


(74) Representative: Beil, Hans Christoph, Dr. et al
Beil, Wolff und Beil Rechtsanwälte
Adelonstrasse 58
W-6230 Frankfurt am Main 80(DE)

(54) N-arylheteroarylalkyl imidazol-2-one compounds for treatment of circulatory disorders.

(57) A class of N-arylheteroarylalkyl imidazol-2-one compounds is described for use in treatment of circulatory disorders such as hypertension and congestive heart failure. Compounds of particular interest are angiotensin II antagonists of the formula




wherein A is selected from



EP 0 508 393 A1

wherein m is one;

wherein R¹ is selected from methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, 4-methylbutyl, 2-ethylbutyl, n-pentyl, neopentyl, phenyl, methylphenyl, difluorophenyl, benzyl, phenethyl, cyclohexyl, cyclohexylmethyl, cyclohexylethyl, cyclohexanoyl, 1-oxo-2-cyclohexylethyl, benzoyl, 1-oxo-2-phenethyl, 1-oxopropyl, 1-oxobutyl, 1-oxopentyl and 2-hydroxybutyl; wherein R⁰ is hydrido; wherein R² is selected from methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, n-pentyl, isopentyl, neopentyl, phenyl, benzyl, phenethyl, cyclohexyl, cyclohexylmethyl, propylthio, butylthio, and hydroxyalkyl; wherein each of R³, R⁴, R⁶, R⁷, R⁸, R⁹, R¹⁰ and R¹¹ is hydrido and R⁵ must be selected from COOH, SH, PO₃H₂, SO₃H, CONHNH₂, CONHNHSO₂CF₃, OH,



wherein each of R⁴² and R⁴³ is independently selected from chloro, cyano, nitro, trifluoromethyl, methoxycarbonyl and trifluoromethylsulfonyl; or a tautomer thereof or a pharmaceutically-acceptable salt thereof.

Related Application

This is a continuation-in-part of U.S. Application Ser. No. 07/681,011 filed 5 April 1991.

Field of the Invention

Non-peptidic N-arylheteroarylalkyl imidazol-2-one compounds are described for use in treatment of circulatory disorders such as hypertension and congestive heart failure. Of particular interest are angiotensin II antagonist compounds provided by an imidazol-2-one having a arylheteroarylmethyl moiety attached to a nitrogen atom of the imidazol-2-one.

Background of the Invention

The renin-angiotensin system is one of the hormonal mechanisms involved in regulation of pressure/volume homeostasis and in expression of hypertension. Activation of the renin-angiotensin cascade begins with renin secretion from the juxtaglomerular apparatus of the kidney and culminates in the formation of angiotensin II, the primary active species of this system. This octapeptide, angiotensin II, is a potent vasoconstrictor agent and also produces other physiological effects such as promoting aldosterone secretion, promoting sodium and fluid retention, inhibiting renin secretion, increasing sympathetic nervous system activity, increasing vasopressin secretion, causing positive cardiac inotropic effect and modulating other hormonal systems.

Previous studies have shown that antagonizing angiotensin II at its receptors is a viable approach to inhibit the renin-angiotensin system, given the pivotal role of this octapeptide which mediates the actions of the renin-angiotensin system through interaction with various tissue receptors. There are several known angiotensin II antagonists, most of which are peptidic in nature. Such peptidic compounds are of limited use due to their lack of oral bioavailability or their short duration of action. Also, commercially-available peptidic angiotensin II antagonists (e.g., Saralasin) have a significant residual agonist activity which further limit their therapeutic application.


Non-peptidic compounds with angiotensin II antagonist properties are known. For example, the sodium salt of 2-n-butyl-4-chloro-1-(2-chlorobenzyl)imidazole-5-acetic acid has specific competitive angiotensin II antagonist activity as shown in a series of binding experiments, functional assays and in vivo tests [P. C. Wong et al, *J. Pharmacol. Exp. Ther.*, 247(1), 1-7 (1988)]. Also, the sodium salt of 2-butyl-4-chloro-1-(2-nitrobenzyl)imidazole-5-acetic acid has specific competitive angiotensin II antagonist activity as shown in a series of binding experiments, functional assays and in vivo tests [A. T. Chiu et al, *European J. Pharmacol.*, 157, 13-21 (1988)]. A family of 1-benzylimidazole-5-acetate derivatives has been shown to have competitive angiotensin II antagonist properties [A. T. Chiu et al, *J. Pharmacol. Exp. Ther.*, 250(3), 867-874 (1989)]. U.S. Patent No. 4,816,463 to Blankey et al describes a family of 4,5,6,7-tetrahydro-1H-imidazo(4,5-c)-tetrahydropyridine derivatives useful as antihypertensives, some of which are reported to antagonize the binding of labelled angiotensin II to rat adrenal receptor preparation and thus cause a significant decrease in mean arterial blood pressure in conscious hypertensive rats. EP No. 253,310, published 20 January 1988, describes a series of aralkyl imidazole compounds, including in particular a family of biphenylmethyl substituted imidazoles, as antagonists to the angiotensin II receptor. EP No. 323,841 published 12 July 1989 describes four classes of angiotensin II antagonists, namely, biphenylmethylpyrroles, biphenylmethylpyrazoles, biphenylmethyl-1,2,3-triazoles and biphenylmethyl 4-substituted-4H-1,2,4-triazoles, including the compound 3,5-dibutyl-4-[(2'-carboxybiphenyl-4-yl)methyl]-4H-1,2,4-triazole. U.S. Patent No. 4,880,804 to Carini et al describes a family of biphenylmethylbenzimidazole compounds as angiotensin II receptor blockers for use in treatment of hypertension and congestive heart failure.

There are several families of known compounds having one or two oxo substituents on a triazole ring. For example, East German Patent No. 160,447 published 3 August 1983 describes a family of 1,2,4-triazolin-5-one compounds, specifically 2,4-dihydro-4,5-bis(phenylmethyl)-3H-1,2,4-triazol-3-one, for use as herbicides. Belgian Patent No. 806,146 published 16 October 1972 describes a family of triazolinone compounds, including the compound (3-(4-m-chlorophenyl-1-piperazinyl)-propyl)-3,4-diethyl-1,2,4-triazolin-5-one, having tranquilizer, hypotensive and analgesic activities. Belgian Patent No. 631,842 published 28 February 1963 describes a family of 1,2,4-triazolones having hypnotic, tranquilizer, narcotic, sedative and analgetic activities, which includes a class of 4-N-aralkyl-1,2,4-triazol-5-one compounds. EP #7,180 published 15 June 1978 describes a family of 1,2-disubstituted-4-alkyl-1,2,4-triazolidine-3,5-dione compounds having a wide variety of activities, such as antiulcer, bronchodilator, antifertility and cardiovascular-related activities which include antihypertensive, antiarrhythmic, platelet aggregation inhibition and smooth muscle


activities. EP #283,310 published 18 March 1987 describes a family of N¹-diarylmethyl-N²-aminoalkyl-diaza-heterocyclic derivatives for treating cerebral vascular and ischemic diseases and for protecting against anoxia.

5 DESCRIPTION OF THE INVENTION

A class of N-substituted arylheteroarylalkyl imidazol-2-one compounds useful in treating circulatory disorders, particularly cardiovascular disorders, is defined by Formula I:




wherein A is selected from



wherein m is a number selected from one to four, inclusive;


wherein R¹ is selected from alkyl, hydroxyalkyl, haloalkyl, cycloalkyl, cycloalkylalkyl, formyl, thienylalkyl, phenylalkyl, polycycloalkyl, polycycloalkylalkyl, phenyl, halophenyl, alkylphenyl, alkoxyphenyl, cycloalkenyl, cycloalkenylalkyl, aroyl, alkoxyalkyl, alkylcarbonyl, alkylcarbonylalkyl, cycloalkylcarbonyl, cycloalkylalkylcarbonyl, aralkylcarbonyl, alkoxyalkyl, alkenyl, cycloalkenyl, aralkoxycarbonyl, alkynyl, alkylthiocarbonyl, alkylthiothiocarbonyl, arylthiocarbonyl, arylthiothiocarbonyl, aralkylthiocarbonyl, alkylthiocarbonyl, alkylsulfonyl, aralkylsulfonyl, aralkylsulfonyl, arylsulfonyl, arylsulfonyl, heteroaryl having one or more ring atoms selected from oxygen, sulfur and nitrogen atoms, and amido radicals of the formula



wherein X is oxygen atom or sulfur atom;


wherein each of R¹² and R¹³ is independently selected from hydrido, alkyl, cycloalkyl, cyano, amino, monoalkylamino, dialkylamino, hydroxyalkyl, cycloalkylalkyl, alkoxyalkyl, aralkyl and aryl, and wherein R¹² and R¹³ taken together may form a heterocyclic group having five to seven ring members including the nitrogen atom of said amido radical and which heterocyclic group may further contain one or more hetero atoms as ring members selected from oxygen, nitrogen and sulfur atoms and which heterocyclic group may be saturated or partially unsaturated; wherein each of R¹² and R¹³ taken together may form an aromatic heterocyclic group having five ring members including the nitrogen atom of said amido radical and which

wherein each of R⁰ and R² through R¹¹ is independently selected from hydrido, alkyl, hydroxyalkyl, formyl, halo, haloalkyl, cycloalkyl, cycloalkylalkyl, cycloalkylhaloalkyl, cycloalkylcarbonyl, alkoxy, thienylalkyl, phenylalkyl, polycycloalkyl, polycycloalkylalkyl, phenyl, halophenyl, alkylphenyl, alkoxyphenyl, cycloalkenyl, cycloalkenylalkyl, aralkylhaloalkyl, aroyl, aryloxy, aryloxyalkyl, aralkoxy, alkoxyalkyl, alkylcarbonyl, alkylcarbonylalkyl, alkoxycarbonyl, alkenyl, cycloalkenyl, alkynyl, cyano, nitro, carboxyl, carboxyalkyl, alkylcarbonyloxy, alkylcarbonyloxyalkyl, alkoxycarbonylalkyl, aralkoxycarbonylalkyl, aralkylcarbonyloxyalkyl, mercaptocarbonyl, mercaptothiocarbonyl, mercaptoalkyl, alkoxycarbonyloxy, alkylthio, cycloalkylthio, cycloalkylalkylthio, alkylthiocarbonyl, alkylcarbonylthio, alkylthiocarbonyloxy, alkylthiocarbonylthio, alkylthiothiocarbonyl, alkylthiothiocarbonylthio, arylthio, arylthiocarbonyl, arylcarbonylthio, arylthiocarbonyloxy, arylthiocarbonylthio, arylthiothiocarbonyl, arylthiothiocarbonylthio, aralkylthio, aralkylthiocarbonyl, aralkylcarbonylthio, aralkylthiocarbonyloxy, aralkylthiocarbonylthio, alkylthiocarbonyl, aralkylthiocarbonylthio, mercapto, alkylsulfanyl, alkylsulfonyl, aralkylsulfanyl, aralkylsulfonyl, arylsulfanyl, arylsulfonyl, phthalimido, phthalimidoalkyl, heteroaryl, heteroarylalkyl, cycloheteroalkyl, cycloheteroalkylalkyl and cycloheteroalkylcarbonylalkyl wherein each of said heteroaryl- and cyclohetero-containing groups has one or more ring atoms selected from oxygen, sulfur and nitrogen atoms, and wherein each of R⁰ and R² through R¹¹ may be further independently selected from amino and amido radicals of the formula




and wherein any of the foregoing R⁰ through R²⁶, Y and A groups having a substitutable position may be substituted by one or more groups independently selected from hydroxy, alkyl, alkenyl, alkynyl, aralkyl, hydroxyalkyl, haloalkyl, halo, oxo, alkoxy, aryloxy, aralkoxy, aralkylthio, alkoxyalkyl, cycloalkyl, cycloalkylalkyl, aryl, aroyl, cycloalkenyl, cyano, cyanoamino, nitro, alkylcarbonyloxy, alkoxy carbonyloxy, alkylcarbonyl, alkoxy carbonyl, aralkoxy carbonyl, carboxyl, mercapto, mercapto carbonyl, alkylthio, arylthio, alkylthiocar-


bonyl, alkylsulfinyl, alkylsulfonyl, haloalkylsulfonyl, aralkylsulfinyl, aralkylsulfonyl, arylsulfinyl, arylsulfonyl, heteroaryl having one or more ring atoms selected from oxygen, sulfur and nitrogen atoms, and amino and amido radicals of the formula



wherein X is oxygen atom or sulfur atom; wherein each of R²⁷ through R³¹ is independently selected from hydrido, alkyl, cycloalkyl, cycloalkylalkyl, aralkyl, aryl, DR³² and



wherein D is selected from oxygen atom and sulfur atom and R³² is selected from hydrido, alkyl, cycloalkyl, cycloalkylalkyl, aralkyl and aryl; wherein each of R²⁷, R²⁸, R²⁹, R³⁰, R³¹, R³³ and R³⁴ is independently selected from hydrido, alkyl, cycloalkyl, cyano, hydroxyalkyl, haloalkyl, cycloalkylalkyl, alkoxyalkyl, alkylcarbonyl, alkoxy carbonyl, carboxyl, alkylsulfinyl, alkylsulfonyl, arylsulfinyl, arylsulfonyl, haloalkylsulfinyl, haloalkylsulfonyl, aralkyl and aryl, and wherein each of R²⁷, R²⁸, R²⁹, R³⁰, R³¹, R³³ and R³⁴ is further independently selected from amino and amido radicals of the formula



wherein X is oxygen atom or sulfur atom;


wherein each of R³⁵, R³⁶, R³⁷, R³⁸, R³⁹ and R⁴⁰ is independently selected from hydrido, alkyl, cycloalkyl, cyano, hydroxyalkyl, cycloalkylalkyl, alkoxyalkyl, haloalkylsulfinyl, haloalkylsulfonyl, aralkyl and aryl, and wherein each of R²⁸ and R²⁹ taken together and each of R³⁰ and R³¹ taken together may form a heterocyclic group having five to seven ring members including the nitrogen atom of said amino or amido radical, which heterocyclic group may further contain one or more hetero atoms as ring members selected from oxygen, nitrogen and sulfur atoms and which heterocyclic group may be saturated or partially unsaturated; wherein each of R²⁸ and R²⁹ taken together and each of R³³ and R³⁴ taken together may form an aromatic heterocyclic group having five ring members including the nitrogen atom of said amino or amido radical and which aromatic heterocyclic group may further contain one or more additional nitrogen atoms; or a tautomer thereof or a pharmaceutically-acceptable salt thereof.

Compounds of Formula I would be useful in treating a variety of circulatory disorders, including cardiovascular disorders, such as hypertension, congestive heart failure and arteriosclerosis, and to treat other disorders such as glaucoma. These compounds would also be useful as adjunctive therapies. For example, compounds of Formula I may be used in combination with other drugs, such as a diuretic, to treat hypertension. Also, compounds of Formula I could be used in conjunction with certain surgical procedures. For example, these compounds could be used to prevent post-angioplasty re-stenosis. Compounds of Formula I are therapeutically effective in treatment of cardiovascular disorders by acting as antagonists to, or blockers of, the angiotensin II (Ang II) receptor. Compounds of Formula I would be therapeutically effective in treatment of the above-mentioned circulatory and cardiovascular disorders or would be precursors to, or prodrugs of, therapeutically-effective compounds.


The phrase "acidic group selected to contain at least one acidic hydrogen atom", as used to define the -Y_nAmoiety, is intended to embrace chemical groups which, when attached to any of the R⁰ and R³ through

R¹¹ positions of Formula I, confers acidic character to the compound of Formula I. "Acidic character" means proton-donor capability, that is, the capacity of the compound of Formula I to be a proton donor in the presence of a proton-receiving substance such as water. Typically, the acidic group should be selected to have proton-donor capability such that the product compound of Formula I has a pK_a in a range from about one to about twelve. More typically, the Formula I compound would have a pK_a in a range from about two to about seven. An example of an acidic group containing at least one acidic hydrogen atom is carboxyl group (-COOH). Where n is zero and A is -COOH, in the -Y_nA moiety, such carboxyl group would be attached directly to one of the R⁰ and R³ through R¹¹ positions. The Formula I compound may have one -Y_nA moiety attached at one of the R³ through R¹¹ positions, or may have a plurality of such -Y_nA moieties attached at more than one of the R⁰ and R³ through R¹¹ positions, up to a maximum of ten such -Y_nA moieties. There are many examples of acidic groups other than carboxyl group, selectable to contain at least one acidic hydrogen atom. Such other acidic groups may be collectively referred to as "bioisosteres of carboxylic acid" or referred to as "acidic bioisosteres". Specific examples of such acidic bioisosteres are described hereinafter. Compounds of Formula I having the -Y_nA moiety attached at one of positions R⁰, R⁵, R⁶, R⁸, R⁹ and R⁹ would be expected to have preferred properties, while attachment at R⁵ or R⁹ would be more preferred. Compounds of Formula I having the -Y_nA moiety attached at one of positions R⁵, R⁶, R⁸ and R⁹ would be expected to have preferred properties, while attachment at R⁵ or R⁹ would be more preferred. Compounds of Formula I may have one or more acidic protons and, therefore, may have one or more pK_a values. It is preferred, however, that at least one of these pK_a values of the Formula I compound as conferred by the -Y_nA moiety be in a range from about two to about seven. The -Y_nA moiety may be attached to one of the R³ through R¹¹ positions through any portion of the -Y_nA moiety which results in a Formula I compound being relatively stable and also having a labile or acidic proton to meet the foregoing pK_a criteria. For example, where the -Y_nA acid moiety is tetrazole, the tetrazole is attached at the ring carbon tetrazole atom.

A preferred class of compounds consists of those compounds of Formula I wherein m is one; wherein R¹ is selected from alkyl, hydroxyalkyl, haloalkyl, cycloalkyl, cycloalkylalkyl, thienylalkyl, phenylalkyl, polycycloalkyl, polycycloalkylalkyl, phenyl, halophenyl, alkylphenyl, alkoxyphenyl, cycloalkenyl, cycloalkenylalkyl, aroyl, alkoxyalkyl, alkylcarbonyl, cycloalkylcarbonyl, cycloalkylalkylcarbonyl, aralkylcarbonyl, alkoxy carbonyl, alkenyl, cycloalkenyl, alkynyl, cycloalkynyl, alkylthiocarbonyl, arylthiocarbonyl, arylthiothiocarbonyl, aralkylthiocarbonyl, alkylsulfonyl, aralkylsulfonyl, arylsulfonyl, heteroaryl having one or more ring atoms selected from oxygen, sulfur and nitrogen atoms, and amido radicals of the formula



wherein X is oxygen atom or sulfur atom;
 wherein each of R¹² and R¹³ is independently selected from hydrido, alkyl, cycloalkyl, cyano, amino, monoalkylamino, dialkylamino, hydroxyalkyl, cycloalkylalkyl, alkoxyalkyl, aralkyl and aryl;
 wherein each of R⁰ and R² is independently selected from hydrido, alkyl, hydroxyalkyl, formyl, halo, haloalkyl, cycloalkyl, cycloalkylalkyl, cycloalkylhaloalkyl, cycloalkylcarbonyl, alkoxy, thienylalkyl, phenylalkyl, polycycloalkyl, polycycloalkylalkyl, phenyl, halophenyl, alkylphenyl, alkoxyphenyl, cycloalkenyl, cycloalkenylalkyl, aralkylhaloalkyl, aroyl, aryloxy, aryloxyalkyl, aralkoxy, alkoxyalkyl, alkylcarbonyl, alkoxy carbonyl, alkenyl, cycloalkenyl, alkynyl, cycloalkynyl, cyano, nitro, carboxyl, carboxyalkyl, alkylcarbonyloxy, alkylcarbonyloxyalkyl, alkoxy carbonylalkyl, aralkoxy carbonylalkyl, aralkylcarbonyloxyalkyl, mercaptocarbonyl, mercaptothiocarbonyl, mercaptoalkyl, alkoxy carbonyloxy, alkylthio, cycloalkylthio, cycloalkylalkylthio, alkylthiocarbonyl, alkylcarbonylthio, alkylthiocarbonyloxy, alkylthiocarbonylthio, alkylthiothiocarbonyl, arylthio, arylthiocarbonyl, arylcarbonylthio, arylthiocarbonyloxy, arylthiocarbonylthio, arylthiothiocarbonyl, arylthiothiocarbonylthio, aralkylthio, aralkylthiocarbonyl, aralkylthiocarbonyloxy, aralkylthiocarbonylthio, aralkylthiocarbonylthio, mercapto, alkylsulfinyl, alkylsulfonyl, aralkylsulfinyl, aralkylsulfonyl, arylsulfinyl, arylsulfonyl, phthalimido, phthalimidoalkyl, heteroaryl, heteroarylalkyl, cycloheteroalkyl, cycloheteroalkylalkyl and cycloheteroalkylcarbonylalkyl wherein each of said heteroaryl- and cycloheteroalkyl-containing groups has one or more hetero ring atoms selected from oxygen, sulfur and nitrogen atoms, and wherein each of R⁰ and R² through R¹¹ may be further independently selected from amino and amido radicals of the formula




wherein X is selected from oxygen atom or sulfur atom;

15 wherein each n is a number independently selected from zero to six, inclusive;

wherein each of R¹⁴ through R²⁶ is independently selected from hydrido, alkyl, cycloalkyl, cyano, amino, monoalkylamino, dialkylamino, hydroxyalkyl, cycloalkylalkyl, alkoxyalkyl, aralkyl and aryl;

wherein each of R³ through R¹¹ is independently selected from hydrido, hydroxy, alkyl, hydroxyalkyl, halo, haloalkyl, cycloalkyl, cycloalkylalkyl, alkoxy, aralkyl, aryl, aryloxy, aralkoxy, alkoxyalkyl, alkylcarbonyl, alkoxyalkyl, alkenyl, cycloalkenyl, alkynyl, cycloalkynyl, cyano, nitro, carboxyl, alkylcarbonyloxy, mercaptoalkyl, mercaptothiocarbonyl, alkoxyalkoxy, alkylthio, alkylthiocarbonyl, alkylcarbonylthio, alkylthiocarbonyloxy, alkylthiocarbonylthio, alkylthiothiocarbonyl, arylthio, arylthiocarbonyl, arylcarbonylthio, arylthiocarbonyloxy, arylthiothiocarbonyl, aralkylthio, aralkylthiocarbonyl, aralkylcarbonylthio, aralkylthiocarbonyloxy, aralkylthiocarbonylthio, aralkylthiocarbonyl, aralkylthiocarbonylthio, mercapto, alkylsulfonyl, aralkylsulfonyl and arylsulfonyl, and amino and amido radicals of the formula



35 wherein X is oxygen atom or sulfur atom;


wherein each of R¹⁴, R¹⁵, R¹⁶, R¹⁷, R¹⁸ and R¹⁹ is independently selected from hydrido, alkyl, cycloalkyl, cyano, hydroxyalkyl, cycloalkylalkyl, alkoxyalkyl, aralkyl and aryl;

and wherein each of R⁰ and R³ through R¹¹ may be further independently selected from acidic moieties of the formula




wherein n is a number selected from zero through three, inclusive; wherein A is an acidic group selected from acids containing one or more atoms selected from oxygen, sulfur, phosphorus and nitrogen atoms, and wherein said acidic group is selected to contain at least one acidic hydrogen atom, and the amide, ester and salt derivatives of said acidic moieties; wherein Y is a spacer group independently selected from one or more of alkyl, cycloalkyl, cycloalkylalkyl, alkenyl, alkynyl, aryl, aralkyl and heteroaryl having one or more ring atoms selected from oxygen, sulfur and nitrogen atoms;

and wherein any of the foregoing R⁰ through R²⁶, Y and A groups having a substitutable position may be substituted by one or more groups independently selected from hydroxy, alkyl, alkenyl, aralkyl, hydroxyalkyl, halo, haloalkyl, oxo, alkoxy, aryloxy, aralkoxy, alkoxyalkyl, alkylcarbonyl, alkoxyalkoxy, carboxyl, cyano, nitro, alkylsulfonyl, haloalkylsulfonyl, aryl, aralkyl, mercaptoalkyl, alkylthio and alkylthiocarbonyl, and amino and amido radicals of the formula




wherein X is oxygen atom or sulfur atom; wherein each of R²⁷ through R³¹ is independently selected from hydrido, alkyl, cycloalkyl, cycloalkylalkyl, aralkyl, aryl, and DR³² and




wherein D is selected from oxygen atom and sulfur atom, and R³² is selected from hydrido, alkyl, cycloalkyl, cycloalkylalkyl, aralkyl and aryl; wherein each of R²⁷, R²⁸, R²⁹, R³⁰, R³¹, R³³ and R³⁴ is independently selected from hydrido, alkyl, cycloalkyl, cyano, hydroxyalkyl, haloalkyl, cycloalkylalkyl, alkoxyalkyl, alkanoyl, alkoxy carbonyl, carboxyl, haloalkylsulfinyl, haloalkylsulfonyl, aralkyl and aryl; or a tautomer thereof or a pharmaceutically-acceptable salt thereof.

A more preferred class of compounds consists of those compounds of Formula I wherein m is one; wherein R¹ is selected from alkyl, hydroxyalkyl, haloalkyl, cycloalkyl, cycloalkylalkyl, thienylalkyl, phenylalkyl, polycycloalkyl, polycycloalkylalkyl, phenyl, halophenyl, alkylphenyl, alkoxyphenyl, cycloalkenyl, cycloalkenylalkyl, aroyl, alkoxyalkyl, alkylcarbonyl, cycloalkylcarbonyl, cycloalkylalkylcarbonyl, aralkylcarbonyl, alkoxy carbonyl, alkenyl, cycloalkenyl, alkynyl, cycloalkynyl, alkylsulfonyl, aralkylsulfonyl and arylsulfonyl, and amido radicals of the formula



wherein each of R¹² and R¹³ is independently selected from hydrido, alkyl, cycloalkyl, cyano, amino, monoalkylamino, dialkylamino, hydroxyalkyl, cycloalkylalkyl, alkoxyalkyl, aralkyl and aryl; wherein each of R⁰ and R² is independently selected from hydrido, alkyl, hydroxyalkyl, formyl, halo, haloalkyl, cycloalkyl, cycloalkylalkyl, cycloalkylhaloalkyl, cycloalkylcarbonyl, alkoxy, thienylalkyl, phenylalkyl, polycycloalkyl, polycycloalkylalkyl, phenyl, halophenyl, alkylphenyl, alkoxyphenyl, cycloalkenyl, cycloalkenylalkyl, aralkylhaloalkyl, aroyl, aryloxy, aryloxyalkyl, aralkoxy, alkoxyalkyl, alkylcarbonyl, alkoxy carbonyl, alkenyl, cycloalkenyl, alkynyl, cycloalkynyl, cyano, nitro, carboxyl, carboxyalkyl, alkylcarbonyloxy, mercapto carbonyl, alkoxy carbonyloxy, alkylcarbonyloxyalkyl, alkoxy carbonylalkyl, aralkoxy carbonylalkyl, aralkylcarbonyloxyalkyl, alkylthio, cycloalkylthio, cycloalkylalkylthio, arylthio, aralkylthio, aralkylthiocarbonylthio, mercapto, alkylsulfinyl, alkylsulfonyl, aralkylsulfinyl, aralkylsulfonyl, arylsulfinyl, arylsulfonyl, phthalimido, phthalimidoalkyl, heteroaryl, heteroarylalkyl, cycloheteroalkyl, cycloheteroalkylalkyl and cycloheteroalkylcarbonylalkyl wherein each of said heteroaryl- and cycloheteroalkyl-containing groups has one or more hetero ring atoms selected from oxygen, sulfur and nitrogen atoms, and wherein each of R⁰ and R² may be further independently selected from amino and amido radicals of the formula




wherein X is selected from oxygen atom or sulfur atom;

15 wherein each n is a number independently selected from zero to six, inclusive;

wherein each of R¹⁴ through R²⁶ is independently selected from hydrido, alkyl, cycloalkyl, cyano, amino, monoalkylamino, dialkylamino, hydroxyalkyl, cycloalkylalkyl, alkoxyalkyl, aralkyl and aryl;

wherein each of R³ through R¹¹ is independently selected from hydrido, hydroxy, alkyl, hydroxyalkyl, halo, haloalkyl, cycloalkyl, alkoxy, aralkyl, aryl, aroyl, aryloxy, aralkoxy, alkoxyalkyl, alkylcarbonyl, alkoxy carbonyl, alkenyl, cycloalkenyl, alkynyl, cyano, nitro, carboxyl, alkylcarbonyloxy, mercaptocarbonyl, alkoxy carbonyloxy, alkylthio, arylthio, aralkylthio, mercapto, alkylsulfonyl, aralkylsulfonyl and arylsulfonyl, and amino and amido radicals of the formula



wherein each of R¹⁴, R¹⁵, R¹⁶, R¹⁷, R¹⁸ and R¹⁹ is independently selected from hydrido, alkyl, cycloalkyl, cyano, amino, monoalkylamino, dialkylamino, hydroxyalkyl, cycloalkylalkyl, alkoxyalkyl, aralkyl and aryl;

35 and wherein each of R⁰ and R³ through R¹¹ may be further independently selected from acidic moieties of the formula

$$-Y_{PA}$$

wherein n is a number selected from zero through three, inclusive;

40 wherein A is selected from carboxylic acid and bioisosteres of carboxylic acid selected from -OH, -SH.




-S-WH,




wherein each W is independently selected from oxygen atom, sulfur atom and NR³⁹; wherein each of R³⁵, R³⁶, R³⁷, R³⁸ and R³⁹ is independently selected from hydrido, alkyl, haloalkyl, haloalkylsulfonyl, haloalkyl-


carbonyl, cycloalkyl, cycloalkylalkyl, aryl and aralkyl; wherein each of R^{35} , R^{36} , R^{37} and R^{39} may be further independently selected from amino radical of the formula



- wherein each of R^{40} and R^{41} is independently selected from hydrido, alkyl, cycloalkyl, hydroxyalkyl, haloalkyl, cycloalkylalkyl, alkoxyalkyl, aralkyl and aryl, and wherein R^{40} and R^{41} taken together may form a heterocyclic group having five to seven ring members including the nitrogen atom of said amino radical, which heterocyclic group may further contain one or more hetero atoms as ring members selected from oxygen, nitrogen and sulfur atoms and which heterocyclic group may be saturated or partially unsaturated;
- wherein R^{40} and R^{41} taken together may form an aromatic heterocyclic group having five ring members including the nitrogen atom of said amino radical and which aromatic heterocyclic group may further contain one or more hetero atoms as ring atoms selected from oxygen, nitrogen and sulfur atoms; wherein each of R^{36} and R^{37} may be further independently selected from hydroxy, alkoxy, alkylthio, aryloxy, arylthio, aralkylthio and aralkoxy; and the amide, ester and salt derivatives of said acidic groups;
- wherein said bioisostere of carboxylic acid may be further selected from heterocyclic acidic groups consisting of heterocyclic rings of four to about nine ring members, which heterocyclic ring contains at least one hetero atom selected from oxygen, sulfur and nitrogen atoms, which heterocyclic ring may be saturated, fully unsaturated or partially unsaturated, and which heterocyclic ring may be attached at a single position selected from R^3 through R^{11} or may be attached at any two adjacent positions selected from R^3 through R^{11} so as to form a fused-ring system with one of the phenyl rings of Formula I; and the amide, ester and salt derivatives of said heterocyclic acidic groups;
- wherein Y is a spacer group independently selected from one or more of alkyl, cycloalkyl, cycloalkylalkyl, alkenyl, aryl and aralkyl;
- and wherein any of the foregoing R^0 through R^{26} and R^{35} through R^{41} , Y and A groups having a substitutable position may be substituted by one or more groups independently selected from hydroxy, alkyl, alkenyl, aralkyl, hydroxyalkyl, halo, oxo, haloalkyl, alkoxy, aryloxy, aralkoxy, alkoxyalkyl, alkylcarbonyl, alkoxyalkyl, carboxyl, cyano, nitro, alkylsulfonyl, haloalkylsulfonyl, aryl, aralkyl, mercaptocarbonyl, alkylthio and alkylthiocarbonyl, and amino and amido radicals of the formula



wherein X is selected from oxygen atom and sulfur atom;
wherein R^{27} is selected from hydrido, alkyl, cycloalkyl, cycloalkylalkyl, aralkyl, aryl and DR^{32} and



- wherein D is selected from oxygen atom and sulfur atom;
wherein R^{32} is selected from hydrido, alkyl, cycloalkyl, cycloalkylalkyl, aralkyl and aryl;
wherein each of R^{27} , R^{28} , R^{29} , R^{30} , R^{31} , R^{33} and R^{34} is independently selected from hydrido, alkyl, cycloalkyl, cyano, hydroxyalkyl, haloalkyl, cycloalkylalkyl, alkoxyalkyl, alkanoyl, alkoxyalkyl, carboxyl, haloalkylsulfonyl, haloalkylsulfonyl, aralkyl and aryl;
or a tautomer thereof or a pharmaceutically-acceptable salt thereof.

An even more preferred class of compounds consists of those compounds of Formula I wherein m is one; wherein R^1 is selected from alkyl, hydroxyalkyl, haloalkyl, cycloalkyl, cycloalkylalkyl, thienylalkyl,


phenylalkyl, polycycloalkyl, polycycloalkylalkyl, phenyl, halophenyl, alkylphenyl, alkoxyphenyl, cycloalkenyl, cycloalkenylalkyl, aroyl, alkoxyalkyl, alkylcarbonyl, cycloalkylcarbonyl, cycloalkylalkylcarbonyl, aralkylcarbonyl, alkenyl, cycloalkenyl, alkynyl, mercaptocarbonyl, alkylsulfonyl, aralkylsulfonyl, arylsulfonyl and amido radicals of the formula

5



wherein each of R¹² and R¹³ is independently selected from hydrido, alkyl, cycloalkyl, cyano, amino, monoalkylamino, dialkylamino, hydroxyalkyl, cycloalkylalkyl, alkoxyalkyl, aralkyl and aryl;

15 wherein each of R⁰ and R² is independently selected from hydrido, alkyl, hydroxyalkyl, formyl, halo, haloalkyl, cycloalkyl, cycloalkylalkyl, cycloalkylhaloalkyl, cycloalkylcarbonyl, alkoxy, thienylalkyl, phenylalkyl, polycycloalkyl, polycycloalkylalkyl, phenyl, halophenyl, alkylphenyl, alkoxyphenyl, cycloalkenyl, cycloalkenylalkyl, aralkylhaloalkyl, aroyl, aryloxy, aryloxyalkyl, aralkoxy, alkoxyalkyl, alkylcarbonyl, alkoxy carbonyl, alkenyl, cycloalkenyl, alkynyl, cycloalkynyl, cyano, nitro, carboxyl, carboxyalkyl, alkylcarbonyloxy, alkylcarbonyloxyalkyl, alkoxy carbonylalkyl, aralkoxy carbonylalkyl, aralkylcarbonyloxyalkyl, mercaptocarbonyl, mercaptoalkyl, alkoxy carbonyloxy, alkylthio, cycloalkylthio, cycloalkylalkylthio, arylthio, aralkylthio, mercapto, alkylsulfanyl, alkylsulfanyl, aralkylsulfanyl, aralkylsulfonyl, arylsulfanyl, arylsulfonyl, phthalimido, phthalimidoalkyl, heteroaryl, heteroarylalkyl, cycloheteroalkyl, cycloheteroalkylalkyl and cycloheteroalkylcarbonylalkyl
20 wherein each of said heteroaryl- and cycloheteroalkyl-containing groups has one or more hetero ring atoms selected from oxygen, sulfur and nitrogen atoms, and wherein each of R² through R¹¹ may be further
25 independently selected from amino and amido radicals of the formula



wherein X is selected from oxygen atom and sulfur atom;

wherein each n is a number independently selected from zero to six, inclusive;



wherein each of R¹⁴ through R²⁶ is independently selected from hydrido, alkyl, cycloalkyl, cyano, amino, monoalkylamino, dialkylamino, hydroxyalkyl, cycloalkylalkyl, alkoxyalkyl, aralkyl and aryl;

45 wherein each of R³ through R¹¹ is independently selected from hydrido, hydroxy, alkyl, hydroxyalkyl, halo, haloalkyl, cycloalkyl, alkoxy, aralkyl, aryl, aroyl, aryloxy, aralkoxy, alkoxyalkyl, alkylcarbonyl, alkoxy carbonyl, alkenyl, cycloalkenyl, alkynyl, cyano, nitro, carboxyl, alkylthio, aralkylthio and mercapto; and wherein each of R⁰ and R³ through R¹¹ may be further independently selected from acidic moieties of the formula


$$-Y_D A$$

wherein n is a number selected from zero through three, inclusive; wherein A is selected from carboxylic acid and bioisosteres of carboxylic acid selected from

-OH, -SH,




wherein each W is independently selected from oxygen atom, sulfur atom and NR³⁹; wherein each of R³⁵, R³⁸ and R³⁹ is independently selected from hydrido, alkyl, haloalkyl, haloalkylsulfonyl, haloalkylcarbonyl, cycloalkyl, cycloalkylalkyl, aryl and aralkyl; wherein each of R³⁵ and R³⁹ may be further independently selected from amino radical of the formula




wherein each of R⁴⁰ and R⁴¹ is independently selected from hydrido, alkyl, cycloalkyl, hydroxyalkyl, haloalkyl, cycloalkylalkyl, alkoxyalkyl, aralkyl and aryl, and wherein R⁴⁰ and R⁴¹ taken together may form a heterocyclic group having five to seven ring members including the nitrogen atom of said amino radical, which heterocyclic group may further contain one or more hetero atoms as ring members selected from oxygen, nitrogen and sulfur atoms, and which heterocyclic group may be saturated or partially unsaturated; wherein R⁴⁰ and R⁴¹ taken together may form an aromatic heterocyclic group having five ring members including the nitrogen atom of said amino radical and which aromatic heterocyclic group may further contain one or more hetero atoms as ring atoms selected from oxygen, nitrogen and sulfur atoms; and the amide, ester and salt derivatives of said acidic groups; wherein said bioisostere of carboxylic acid may be further selected from heterocyclic acidic groups consisting of heterocyclic rings of four to about nine ring members, which ring contains at least one hetero atom, selected from oxygen, sulfur and nitrogen atoms, which heterocyclic ring may be saturated, fully unsaturated or partially unsaturated, and which heterocyclic ring may be attached at a single position selected from R³ through R¹¹ or may be attached at any two adjacent positions selected from R³ through R¹¹ so as to form a fused-ring system with one of the phenyl rings of Formula I; and the amide, ester and salt derivatives of said heterocyclic acidic groups; wherein Y is a spacer group independently selected from one or more of alkyl, cycloalkyl, cycloalkylalkyl, alkenyl, aryl and aralkyl; wherein each of R⁰ through R²⁶, R³⁵ and R³⁸ through R⁴¹, Y and A independently may be substituted at any substitutable position with one or more groups selected from alkyl, hydroxy, halo, oxo, haloalkyl, alkoxycarbonyl, cyano, nitro, alkylsulfonyl, haloalkylsulfonyl, aryl, aralkyl, alkoxy, aryloxy and aralkoxy; or a tautomer thereof or a pharmaceutically-acceptable salt thereof.

A highly preferred class of compounds within Formula I consists of those compounds wherein m is one; wherein R¹ is selected from alkyl, hydroxyalkyl, haloalkyl, cycloalkyl, cycloalkylalkyl, thienylalkyl, phenylalkyl, polycycloalkyl, polycycloalkylalkyl, phenyl, halophenyl, alkylphenyl, alkoxyphenyl, cycloalkenyl, cycloalkenylalkyl, aroyl, alkoxyalkyl, alkylcarbonyl, cycloalkylcarbonyl, cycloalkylalkylcarbonyl, aralkylcarbonyl, alkenyl, alkynyl, alkylsulfonyl, aralkylsulfonyl, arylsulfonyl and amido radicals of the formula



wherein each of R^{12} and R^{13} is independently selected from hydrido, alkyl, cycloalkyl, cyano, amino, hydroxyalkyl, alkoxyalkyl, phenalkyl and phenyl;
 wherein each of R^0 and R^2 is independently selected from hydrido, alkyl, hydroxyalkyl, formyl, halo, haloalkyl, cycloalkyl, cycloalkylalkyl, cycloalkylhaloalkyl, cycloalkylcarbonyl, alkoxy, thienylalkyl, phenylalkyl, polycycloalkyl, polycycloalkylalkyl, phenyl, halophenyl, alkylphenyl, alkoxyphenyl, cycloalkenyl, cycloalkenylalkyl, aralkylhaloalkyl, benzoyl, phenoxy, phenoxyalkyl, phenalkyloxy, phenylthio, phenalkylthio, aralkoxy, alkoxyalkyl, alkylcarbonyl, alkoxy carbonyl, alkenyl, cycloalkenyl, alkynyl, cyano, nitro, carboxyl, carboxyalkyl, alkylcarbonyloxy, alkylcarbonyloxyalkyl, alkoxy carbonylalkyl, aralkoxy carbonylalkyl, aralkylcarbonyloxyalkyl, mercaptocarbonyl, mercaptoalkyl, alkoxy carbonyloxy, alkylthio, cycloalkylthio, cycloalkylalkylthio, phthalimido, phthalimidoalkyl, heteroaryl, heteroarylalkyl, cycloheteroalkyl, cycloheteroalkylalkyl and cycloheteroalkylcarbonylalkyl wherein each of said heteroaryl- and cycloheteroalkyl- containing groups has one or more hetero ring atoms selected from oxygen, sulfur and nitrogen atoms, and wherein each of R^0 and R^2 through R^{11} may be further independently selected from amino and amido radicals of the formula



wherein X is selected from oxygen atom and sulfur atom;
 wherein each n is a number independently selected from zero to six, inclusive;
 wherein each of R^{14} through R^{26} is independently selected from hydrido, alkyl, cycloalkyl, cyano, amino, hydroxyalkyl, alkoxyalkyl, phenalkyl and phenyl;
 wherein each of R^3 through R^{11} is independently selected from hydrido, hydroxy, alkyl, hydroxyalkyl, halo, haloalkyl, cycloalkyl, alkoxy, phenalkyl, phenyl, benzoyl, phenoxy, phenalkyloxy, alkoxyalkyl, alkylcarbonyl, alkoxy carbonyl, alkenyl, cyano, nitro, carboxyl, alkylthio and mercapto;
 and wherein each of R^0 and R^3 through R^{11} may be further independently selected from acidic moieties of the formula

$-Y_nA$

wherein n is a number selected from zero through two, inclusive; wherein A is selected from carboxylic acid and bioisosteres of carboxylic acid selected from $-OH$, $-SH$,




$-S-WH$,



wherein each W is independently selected from oxygen atom, sulfur atom and NR^{39} ; wherein each of R^{35} ,

R^{38} and R^{39} is independently selected from hydrido, alkyl, haloalkyl, haloalkylsulfonyl, haloalkylcarbonyl, cycloalkyl, phenyl and benzyl; wherein each of R^{35} and R^{39} may be further independently selected from amino radical of the formula



10

wherein each of R^{40} and R^{41} is independently selected from hydrido, alkyl, cycloalkyl, hydroxyalkyl, haloalkyl, alkoxyalkyl, benzyl and phenyl; and the amide, ester and salt derivatives of said acidic groups; wherein said bioisostere of carboxylic acid may be further selected from heterocyclic acidic groups consisting of heterocyclic rings of four to about nine ring members, which ring contains at least one hetero

15

atom, selected from oxygen, sulfur and nitrogen atoms, which heterocyclic ring may be saturated, fully unsaturated or partially unsaturated, and which heterocyclic ring may be attached at a single position selected from R^3 through R^{11} or may be attached at any two adjacent positions selected from R^3 through R^{11} so as to form a fused-ring system with one of the phenyl rings of Formula I; and the amide, ester and salt derivatives of said heterocyclic acidic groups;

20

wherein Y is a spacer group independently selected from one or more of alkyl, cycloalkyl, cycloalkylalkyl, alkenyl, phenyl, phenalkyl and aralkyl;

wherein each of R^0 through R^{26} , R^{35} and R^{38} through R^{41} , Y and A and independently may be substituted at any substitutable position with one or more groups selected from alkyl, cycloalkyl, cycloalkylalkyl, hydroxy, halo, oxo, haloalkyl, alkoxy, carbonyl, cyano, nitro, alkylsulfonyl, haloalkylsulfonyl, aryl, aralkyl, alkoxy, aryloxy and aralkoxy;

25


or a tautomer thereof or a pharmaceutically-acceptable salt thereof.

An even more highly preferred class of compounds consists of those compounds of Formula I wherein m is one; wherein R^1 is selected from alkyl, hydroxyalkyl, haloalkyl, cycloalkyl, cycloalkylalkyl, thienylalkyl, phenylalkyl, polycycloalkyl, polycycloalkylalkyl, phenyl, halophenyl, alkylphenyl, alkoxyphenyl, cycloalkenyl, cycloalkenylalkyl, benzoyl, alkoxyalkyl, alkylcarbonyl, alkoxy, carbonyl, cycloalkylcarbonyl, cycloalkylalkylcarbonyl, aralkylcarbonyl, alkenyl and alkynyl;

30

where each of R^0 and R^2 is independently selected from hydrido, alkyl, aminoalkyl, hydroxyalkyl, formyl, halo, haloalkyl, cycloalkyl, cycloalkylalkyl, cycloalkylhaloalkyl, cycloalkylcarbonyl, alkoxy, thienylalkyl, phenylalkyl, polycycloalkyl, polycycloalkylalkyl, phenyl, halophenyl, alkylphenyl, alkoxyphenyl, cycloalkenyl, cycloalkenylalkyl, aralkylhaloalkyl, benzoyl, phenoxy, phenoxyalkyl, phenalkyloxy, phenylthio, phenalkylthio, aralkoxy, alkoxyalkyl, acetyl, alkoxy, carbonyl, alkenyl, cycloalkenyl, alkynyl, cyano, nitro, carboxyl, carboxyalkyl, alkylcarbonyl, alkylcarbonyloxy, mercaptoalkyl, mercaptocarbonyl, alkoxy, carbonyloxy, alkylcarbonyloxyalkyl, alkoxy, carbonylalkyl, aralkoxy, carbonylalkyl, aralkylcarbonyloxyalkyl, phthalimido, phthalimidoalkyl, imidazoalkyl, tetrazole, tetrazolealkyl, alkylthio, cycloalkylthio, cycloalkylalkylthio, and amino and amido radicals of the formula

40



55


wherein X is selected from oxygen atom and sulfur atom;

wherein each n is a number independently selected from zero to six, inclusive;


wherein each of R^{14} through R^{26} is independently selected from hydrido, alkyl, cycloalkyl, cyano, amino, hydroxyalkyl, alkoxyalkyl, phenalkyl and phenyl;

wherein each of R^3 through R^{11} is independently selected from hydrido, hydroxy, alkyl, hydroxyalkyl, halo, haloalkyl, alkoxy, phenyl, benzoyl, phenoxy, alkoxyalkyl, acetyl, alkoxycarbonyl, alkenyl, cyano, nitro, carboxyl, alkylthio and mercapto;

and wherein each of R^0 and R^3 through R^{11} may be further independently selected from acidic moieties consisting of CO_2H , CO_2CH_3 , SH , CH_2SH , $\text{C}_2\text{H}_4\text{SH}$, PO_3H_2 , NHSO_2CF_3 , $\text{NHSO}_2\text{C}_6\text{F}_5$, SO_3H , CONHNH_2 , $\text{CONHNHSO}_2\text{CF}_3$, CONHOCH_3 , $\text{CONHOC}_2\text{H}_5$, CONHCF_3 , OH , CH_2OH , $\text{C}_2\text{H}_4\text{OH}$, OPO_3H_2 , OSO_3H ,



wherein each of R^{42} , R^{43} and R^{44} is independently selected from H, Cl, CN, NO_2 , CF_3 , C_2F_5 , C_3F_7 , CHF_2 , CH_2F , CO_2CH_3 , $\text{CO}_2\text{C}_2\text{H}_5$, SO_2CH_3 , SO_2CF_3 and $\text{SO}_2\text{C}_6\text{F}_5$; wherein Z is selected from O, S, NR^{45} and CH_2 ; wherein R^{45} is selected from hydrido, CH_3 and $\text{CH}_2\text{C}_6\text{H}_5$; and wherein said acidic moiety may be a heterocyclic acidic group attached at any two adjacent positions of R^3 through R^{11} so as to form a fused ring system with one of the phenyl rings of the biphenyl moiety of Formula I, said biphenyl fused ring system selected from



30 and the esters, amides and salts of said acidic moieties;
or a tautomer thereof or a pharmaceutically-acceptable salt thereof.

A class of compounds of particular interest consists of those compounds of Formula I wherein m is one; wherein R¹ is selected from methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, n-pentyl, isopentyl, neopentyl, phenyl, benzyl, phenethyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cyclopropylmethyl, cyclopropylethyl, cyclobutylmethyl, cyclobutylethyl, cyclopentylmethyl, cyclopentylethyl, cyclohexylmethyl, cyclohexylethyl, cyclohexanoyl, methylbutyl, ethylbutyl, dimethylbutyl, thienylmethyl, thienylethyl, thienylpropyl, cyclopentenylmethyl, cyclopentenylethyl, cyclopentenylpropyl, cyclohexenyl, cyclohexenylmethyl, cyclohexenylethyl, adamantyl, adamantylmethyl, adamantylethyl, phenyl, chlorophenyl, dichlorophenyl, fluorophenyl, difluorophenyl, methoxyphenyl, ethoxyphenyl, methylphenyl, ethylphenyl, propylphenyl, isopropylphenyl, dimethylphenyl, diethylphenyl, benzyl, 1-oxo-2-cyclohexylethyl, benzoyl, 1-oxo-2-phenethyl, 1-oxoethyl, 1-oxopropyl, 1-oxobutyl, 1-oxopentyl, 2-butenyl, 3-butenyl, 2-butyryl, 3-butyryl and 2-hydroxybutyl; wherein R⁰ is selected from hydrido, C₄H₉(n), CH₃CH₂CH=CH, C₃H₇(n), SC₃H₇,

45




50




55 C₂H₅, C₅H₁₁(n), C₆H₁₃(n), SC₄H₉,

55




CH₃CH=CH, CH₃CH₂CH₂CH=CH-, amino, aminomethyl, aminoethyl, aminopropyl, CH₂OH, CH₂OCOCH₃, CH₂Cl, CH₂OCH₃, CH₂OCH(CH₃)₂, CHO, CH₂CO₂H, CH(CH₃)CO₂H, NO₂, Cl,

5




10



15

-CO₂CH₃, -CONH₂, -CONHCH₃, CON(CH₃)₂, -CH₂-NHCO₂C₂H₅,

20



25


-CH₂NHCO₂CH₃, -CH₂NHCO₂C₃H₇, -CH₂NHCO₂CH₂(CH₃)₂, -CH₂NHCO₂C₄H₉, CH₂NHCO₂-adamantyl, -CH₂NHCO₂-(1-naphthyl), -CH₂NHCONHCH₃, -CH₂NHCONHC₂H₅, -CH₂NHCONHC₃H₇, -CH₂NHCONHC₄H₉, -CH₂NHCONHCH(CH₃)₂, -CH₂NHCONH(1-naphthyl), -CH₂NHCONH(1-adamantyl),



30


-CH₂CH₂CH₂CO₂H, -CH₂CH₂F, -CH₂OCONHCH₃, -CH₂OCSNHCH₃, -CH₂NHCSOC₃H₇, -CH₂CH₂CH₂F, -CH₂ONO₂,

35



40

-CH₂SH,




45

Cl, NO₂, CF₃, CH₂OH, Br, F, I, methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, n-pentyl, isopentyl, neopentyl, phenyl, cyclohexyl, cyclohexylmethyl, carboxyl, formyl, 1-oxoethyl, 1-oxopropyl, 1-oxobutyl, 1-oxopentyl, dimethoxymethyl, 1,1-dimethoxypropyl, 1,1-dimethoxypentyl, hydroxyalkyl, halo, 1-oxo-2-phenylethyl, 1-oxo-2-cyclohexylethyl, 1,1-difluoro-2-phenylethyl, monofluoromethyl, 1,1-difluoro-2-cyclohexylethyl, 2-cyclohexylethyl, 1,1-difluoro-3-cyclohexylpropyl, 1,1-dimethoxybutyl, 1,1-difluoroethyl, 1,1-difluoropropyl, 1,1-difluorobutyl, 1,1-difluoropentyl, benzyl, 2-phenylethyl, 1,1-difluoro-3-phenylpropyl, cyclohexylmethyl, cyclohexylethyl, cyclohexanoyl, 1-butenyl, 2-butenyl, 3-butenyl, 1-butenyl, 2-butenyl, 3-butenyl, difluoromethyl, CO₂H, SH, PO₃H₂, SO₃H, CONHNH₂, CONHNHSO₂CF₃, OH,


55

5



10 wherein each of R⁴² and R⁴³ is independently selected from chloro, cyano, nitro, trifluoromethyl, methoxycarbonyl and trifluoromethylsulfonyl; wherein R² is selected from methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, n-pentyl, isopentyl, neopentyl, phenyl, benzyl, phenethyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cyclopropylmethyl, cyclopropylethyl, cyclobutylmethyl, cyclobutylethyl, cyclopentylmethyl, cyclopentylethyl, cyclohexylmethyl, cyclohexylethyl, methylbutyl, ethylbutyl, dimethylbutyl, thienylmethyl, thienylethyl, thienylpropyl, cyclopentenylmethyl, cyclopentenylethyl, cyclopentenylpropyl, cyclohexenyl, cyclohexenylmethyl, cyclohexenylethyl, adamantyl, adamantylmethyl, adamantylethyl, phenyl, chlorophenyl, dichlorophenyl, fluorophenyl, difluorophenyl, methoxyphenyl, ethoxyphenyl, methylphenyl, ethylphenyl, propylphenyl, isopropylphenyl, dimethylphenyl, diethylphenyl, benzyl, propylthio, butylthio and hydroxyalkyl; wherein each of R³ through R¹¹ is hydrido with the proviso that at least one of R⁵, R⁶, R⁸ and R⁹ is an acidic group selected from CO₂H, SH, PO₃H₂, SO₃H, CONHNH₂, CONHNHSO₂CF₃, OH,

25




30

wherein each of R⁴² and R⁴³ is independently selected from chloro, cyano, nitro, trifluoromethyl, methoxycarbonyl and trifluoromethylsulfonyl; or a tautomer thereof or a pharmaceutically-acceptable salt thereof.

A class of compounds of more particular interest consists of those compounds of Formula I wherein m is one; wherein R¹ is selected from methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, 4-methylbutyl, n-pentyl, isopentyl, neopentyl, phenyl, benzyl, phenethyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cyclopropylmethyl, cyclopropylethyl, cyclobutylmethyl, cyclobutylethyl, cyclopentylmethyl, cyclopentylethyl, cyclohexylmethyl, cyclohexylethyl, cyclohexanoyl, methylbutyl, ethylbutyl, dimethylbutyl, thienylmethyl, thienylethyl, thienylpropyl, cyclopentenylmethyl, cyclopentenylethyl, cyclopentenylpropyl, cyclohexenyl, cyclohexenylmethyl, cyclohexenylethyl, adamantyl, adamantylmethyl, adamantylethyl, phenyl, chlorophenyl, dichlorophenyl, fluorophenyl, difluorophenyl, methoxyphenyl, ethoxyphenyl, methylphenyl, ethylphenyl, propylphenyl, isopropylphenyl, dimethylphenyl, diethylphenyl, benzyl, 1-oxo-2-cyclohexylethyl, benzoyl, 1-oxo-2-phenethyl, 1-oxoethyl, 1-oxopropyl, 1-oxobutyl, 1-oxopentyl, 2-butenyl, 3-butenyl, 2-butyryl, 3-butyryl and 2-hydroxybutyl; wherein R⁰ is selected from hydrido, methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, 4-methylbutyl, tert-butyl, n-pentyl, neopentyl, 1-oxoethyl, 1-oxopropyl, 1-oxobutyl, 1-oxopentyl, 1-oxo-2-phenylethyl, 1-oxo-2-cyclohexylethyl, 1,1-difluoro-2-phenylethyl, 1,1-difluoro-2-cyclohexylethyl, 2-cyclohexylethyl, 1,1-difluoro-3-cyclohexylpropyl, fluoro, chloro, monofluoromethyl, difluoromethyl, trifluoromethyl, formyl, carboxyl, dimethoxymethyl, 1,1-dimethoxybutyl, 1,1-difluoroethyl, 1,1-difluoropropyl, 1,1-difluorobutyl, 1,1-difluoropentyl, benzyl, 2-phenylethyl, 1,1-difluoro-3-phenylpropyl, cyclohexylmethyl, cyclohexylethyl, cyclohexanoyl, 1-butenyl, 2-butenyl, 3-butenyl, 1-butyryl, 2-butyryl, 3-butyryl, propylthio, butylthio, CO₂H, SH, PO₃H₂, SO₃H, CONHNH₂, CONHNHSO₂CF₃, OH,


55

5



- 10 wherein each of R⁴² and R⁴³ is independently selected from chloro, cyano, nitro, trifluoromethyl, methoxycarbonyl and trifluoromethylsulfonyl; wherein R² is selected from methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, n-pentyl, isopentyl, neopentyl, phenyl, benzyl, phenethyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cyclopropylmethyl, cyclopropylethyl, cyclobutylmethyl, cyclobutylethyl, cyclopentylmethyl, cyclopentylethyl, cyclohexylmethyl, cyclohexylethyl, methylbutyl, ethylbutyl, dimethylbutyl, thienylmethyl, thienylethyl, thienylpropyl, cyclopentenylmethyl, cyclopentenylethyl, cyclopentenylpropyl, cyclohexenyl, cyclohexenylmethyl, cyclohexenylethyl, adamantyl, adamantylmethyl, adamantylethyl, phenyl, chlorophenyl, dichlorophenyl, fluorophenyl, difluorophenyl, methoxyphenyl, ethoxyphenyl, methylphenyl, ethylphenyl, propylphenyl, isopropylphenyl, dimethylphenyl, diethylphenyl, benzyl, propylthio, butylthio and hydroxyalkyl; wherein each of R³ through R¹¹ is hydrido with the proviso that at least one of R⁵, R⁶, R⁸ and R⁹ is an acidic group selected from CO₂H, SH, PO₃H₂, SO₃H, CONHNH₂, CONHNHOSO₂CF₃, OH,

25




30

wherein each of R⁴² and R⁴³ is independently selected from chloro, cyano, nitro, trifluoromethyl, methoxycarbonyl and trifluoromethylsulfonyl; or a tautomer thereof or a pharmaceutically-acceptable salt thereof.

- A class of compounds of even more particular interest consists of those compounds of Formula I
 35 wherein m is one; wherein R¹ is selected from methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tertbutyl, 4-methylbutyl, n-pentyl, isopentyl, neopentyl, phenyl, benzyl, phenethyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cyclopropylmethyl, cyclopropylethyl, cyclobutylmethyl, cyclobutylethyl, cyclopentylmethyl, cyclopentylethyl, cyclohexylmethyl, cyclohexylethyl, cyclohexanoyl, methylbutyl, ethylbutyl, dimethylbutyl, thienylmethyl, thienylethyl, thienylpropyl, cyclopentenylmethyl, cyclopentenylethyl, cyclopentenylpropyl, cyclohexenyl, cyclohexenylmethyl, cyclohexenylethyl, adamantyl, adamantylmethyl, adamantylethyl, phenyl, chlorophenyl, dichlorophenyl, fluorophenyl, difluorophenyl, methoxyphenyl, ethoxyphenyl, methylphenyl, ethylphenyl, propylphenyl, isopropylphenyl, dimethylphenyl, diethylphenyl, benzyl, 1-oxo-2-cyclohexylethyl, benzoyl, 1-oxo-2-phenethyl, 1-oxopropyl, 1-oxobutyl, 1-oxopentyl and 2-hydroxybutyl; wherein R⁰ is selected from hydrido, methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, 4-methylbutyl, n-pentyl, 1-oxo-2-phenylethyl, 1-oxo-2-cyclohexylethyl, fluoro, chloro, monofluoromethyl, difluoromethyl, trifluoromethyl, formyl, carboxyl, dimethoxymethyl, 1,1-difluoro-2-phenylethyl, 1,1-difluoro-2-cyclohexylethyl, 2-cyclohexylethyl, 1,1-difluoro-3-cyclohexylpropyl, dimethoxymethyl, 1,1-dimethoxybutyl, 1,1-difluoroethyl, 1,1-difluoropropyl, 1,1-difluorobutyl, 1,1-difluoropentyl, benzyl, 2-phenylethyl, 1,1-difluoro-3-phenylpropyl, cyclohexylmethyl, cyclohexylethyl, cyclohexanoyl, 1-butenyl, 2-butenyl, 3-butenyl, 1-butylnyl, 2-butylnyl, 3-butylnyl, propylthio and butylthio; wherein R² is selected from methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, n-pentyl, isopentyl, neopentyl, phenyl, benzyl, phenethyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cyclopropylmethyl, cyclopropylethyl, cyclobutylmethyl, cyclobutylethyl, cyclopentylmethyl, cyclopentylethyl, cyclohexylmethyl, cyclohexylethyl, methylbutyl, ethylbutyl, dimethylbutyl, thienylmethyl, thienylethyl, thienylpropyl, cyclopentenylmethyl, cyclopentenylethyl, cyclopentenylpropyl, cyclohexenyl, cyclohexenylmethyl, cyclohexenylethyl, adamantyl, adamantylmethyl, adamantylethyl, phenyl, chlorophenyl, dichlorophenyl, fluorophenyl, difluorophenyl, methoxyphenyl, ethoxyphenyl, methylphenyl, ethylphenyl, propylphenyl, isopropylphenyl, dimethylphenyl, diethylphenyl, benzyl, propylthio, butylthio and hydroxyalkyl; wherein each of R³, R⁴, R⁶, R⁷, R⁸, R¹⁰ and R¹¹ is hydrido; wherein


one of R^5 and R^9 is hydrido and the other of R^5 and R^9 is an acidic group selected from COOH , SH , PO_3H_2 , SO_3H , CONHNH_2 , $\text{CONHNHSO}_2\text{CF}_3$, OH ,



10 wherein each of R^{42} and R^{43} is independently selected from chloro, cyano, nitro, trifluoromethyl, methoxycarbonyl and trifluoromethylsulfonyl;

or a tautomer thereof or a pharmaceutically-acceptable salt thereof.

A class of compounds of even more particular interest consists of those compounds of Formula I
 15 wherein m is one; wherein R^1 is selected from methyl, ethyl, n -propyl, isopropyl, n -butyl, sec -butyl, isobutyl, $tert$ -butyl, 4-methylbutyl, n -pentyl, isopentyl, neopentyl, phenyl, benzyl, phenethyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cyclopropylmethyl, cyclopropylethyl, cyclobutylmethyl, cyclobutylethyl, cyclopentylmethyl, cyclopentylethyl, cyclohexylmethyl, cyclohexylethyl, cyclohexanoyl, methylbutyl, ethylbutyl, dimethylbutyl, thienylmethyl, thienylethyl, thienylpropyl, cyclopentenylmethyl, cyclopentenylethyl, cyclopentenylpropyl, cyclohexenyl, cyclohexenylmethyl, cyclohexenylethyl, adamantyl, adamantylmethyl, adamantylethyl, phenyl, chlorophenyl, dichlorophenyl, fluorophenyl, difluorophenyl, methoxyphenyl, ethoxyphenyl, methylphenyl, ethylphenyl, propylphenyl, isopropylphenyl, dimethylphenyl, diethylphenyl, benzyl, 1-oxo-2-cyclohexylethyl, benzoyl, 1-oxo-2-phenethyl, 1-oxopropyl, 1-oxobutyl, 1-oxopentyl and 2-hydroxybutyl;
 20 wherein R^0 is selected from hydrido, methyl, fluoro, chloro, monofluoromethyl, difluoromethyl, trifluoromethyl, formyl, carboxyl and dimethoxymethyl; wherein R^2 is selected from methyl, ethyl, n -propyl, isopropyl, n -butyl, sec -butyl, isobutyl, $tert$ -butyl, n -pentyl, isopentyl, neopentyl, phenyl, benzyl, phenethyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cyclopropylmethyl, cyclopropylethyl, cyclobutylmethyl, cyclobutylethyl, cyclopentylmethyl, cyclopentylethyl, cyclohexylmethyl, cyclohexylethyl, methylbutyl, ethylbutyl, dimethylbutyl, thienylmethyl, thienylethyl, thienylpropyl, cyclopentenylmethyl, cyclopentenylethyl, cyclopentenylpropyl, cyclohexenyl, cyclohexenylmethyl, cyclohexenylethyl, adamantyl, adamantylmethyl, adamantylethyl, phenyl, chlorophenyl, dichlorophenyl, fluorophenyl, difluorophenyl, methoxyphenyl, ethoxyphenyl, methylphenyl, ethylphenyl, propylphenyl, isopropylphenyl, dimethylphenyl, diethylphenyl, benzyl, propylthio, butylthio and hydroxyalkyl; wherein each of R^3 , R^4 , R^6 , R^7 , R^8 , R^{10} and R^{11} is hydrido; with the proviso that at least one of R^5 and R^9 must be selected from COOH , SH , PO_3H_2 , SO_3H , CONHNH_2 , $\text{CONHNHSO}_2\text{CF}_3$, OH ,
 35




wherein each of R^{42} and R^{43} is independently selected from chloro, cyano, nitro, trifluoromethyl, methoxycarbonyl and trifluoromethylsulfonyl;

or a tautomer thereof or a pharmaceutically-acceptable salt thereof.


A class of compounds of even greater particular interest consists of those compounds of Formula I
 50 wherein m is one; wherein R^1 is selected from methyl, ethyl, n -propyl, isopropyl, n -butyl, sec -butyl, isobutyl, $tert$ -butyl, 4-methylbutyl, n -pentyl, isopentyl, neopentyl, phenyl, benzyl, phenethyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cyclopropylmethyl, cyclopropylethyl, cyclobutylmethyl, cyclobutylethyl, cyclopentylmethyl, cyclopentylethyl, cyclohexylmethyl, cyclohexylethyl, cyclohexanoyl, methylbutyl, ethylbutyl, dimethylbutyl, thienylmethyl, thienylethyl, thienylpropyl, cyclopentenylmethyl, cyclopentenylethyl, cyclopentenylpropyl, cyclohexenyl, cyclohexenylmethyl, cyclohexenylethyl, adamantyl, adamantylmethyl, adamantylethyl, phenyl, chlorophenyl, dichlorophenyl, fluorophenyl, difluorophenyl, methoxyphenyl, ethoxyphenyl, methylphenyl, ethylphenyl, propylphenyl, isopropylphenyl, dimethylphenyl, diethylphenyl, benzyl, 1-oxo-2-cyclohexylethyl, benzoyl, 1-oxo-2-phenethyl, 1-oxopropyl, 1-oxobutyl, 1-oxopentyl and 2-hydroxybutyl;
 55 wherein R^0 is selected from hydrido, methyl, fluoro, chloro, monofluoromethyl, difluoromethyl, trifluoromethyl, formyl, carboxyl and dimethoxymethyl; wherein R^2 is selected from methyl, ethyl, n -propyl,

isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, n-pentyl, isopentyl, neopentyl, phenyl, benzyl, phenethyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cyclopropylmethyl, cyclopropylethyl, cyclobutylmethyl, cyclobutylethyl, cyclopentylmethyl, cyclopentylethyl, cyclohexylmethyl, cyclohexylethyl, methylbutyl, ethylbutyl, dimethylbutyl, thienylmethyl, thienylethyl, thienylpropyl, cyclopentenylmethyl, cyclopentenylethyl, cyclopentenylpropyl, cyclohexenyl, cyclohexenylmethyl, cyclohexenylethyl, adamantyl, adamantylmethyl, adamantylethyl, phenyl, chlorophenyl, dichlorophenyl, fluorophenyl, difluorophenyl, methoxyphenyl, ethoxyphenyl, methylphenyl, ethylphenyl, propylphenyl, isopropylphenyl, dimethylphenyl, diethylphenyl, benzyl, propylthio, butylthio and hydroxyalkyl; wherein each of R³, R⁴, R⁵, R⁷, R⁸, R¹⁰ and R¹¹ is hydrido; wherein one of R⁵ and R⁹ is hydrido and the other of R⁵ and R⁹ is an acidic group selected from CO₂H and




or a tautomer thereof or a pharmaceutically-acceptable salt thereof.

Within Formula I there is a sub-class of compounds of high interest as represented by Formula II:




wherein R¹ is selected from methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, 4-methylbutyl, n-pentyl, isopentyl, neopentyl, phenyl, benzyl, phenethyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cyclopropylmethyl, cyclopropylethyl, cyclobutylmethyl, cyclobutylethyl, cyclopentylmethyl, cyclopentylethyl, cyclohexylmethyl, cyclohexylethyl, cyclohexanoyl, methylbutyl, ethylbutyl, dimethylbutyl, thienylmethyl, thienylethyl, thienylpropyl, cyclopentenylmethyl, cyclopentenylethyl, cyclopentenylpropyl, cyclohexenyl, cyclohexenylmethyl, cyclohexenylethyl, adamantyl, adamantylmethyl, adamantylethyl, phenyl, chlorophenyl, dichlorophenyl, fluorophenyl, difluorophenyl, methoxyphenyl, ethoxyphenyl, methylphenyl, ethylphenyl, propylphenyl, isopropylphenyl, dimethylphenyl, diethylphenyl, benzyl, 1-oxo-2-cyclohexylethyl, benzoyl, 1-oxo-2-phenethyl, 1-oxopropyl, 1-oxobutyl, 1-oxopentyl and 2-hydroxybutyl; wherein R⁰ is selected from hydrido, methyl, fluoro, chloro, monofluoromethyl, difluoromethyl, trifluoromethyl, formyl, carboxyl and dimethoxymethyl; wherein R² is selected from methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, n-pentyl, isopentyl, neopentyl, phenyl, benzyl, phenethyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cyclopropylmethyl, cyclopropylethyl, cyclobutylmethyl, cyclobutylethyl, cyclopentylmethyl, cyclopentylethyl, cyclohexylmethyl, cyclohexylethyl, methylbutyl, ethylbutyl, dimethylbutyl, thienylmethyl, thienylethyl, thienylpropyl, cyclopentenylmethyl, cyclopentenylethyl, cyclopentenylpropyl, cyclohexenyl, cyclohexenylmethyl, cyclohexenylethyl, adamantyl, adamantylmethyl, adamantylethyl, phenyl, chlorophenyl, dichlorophenyl, fluorophenyl, difluorophenyl, methoxyphenyl, ethoxyphenyl, methylphenyl, ethylphenyl, propylphenyl, isopropylphenyl, dimethylphenyl, diethylphenyl and benzyl; wherein R⁵ is an acidic group selected from CO₂H and



or a tautomer thereof or a pharmaceutically-acceptable salt thereof.

Within Formula II there is a more preferred sub-class of compounds of high interest wherein R¹ is

selected from methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, 4-methylbutyl, n-pentyl, isopentyl, neopentyl, phenyl, benzyl, phenethyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cyclopropylmethyl, cyclopropylethyl, cyclobutylmethyl, cyclobutylethyl, cyclopentylmethyl, cyclopentylethyl, cyclohexylmethyl, cyclohexylethyl, cyclohexanoyl, methylbutyl, ethylbutyl, dimethylbutyl, thienylmethyl, thienylethyl, thienylpropyl, cyclopentenylmethyl, cyclopentenylethyl, cyclopentenylpropyl, cyclohexenyl, cyclohexenylmethyl, cyclohexenylethyl, adamantyl, adamantylmethyl, adamantylethyl, phenyl, chlorophenyl, dichlorophenyl, fluorophenyl, difluorophenyl, methoxyphenyl, ethoxyphenyl, methylphenyl, ethylphenyl, propylphenyl, isopropylphenyl, dimethylphenyl, diethylphenyl, benzyl, 1-oxo-2-cyclohexylethyl, benzoyl, 1-oxo-2-phenethyl, 1-oxopropyl, 1-oxobutyl, 1-oxopentyl and 2-hydroxybutyl; wherein R⁰ is hydrido; wherein R² is selected from methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, n-pentyl, isopentyl, neopentyl, phenyl, benzyl, phenethyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cyclopropylmethyl, cyclopropylethyl, cyclobutylmethyl, cyclobutylethyl, cyclopentylmethyl, cyclopentylethyl, cyclohexylmethyl, cyclohexylethyl, methylbutyl, ethylbutyl, dimethylbutyl, thienylmethyl, thienylethyl, thienylpropyl, cyclopentenylmethyl, cyclopentenylethyl, cyclopentenylpropyl, cyclohexenyl, cyclohexenylmethyl, cyclohexenylethyl, adamantyl, adamantylmethyl, adamantylethyl, phenyl, chlorophenyl, dichlorophenyl, fluorophenyl, difluorophenyl, methoxyphenyl, ethoxyphenyl, methylphenyl, ethylphenyl, propylphenyl, isopropylphenyl, dimethylphenyl, diethylphenyl and benzyl; wherein R⁵ is an acidic group selected from CO₂H and



or a tautomer thereof or a pharmaceutically-acceptable salt thereof.

A family of specific compounds of particular interest within Formula II consists of compounds and pharmaceutically-acceptable salts thereof as follows:

1-methyl-4-propyl-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
1-ethyl-4-propyl-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
1,4-dipropyl-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
1-isopropyl-4-propyl-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
1-butyl-4-propyl-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
1-secbutyl-4-propyl-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
1-isobutyl-4-propyl-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
1-tertbutyl-4-propyl-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
1-pentyl-4-propyl-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
1-isopentyl-4-propyl-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
1-cyclohexyl-4-propyl-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
1-cyclohexylmethyl-4-propyl-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
1-phenyl-4-propyl-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
1-phenylmethyl-4-propyl-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
1-(2-cyclohexylethyl)-4-propyl-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
1-(2-phenylethyl)-4-propyl-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
1-methyl-4-butyl-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
1-ethyl-4-butyl-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
1-propyl-4-butyl-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
1-isopropyl-4-butyl-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
1,4-dibutyl-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
1-secbutyl-4-butyl-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
1-isobutyl-4-butyl-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
1-tertbutyl-4-butyl-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
1-pentyl-4-butyl-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
1-isopentyl-4-butyl-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
1-cyclohexyl-4-butyl-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;

[illegible]

- 1-propyl-4-butyl-5-methyl-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
- 1-isopropyl-4-butyl-5-methyl-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
- 5 1,4-dibutyl-5-methyl-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
- 1-secbutyl-4-butyl-5-methyl-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
- 1-isobutyl-4-butyl-5-methyl-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
- 10 1-tertbutyl-4-butyl-5-methyl-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
- 1-pentyl-4-butyl-5-methyl-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
- 1-isopentyl-4-butyl-5-methyl-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
- 15 1-cyclohexyl-4-butyl-5-methyl-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
- 1-cyclohexylmethyl-4-butyl-5-methyl-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
- 20 1-phenyl-4-butyl-5-methyl-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
- 1-phenylmethyl-4-butyl-5-methyl-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
- 1-(2-cyclohexylethyl)-4-butyl-5-methyl-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
- 25 1-(2-phenylethyl)-4-butyl-5-methyl-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
- 1,5-dimethyl-4-pentyl-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
- 1-ethyl-4-pentyl-5-methyl-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
- 30 1-propyl-4-pentyl-5-methyl-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
- 1-isopropyl-4-pentyl-5-methyl-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
- 35 1-butyl-4-pentyl-5-methyl-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
- 1-secbutyl-4-pentyl-5-methyl-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
- 1-isobutyl-4-pentyl-5-methyl-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
- 40 1-tertbutyl-4-pentyl-5-methyl-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
- 1-phenyl-4-pentyl-5-methyl-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
- 45 1-isopentyl-4-pentyl-5-methyl-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
- 1-cyclohexyl-4-pentyl-5-methyl-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
- 1-cyclohexylmethyl-4-pentyl-5-methyl-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
- 50 1,4-dipentyl-5-methyl-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
- 1-phenylmethyl-4-pentyl-5-methyl-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
- 1-(2-cyclohexylethyl)-4-pentyl-5-methyl-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
- 55 1-(2-phenylethyl)-4-pentyl-5-methyl-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
- 1-methyl-4-propyl-5-chloro-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-

- one;
- 1-ethyl-4-propyl-5-chloro-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
- 1,4-dipropyl-5-chloro-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
- 1-isopropyl-4-propyl-5-chloro-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
- 5 one;
- 1-butyl-4-propyl-5-chloro-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
- 1-secbutyl-4-propyl-5-chloro-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
- 1-isobutyl-4-propyl-5-chloro-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
- 10 one;
- 1-tertbutyl-4-propyl-5-chloro-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
- 1-pentyl-4-propyl-5-chloro-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
- 15 1-isopentyl-4-propyl-5-chloro-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
- 1-cyclohexyl-4-propyl-5-chloro-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
- 1-cyclohexylmethyl-4-propyl-5-chloro-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
- 20 1-phenyl-4-propyl-5-chloro-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
- 1-phenylmethyl-4-propyl-5-chloro-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
- 25 1-(2-cyclohexylethyl)-4-propyl-5-chloro-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
- 1-(2-phenylethyl)-4-propyl-5-chloro-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
- 1-methyl-4-butyl-5-chloro-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
- 30 one;
- 1-ethyl-4-butyl-5-chloro-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
- 1-propyl-4-butyl-5-chloro-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
- 1-isopropyl-4-butyl-5-chloro-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
- 35 1,4-dibutyl-5-chloro-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
- 1-secbutyl-4-butyl-5-chloro-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
- 1-isobutyl-4-butyl-5-chloro-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
- 40 1-tertbutyl-4-butyl-5-chloro-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
- 1-pentyl-4-butyl-5-chloro-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
- 1-isopentyl-4-butyl-5-chloro-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
- 45 1-cyclohexyl-4-butyl-5-chloro-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
- 1-cyclohexylmethyl-4-butyl-5-chloro-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
- 1-phenyl-4-butyl-5-chloro-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
- 50 one;
- 1-phenylmethyl-4-butyl-5-chloro-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
- 1-(2-cyclohexylethyl)-4-butyl-5-chloro-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
- 1-(2-phenylethyl)-4-butyl-5-chloro-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
- 55 1-methyl-4-pentyl-5-chloro-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;

- 1-ethyl-4-pentyl-5-chloro-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
 1-propyl-4-pentyl-5-chloro-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
 1-isopropyl-4-pentyl-5-chloro-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
 5 1-butyl-4-pentyl-5-chloro-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
 1-secbutyl-4-pentyl-5-chloro-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
 1-isobutyl-4-pentyl-5-chloro-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
 10 1-tertbutyl-4-pentyl-5-chloro-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
 1,4-dipentyl-5-chloro-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
 1-isopentyl-4-pentyl-5-chloro-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
 15 1-cyclohexyl-4-pentyl-5-chloro-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
 1-cyclohexylmethyl-4-pentyl-5-chloro-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
 20 1-phenyl-4-pentyl-5-chloro-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
 1-phenylmethyl-4-pentyl-5-chloro-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
 1-(2-cyclohexylethyl)-4-pentyl-5-chloro-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one; and
 25 1-(2-phenylethyl)-4-pentyl-5-chloro-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one.

A family of specific compounds of more particular interest within Formula II consists of compounds and pharmaceutically-acceptable salts thereof as follows:

- 30 1-propyl-4-butyl-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
 1,4-dibutyl-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
 1-pentyl-4-butyl-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
 1-isopentyl-4-butyl-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
 1-cyclohexylmethyl-4-butyl-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
 35 1-phenylmethyl-4-butyl-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
 1-(2-cyclohexylethyl)-4-butyl-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
 1-(2-phenylethyl)-4-butyl-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
 40 1-propyl-4-butyl-5-methyl-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
 1,4-dibutyl-5-methyl-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
 1-pentyl-4-butyl-5-methyl-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
 45 1-isopentyl-4-butyl-5-methyl-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
 1-cyclohexylmethyl-4-butyl-5-methyl-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
 1-phenylmethyl-4-butyl-5-methyl-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
 50 1-(2-cyclohexylethyl)-4-butyl-5-methyl-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
 1-(2-phenylethyl)-4-butyl-5-methyl-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
 55 1-propyl-4-butyl-5-chloro-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
 1,4-dibutyl-5-chloro-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
 1-pentyl-4-butyl-5-chloro-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
 1-isopentyl-4-butyl-5-chloro-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-

one;


1-cyclohexylmethyl-4-butyl-5-chloro-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;

1-phenylmethyl-4-butyl-5-chloro-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;


1-(2-cyclohexylethyl)-4-butyl-5-chloro-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one; and

1-(2-phenylethyl)-4-butyl-5-chloro-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one.

Within Formula I there is second sub-class of compounds of high interest as represented by Formula III:



wherein R¹ is selected from methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, 4-methylbutyl, n-pentyl, neopentyl, phenyl, benzyl, phenethyl, cyclohexyl, cyclohexylmethyl, cyclohexylethyl, cyclohexanoyl, 1-oxo-2-cyclohexylethyl, benzoyl, 1-oxo-2-phenethyl, 1-oxopropyl, 1-oxobutyl, 1-oxopentyl and 2-hydroxybutyl; wherein R⁰ is selected from hydrido, methyl, fluoro, chloro, monofluoromethyl, difluoromethyl, trifluoromethyl, formyl, carboxyl and dimethoxymethyl; wherein R² is selected from methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, n-pentyl, isopentyl, neopentyl, phenyl, benzyl, phenethyl, cyclohexyl, cyclohexylmethyl and cyclohexylethyl; wherein R⁵ is an acidic group selected from CO₂H and



or a tautomer thereof or a pharmaceutically-acceptable salt thereof.

A family of specific compounds of particular interest within Formula III consists of compounds and pharmaceutically-acceptable salts thereof as follows:

1-methyl-4-propyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
 1-ethyl-4-propyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
 1,4-dipropyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
 1-isopropyl-4-propyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
 1-butyl-4-propyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
 1-secbutyl-4-propyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
 1-isobutyl-4-propyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
 1-tertbutyl-4-propyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
 1-pentyl-4-propyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
 1-isopentyl-4-propyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
 1-cyclohexyl-4-propyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
 1-cyclohexylmethyl-4-propyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
 1-phenyl-4-propyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
 1-phenylmethyl-4-propyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
 1-(2-cyclohexylethyl)-4-propyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
 1-(2-phenylethyl)-4-propyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;

[illegible]

- imidazol-2-one;
 1-phenyl-4-propyl-5-methyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
 1-phenylmethyl-4-propyl-5-methyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
 5 1-(2-cyclohexylethyl)-4-propyl-5-methyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
 1-(2-phenylethyl)-4-propyl-5-methyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
 10 1,5-dimethyl-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
 1-ethyl-4-butyl-5-methyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
 1-propyl-4-butyl-5-methyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
 1-isopropyl-4-butyl-5-methyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
 15 1,4-dibutyl-5-methyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
 1-secbutyl-4-butyl-5-methyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
 1-isobutyl-4-butyl-5-methyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
 20 1-tertbutyl-4-butyl-5-methyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
 1-pentyl-4-butyl-5-methyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
 25 1-isopentyl-4-butyl-5-methyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
 1-cyclohexyl-4-butyl-5-methyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
 1-cyclohexylmethyl-4-butyl-5-methyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
 30 1-phenyl-4-butyl-5-methyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
 1-phenylmethyl-4-butyl-5-methyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
 35 1-(2-cyclohexylethyl)-4-butyl-5-methyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
 1-(2-phenylethyl)-4-butyl-5-methyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
 1-methyl-4-pentyl-5-methyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
 40 1-ethyl-4-pentyl-5-methyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
 1-propyl-4-pentyl-5-methyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
 45 1-isopropyl-4-pentyl-5-methyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
 1-butyl-4-pentyl-5-methyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
 1-secbutyl-4-pentyl-5-methyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
 50 1-isobutyl-4-pentyl-5-methyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
 1-tertbutyl-4-pentyl-5-methyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
 55 1,4-dipentyl-5-methyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
 1-isopentyl-4-pentyl-5-methyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
 1-cyclohexyl-4-pentyl-5-methyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-

- 2-one;
 1-cyclohexylmethyl-4-pentyl-5-methyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
 1-phenyl-4-pentyl-5-methyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
 5 1-phenylmethyl-4-pentyl-5-methyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
 1-(2-cyclohexylethyl)-4-pentyl-5-methyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
 10 1-(2-phenylethyl)-4-pentyl-5-methyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
 1-methyl-4-propyl-5-chloro-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
 1-ethyl-4-propyl-5-chloro-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
 15 1,4-dipropyl-5-chloro-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
 1-isopropyl-4-propyl-5-chloro-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
 1-butyl-4-propyl-5-chloro-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
 1-secbutyl-4-propyl-5-chloro-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
 20 1-isobutyl-4-propyl-5-chloro-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
 1-tertbutyl-4-propyl-5-chloro-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
 25 1-pentyl-4-propyl-5-chloro-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
 1-isopentyl-4-propyl-5-chloro-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
 1-cyclohexyl-4-propyl-5-chloro-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
 30 1-cyclohexylmethyl-4-propyl-5-chloro-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
 1-phenyl-4-propyl-5-chloro-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
 35 1-phenylmethyl-4-propyl-5-chloro-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
 1-(2-cyclohexylethyl)-4-propyl-5-chloro-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
 1-(2-phenylethyl)-4-propyl-5-chloro-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
 40 1-methyl-4-butyl-5-chloro-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
 1-ethyl-4-butyl-5-chloro-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
 1-propyl-4-butyl-5-chloro-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
 45 1-isopropyl-4-butyl-5-chloro-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
 1,4-dibutyl-5-chloro-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
 1-secbutyl-4-butyl-5-chloro-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
 50 1-isobutyl-4-butyl-5-chloro-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
 1-tertbutyl-4-butyl-5-chloro-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
 1-pentyl-4-butyl-5-chloro-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
 55 1-isopentyl-4-butyl-5-chloro-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
 1-cyclohexyl-4-butyl-5-chloro-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;

- 1-cyclohexylmethyl-4-butyl-5-chloro-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
 1-phenyl-4-butyl-5-chloro-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
 5 1-phenylmethyl-4-butyl-5-chloro-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
 1-(2-cyclohexylethyl)-4-butyl-5-chloro-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
 1-(2-phenylethyl)-4-butyl-5-chloro-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
 10 1-methyl-4-pentyl-5-chloro-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
 1-ethyl-4-pentyl-5-chloro-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
 1-propyl-4-pentyl-5-chloro-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
 15 1-isopropyl-4-pentyl-5-chloro-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
 1-butyl-4-pentyl-5-chloro-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
 1-secbutyl-4-pentyl-5-chloro-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
 20 1-isobutyl-4-pentyl-5-chloro-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
 1-tertbutyl-4-pentyl-5-chloro-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
 1,4-dipentyl-5-chloro-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
 25 1-isopentyl-4-pentyl-5-chloro-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
 1-cyclohexyl-4-pentyl-5-chloro-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
 1-cyclohexylmethyl-4-pentyl-5-chloro-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
 30 1-phenyl-4-pentyl-5-chloro-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
 1-phenylmethyl-4-pentyl-5-chloro-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
 35 1-(2-cyclohexylethyl)-4-pentyl-5-chloro-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one; and
 1-(2-phenylethyl)-4-pentyl-5-chloro-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one.
 40 or a tautomer thereof or a pharmaceutically-acceptable salt thereof.

A family of specific compounds of particular interest within Formula III consists of compounds and pharmaceutically-acceptable salts thereof as follows:

- 1-propyl-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
 1,4-dibutyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
 45 1-pentyl-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
 1-isopentyl-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
 1-cyclohexylmethyl-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
 1-phenylmethyl-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
 50 1-(2-cyclohexylethyl)-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
 1-(2-phenylethyl)-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
 1-propyl-4-butyl-5-methyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
 55 1,4-dibutyl-5-methyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
 1-pentyl-4-butyl-5-methyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
 1-isopentyl-4-butyl-5-methyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-

one;

1-cyclohexylmethyl-4-butyl-5-methyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;

1-phenylmethyl-4-butyl-5-methyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;

1-(2-cyclohexylethyl)-4-butyl-5-methyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;

1-(2-phenylethyl)-4-butyl-5-methyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;

1-propyl-4-butyl-5-chloro-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;

1,4-dibutyl-5-chloro-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;

1-pentyl-4-butyl-5-chloro-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;

1-isopentyl-4-butyl-5-chloro-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;


11-cyclohexylmethyl-4-butyl-5-chloro-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;

1-phenylmethyl-4-butyl-5-chloro-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;


1-(2-cyclohexylethyl)-4-butyl-5-chloro-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one; and

1-(2-phenylethyl)-4-butyl-5-chloro-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one.

Within Formula I there is third sub-class of compounds of high interest as represented by Formula IV:



wherein R¹ is selected from methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, 4-methylbutyl, n-pentyl, neopentyl, phenyl, benzyl, phenethyl, cyclohexyl, cyclohexylmethyl, cyclohexylethyl, cyclohexanoyl, 1-oxo-2-cyclohexylethyl, benzoyl, 1-oxo-2-phenethyl, 1-oxopropyl, 1-oxobutyl, 1-oxopentyl and 2-hydroxybutyl; wherein R⁰ is selected from hydrido, methyl, fluoro, chloro, monofluoromethyl, difluoromethyl, trifluoromethyl, formyl, carboxyl and dimethoxymethyl; wherein R² is selected from methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, n-pentyl, isopentyl, neopentyl, phenyl, benzyl, phenethyl, cyclohexyl, cyclohexylmethyl and cyclohexylethyl; wherein R⁵ is an acidic group selected from CO₂H and



or a tautomer thereof or a pharmaceutically-acceptable salt thereof.

A family of specific compounds of particular interest within Formula IV consists of compounds and pharmaceutically-acceptable salts thereof as follows:

1-methyl-4-propyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

1-ethyl-4-propyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

1,4-dipropyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

1-isopropyl-4-propyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

1-butyl-4-propyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

1-secbutyl-4-propyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

[illegible]

- 1-secbutyl-4-propyl-5-methyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-isobutyl-4-propyl-5-methyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 5 1-tertbutyl-4-propyl-5-methyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-pentyl-4-propyl-5-methyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-isopentyl-4-propyl-5-methyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 10 1-cyclohexyl-4-propyl-5-methyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-cyclohexylmethyl-4-propyl-5-methyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 15 1-phenyl-4-propyl-5-methyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-phenylmethyl-4-propyl-5-methyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-(2-cyclohexylethyl)-4-propyl-5-methyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 20 1-(2-phenylethyl)-4-propyl-5-methyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1,5-dimethyl-4-butyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-ethyl-4-butyl-5-methyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 25 1-propyl-4-butyl-5-methyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-isopropyl-4-butyl-5-methyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1,4-dibutyl-5-methyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 30 1-secbutyl-4-butyl-5-methyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-isobutyl-4-butyl-5-methyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-tertbutyl-4-butyl-5-methyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 35 1-pentyl-4-butyl-5-methyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-isopentyl-4-butyl-5-methyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 40 1-cyclohexyl-4-butyl-5-methyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-cyclohexylmethyl-4-butyl-5-methyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-phenyl-4-butyl-5-methyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 45 1-phenylmethyl-4-butyl-5-methyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-(2-cyclohexylethyl)-4-butyl-5-methyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 50 1-(2-phenylethyl)-4-butyl-5-methyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1,5-dimethyl-4-pentyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-ethyl-4-pentyl-5-methyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 55 1-propyl-4-pentyl-5-methyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-isopropyl-4-pentyl-5-methyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

- 1-butyl-4-pentyl-5-methyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
- 1-secbutyl-4-pentyl-5-methyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
- 5 1-isobutyl-4-pentyl-5-methyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
- 1-tertbutyl-4-pentyl-5-methyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
- 1,4-dipentyl-5-methyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
- 10 1-isopentyl-4-pentyl-5-methyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
- 1-cyclohexyl-4-pentyl-5-methyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
- 1-cyclohexylmethyl-4-pentyl-5-methyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
- 15 1-phenyl-4-pentyl-5-methyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
- 1-phenylmethyl-4-pentyl-5-methyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
- 20 1-(2-cyclohexylethyl)-4-pentyl-5-methyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
- 1-(2-phenylethyl)-4-pentyl-5-methyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
- 1-methyl-4-propyl-5-chloro-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
- 25 1-ethyl-4-propyl-5-chloro-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
- 1,4-dipropyl-5-chloro-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
- 1-isopropyl-4-propyl-5-chloro-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
- 30 1-butyl-4-propyl-5-chloro-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
- 1-secbutyl-4-propyl-5-chloro-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
- 1-isobutyl-4-propyl-5-chloro-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
- 35 1-tertbutyl-4-propyl-5-chloro-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
- 1-pentyl-4-propyl-5-chloro-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
- 1-isopentyl-4-propyl-5-chloro-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
- 40 1-cyclohexyl-4-propyl-5-chloro-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
- 1-cyclohexylmethyl-4-propyl-5-chloro-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
- 45 1-phenyl-4-propyl-5-chloro-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
- 1-phenylmethyl-4-propyl-5-chloro-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
- 1-(2-cyclohexylethyl)-4-propyl-5-chloro-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
- 50 1-(2-phenylethyl)-4-propyl-5-chloro-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
- 1-methyl-4-butyl-5-chloro-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
- 55 1-ethyl-4-butyl-5-chloro-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
- 1-propyl-4-butyl-5-chloro-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
- 1-isopropyl-4-butyl-5-chloro-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

- 1,4-dibutyl-5-chloro-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-secbutyl-4-butyl-5-chloro-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-isobutyl-4-butyl-5-chloro-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-tertbutyl-4-butyl-5-chloro-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-pentyl-4-butyl-5-chloro-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-isopentyl-4-butyl-5-chloro-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-cyclohexyl-4-butyl-5-chloro-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-cyclohexylmethyl-4-butyl-5-chloro-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-phenyl-4-butyl-5-chloro-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-phenylmethyl-4-butyl-5-chloro-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-(2-cyclohexylethyl)-4-butyl-5-chloro-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-(2-phenylethyl)-4-butyl-5-chloro-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-methyl-4-pentyl-5-chloro-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-ethyl-4-pentyl-5-chloro-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-propyl-4-pentyl-5-chloro-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-isopropyl-4-pentyl-5-chloro-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-butyl-4-pentyl-5-chloro-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-secbutyl-4-pentyl-5-chloro-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-isobutyl-4-pentyl-5-chloro-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-tertbutyl-4-pentyl-5-chloro-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1,4-dipentyl-5-chloro-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-isopentyl-4-pentyl-5-chloro-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-cyclohexyl-4-pentyl-5-chloro-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-cyclohexylmethyl-4-pentyl-5-chloro-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-phenyl-4-pentyl-5-chloro-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-phenylmethyl-4-pentyl-5-chloro-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-(2-cyclohexylethyl)-4-pentyl-5-chloro-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one; and
 1-(2-phenylethyl)-4-pentyl-5-chloro-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one.

A family of compounds of more particular interest within Formula IV consists of compounds and pharmaceutically-acceptable salts thereof as follows:

- 1-propyl-4-butyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1,4-dibutyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-pentyl-4-butyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-isopentyl-4-butyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-cyclohexylmethyl-4-butyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-

one;

1-phenylmethyl-4-butyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
1-(2-cyclohexylethyl)-4-butyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

5 1-(2-phenylethyl)-4-butyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
1-propyl-4-butyl-5-methyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

1,4-dibutyl-5-methyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

1-pentyl-4-butyl-5-methyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

1-isopentyl-4-butyl-5-methyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

1-cyclohexylmethyl-4-butyl-5-methyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

15 1-phenylmethyl-4-butyl-5-methyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

1-(2-cyclohexylethyl)-4-butyl-5-methyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

1-(2-phenylethyl)-4-butyl-5-methyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

1-propyl-4-butyl-5-chloro-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

1,4-dibutyl-5-chloro-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

1-pentyl-4-butyl-5-chloro-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

1-isopentyl-4-butyl-5-chloro-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

1-cyclohexylmethyl-4-butyl-5-chloro-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;


1-phenylmethyl-4-butyl-5-chloro-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

1-(2-cyclohexylethyl)-4-butyl-5-chloro-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one; and

1-(2-phenylethyl)-4-butyl-5-chloro-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one.


Within Formula I there is fourth a sub-class of compounds of high interest as represented by Formula

V:



45 wherein R¹ is selected from methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, 4-methylbutyl, n-pentyl, neopentyl, phenyl, benzyl, phenethyl, cyclohexyl, cyclohexylmethyl, cyclohexylethyl, cyclohexanoyl, 1-oxo-2-cyclohexylethyl, benzoyl, 1-oxo-2-phenethyl, 1-oxopropyl, 1-oxobutyl, 1-oxopentyl and 2-hydroxybutyl; wherein R⁰ is selected from hydrido, methyl, fluoro, chloro, monofluoromethyl, difluoromethyl, trifluoromethyl, formyl, carboxyl and dimethoxymethyl; wherein R² is selected from methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, n-pentyl, isopentyl, neopentyl, phenyl, benzyl, phenethyl, cyclohexyl, cyclohexylmethyl and cyclohexylethyl; wherein R⁵ is an acidic group selected from CO₂H and

5



or a tautomer thereof or a pharmaceutically-acceptable salt thereof.

10 A family of specific compounds of particular interest within Formula V consists of compounds and pharmaceutically-acceptable salts thereof as follows:

- 1-methyl-4-propyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
- 1-ethyl-4-propyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
- 1,4-dipropyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
- 15 1-isopropyl-4-propyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
- 1-butyl-4-propyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
- 1-secbutyl-4-propyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
- 1-isobutyl-4-propyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
- 1-tertbutyl-4-propyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
- 20 1-pentyl-4-propyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
- 1-isopentyl-4-propyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
- 1-cyclohexyl-4-propyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
- 1-cyclohexylmethyl-4-propyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
- 25 1-phenyl-4-propyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
- 1-phenylmethyl-4-propyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
- 1-(2-cyclohexylethyl)-4-propyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
- 1-(2-phenylethyl)-4-propyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
- 30 1-methyl-4-butyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
- 1-ethyl-4-butyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
- 1-propyl-4-butyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
- 1-isopropyl-4-butyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
- 35 1,4-dibutyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
- 1-secbutyl-4-butyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
- 1-isobutyl-4-butyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
- 1-tertbutyl-4-butyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
- 1-pentyl-4-butyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
- 40 1-isopentyl-4-butyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
- 1-cyclohexyl-4-butyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
- 1-cyclohexylmethyl-4-butyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
- 1-phenyl-4-butyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
- 45 1-phenylmethyl-4-butyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
- 1-(2-cyclohexylethyl)-4-butyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
- 1-(2-phenylethyl)-4-butyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
- 1-methyl-4-pentyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
- 50 1-ethyl-4-pentyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
- 1-propyl-4-pentyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
- 1-isopropyl-4-pentyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
- 1-butyl-4-pentyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
- 1-secbutyl-4-pentyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
- 55 1-isobutyl-4-pentyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
- 1-tertbutyl-4-pentyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
- 1,4-dipentyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
- 1-isopentyl-4-pentyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

- 1-cyclohexyl-4-pentyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-cyclohexylmethyl-4-pentyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-phenyl-4-pentyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 5 1-phenylmethyl-4-pentyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-(2-cyclohexylethyl)-4-pentyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-(2-phenylethyl)-4-pentyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 10 1-methyl-4-propyl-5-methyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-ethyl-4-propyl-5-methyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1,4-dipropyl-5-methyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 15 1-isopropyl-4-propyl-5-methyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-butyl-4-propyl-5-methyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-secbutyl-4-propyl-5-methyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 20 1-isobutyl-4-propyl-5-methyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-tertbutyl-4-propyl-5-methyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 25 1-pentyl-4-propyl-5-methyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-isopentyl-4-propyl-5-methyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-cyclohexyl-4-propyl-5-methyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 30 1-cyclohexylmethyl-4-propyl-5-methyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-phenyl-4-propyl-5-methyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 35 1-phenylmethyl-4-propyl-5-methyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-(2-cyclohexylethyl)-4-propyl-5-methyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-(2-phenylethyl)-4-propyl-5-methyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 40 1,5-dimethyl-4-butyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-ethyl-4-butyl-5-methyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-propyl-4-butyl-5-methyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 45 1-isopropyl-4-butyl-5-methyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1,4-dibutyl-5-methyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-secbutyl-4-butyl-5-methyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 50 1-isobutyl-4-butyl-5-methyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-tertbutyl-4-butyl-5-methyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-pentyl-4-butyl-5-methyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 55 1-isopentyl-4-butyl-5-methyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-cyclohexyl-4-butyl-5-methyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-

- 2-one;
 1-cyclohexylmethyl-4-butyl-5-methyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-phenyl-4-butyl-5-methyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 5 1-phenylmethyl-4-butyl-5-methyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-(2-cyclohexylethyl)-4-butyl-5-methyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 10 1-(2-phenylethyl)-4-butyl-5-methyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-methyl-4-pentyl-5-methyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-ethyl-4-pentyl-5-methyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 15 1-propyl-4-pentyl-5-methyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-isopropyl-4-pentyl-5-methyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 20 1-butyl-4-pentyl-5-methyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-secbutyl-4-pentyl-5-methyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-isobutyl-4-pentyl-5-methyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 25 1-tertbutyl-4-pentyl-5-methyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1,4-dipentyl-5-methyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-isopentyl-4-pentyl-5-methyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 30 1-cyclohexyl-4-pentyl-5-methyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-cyclohexylmethyl-4-pentyl-5-methyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 35 1-phenyl-4-pentyl-5-methyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-phenylmethyl-4-pentyl-5-methyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-(2-cyclohexylethyl)-4-pentyl-5-methyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 40 1-(2-phenylethyl)-4-pentyl-5-methyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-methyl-4-propyl-5-chloro-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 45 1-ethyl-4-propyl-5-chloro-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1,4-dipropyl-5-chloro-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-isopropyl-4-propyl-5-chloro-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-butyl-4-propyl-5-chloro-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 50 1-secbutyl-4-propyl-5-chloro-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-isobutyl-4-propyl-5-chloro-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-tertbutyl-4-propyl-5-chloro-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 55 1-pentyl-4-propyl-5-chloro-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-isopentyl-4-propyl-5-chloro-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-

- one;
 1-cyclohexyl-4-propyl-5-chloro-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-cyclohexylmethyl-4-propyl-5-chloro-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 5 1-phenyl-4-propyl-5-chloro-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-phenylmethyl-4-propyl-5-chloro-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 10 1-(2-cyclohexylethyl)-4-propyl-5-chloro-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-(2-phenylethyl)-4-propyl-5-chloro-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-methyl-4-butyl-5-chloro-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 15 1-ethyl-4-butyl-5-chloro-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-propyl-4-butyl-5-chloro-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-isopropyl-4-butyl-5-chloro-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 20 1,4-dibutyl-5-chloro-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-secbutyl-4-butyl-5-chloro-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-isobutyl-4-butyl-5-chloro-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 25 1-tertbutyl-4-butyl-5-chloro-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-pentyl-4-butyl-5-chloro-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-isopentyl-4-butyl-5-chloro-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 30 1-cyclohexyl-4-butyl-5-chloro-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-cyclohexylmethyl-4-butyl-5-chloro-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-phenyl-4-butyl-5-chloro-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 35 1-phenylmethyl-4-butyl-5-chloro-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-(2-cyclohexylethyl)-4-butyl-5-chloro-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 40 1-(2-phenylethyl)-4-butyl-5-chloro-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-methyl-4-pentyl-5-chloro-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-ethyl-4-pentyl-5-chloro-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 45 1-propyl-4-pentyl-5-chloro-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-isopropyl-4-pentyl-5-chloro-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-butyl-4-pentyl-5-chloro-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 50 1-secbutyl-4-pentyl-5-chloro-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-isobutyl-4-pentyl-5-chloro-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-tertbutyl-4-pentyl-5-chloro-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 55 1,4-dipentyl-5-chloro-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-isopentyl-4-pentyl-5-chloro-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

1-cyclohexyl-4-pentyl-5-chloro-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

1-cyclohexylmethyl-4-pentyl-5-chloro-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

5 1-phenyl-4-pentyl-5-chloro-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

1-phenylmethyl-4-pentyl-5-chloro-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

1-(2-cyclohexylethyl)-4-pentyl-5-chloro-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one; and

1-(2-phenylethyl)-4-pentyl-5-chloro-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one.

A family of compounds of more particular interest within Formula V consists of compounds and pharmaceutically acceptable salts thereof as follows:

15 1-propyl-4-butyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

1,4-dibutyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

1-pentyl-4-butyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

1-isopentyl-4-butyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

1-cyclohexylmethyl-4-butyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

1-phenylmethyl-4-butyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

1-(2-cyclohexylethyl)-4-butyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

1-(2-phenylethyl)-4-butyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

25 1-propyl-4-butyl-5-methyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

1,4-dibutyl-5-methyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

1-pentyl-4-butyl-5-methyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

30 1-isopentyl-4-butyl-5-methyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

1-cyclohexylmethyl-4-butyl-5-methyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

1-phenylmethyl-4-butyl-5-methyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

35 1-(2-cyclohexylethyl)-4-butyl-5-methyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

1-(2-phenylethyl)-4-butyl-5-methyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

40 1-propyl-4-butyl-5-chloro-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

1,4-dibutyl-5-chloro-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

1-pentyl-4-butyl-5-chloro-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

1-isopentyl-4-butyl-5-chloro-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;


45 1-cyclohexylmethyl-4-butyl-5-chloro-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

1-phenylmethyl-4-butyl-5-chloro-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;


50 1-(2-cyclohexylethyl)-4-butyl-5-chloro-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one; and

1-(2-phenylethyl)-4-butyl-5-chloro-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one.

Within Formula I there is fifth sub-class of compounds of high interest as represented by Formula VI:



wherein R¹ is selected from methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, 4-methylbutyl, n-pentyl, neopentyl, phenyl, benzyl, phenethyl, cyclohexyl, cyclohexylmethyl, cyclohexylethyl, cyclohexanoyl, 1-oxo-2-cyclohexylethyl, benzoyl, 1-oxo-2-phenethyl, 1-oxopropyl, 1-oxobutyl, 1-oxopentyl and 2-hydroxybutyl; wherein R⁰ is selected from hydrido, methyl, fluoro, chloro, monofluoromethyl, difluoromethyl, trifluoromethyl, formyl, carboxyl and dimethoxymethyl; wherein R² is selected from methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, n-pentyl, isopentyl, neopentyl, phenyl, benzyl, phenethyl, cyclohexyl, cyclohexylmethyl and cyclohexylethyl; wherein R⁵ is an acidic group selected from CO₂H and



or a tautomer thereof or a pharmaceutically-acceptable salt thereof.

A family of specific compounds of particular interest within Formula VI consists of compounds and pharmaceutically-acceptable salts thereof as follows:

- 1-methyl-4-propyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
- 1-ethyl-4-propyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
- 1,4-dipropyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
- 1-isopropyl-4-propyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
- 1-butyl-4-propyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
- 1-secbutyl-4-propyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
- 1-isobutyl-4-propyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
- 1-tertbutyl-4-propyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
- 1-pentyl-4-propyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
- 1-isopentyl-4-propyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
- 1-cyclohexyl-4-propyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
- 1-cyclohexylmethyl-4-propyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
- 1-phenyl-4-propyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
- 1-phenylmethyl-4-propyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
- 1-(2-cyclohexylethyl)-4-propyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
- 1-(2-phenylethyl)-4-propyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
- 1-methyl-4-butyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
- 1-ethyl-4-butyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
- 1-propyl-4-butyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
- 1-isopropyl-4-butyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
- 1,4-dibutyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
- 1-secbutyl-4-butyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
- 1-isobutyl-4-butyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
- 1-tertbutyl-4-butyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
- 1-pentyl-4-butyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
- 1-isopentyl-4-butyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
- 1-cyclohexyl-4-butyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

[illegible]

- 1-propyl-4-butyl-5-methyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
- 1-isopropyl-4-butyl-5-methyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
- 5 1,4-dibutyl-5-methyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
- 1-secbutyl-4-butyl-5-methyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
- 1-isobutyl-4-butyl-5-methyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
- 10 1-tertbutyl-4-butyl-5-methyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
- 1-pentyl-4-butyl-5-methyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
- 1-isopentyl-4-butyl-5-methyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
- 15 1-cyclohexyl-4-butyl-5-methyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
- 1-cyclohexylmethyl-4-butyl-5-methyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
- 20 1-phenyl-4-butyl-5-methyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
- 1-phenylmethyl-4-butyl-5-methyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
- 1-(2-cyclohexylethyl)-4-butyl-5-methyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
- 25 1-(2-phenylethyl)-4-butyl-5-methyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
- 1,5-dimethyl-4-pentyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
- 1-ethyl-4-pentyl-5-methyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
- 30 1-propyl-4-pentyl-5-methyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
- 1-isopropyl-4-pentyl-5-methyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
- 35 1-butyl-4-pentyl-5-methyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
- 1-secbutyl-4-pentyl-5-methyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
- 1-isobutyl-4-pentyl-5-methyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
- 40 1-tertbutyl-4-pentyl-5-methyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
- 1,4-dipentyl-5-methyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
- 1-isopentyl-4-pentyl-5-methyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
- 45 1-cyclohexyl-4-pentyl-5-methyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
- 1-cyclohexylmethyl-4-pentyl-5-methyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
- 50 1-phenyl-4-pentyl-5-methyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
- 1-phenylmethyl-4-pentyl-5-methyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
- 1-(2-cyclohexylethyl)-4-pentyl-5-methyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
- 55 1-(2-phenylethyl)-4-pentyl-5-methyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
- 1-methyl-4-propyl-5-chloro-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-

- one;
 1-ethyl-4-propyl-5-chloro-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1,4-dipropyl-5-chloro-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-isopropyl-4-propyl-5-chloro-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 5 one;
 1-butyl-4-propyl-5-chloro-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-secbutyl-4-propyl-5-chloro-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-isobutyl-4-propyl-5-chloro-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 10 one;
 1-tertbutyl-4-propyl-5-chloro-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-pentyl-4-propyl-5-chloro-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 15 one;
 1-isopentyl-4-propyl-5-chloro-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-cyclohexyl-4-propyl-5-chloro-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-cyclohexylmethyl-4-propyl-5-chloro-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 20 one;
 1-phenyl-4-propyl-5-chloro-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-phenylmethyl-4-propyl-5-chloro-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 25 one;
 1-(2-cyclohexylethyl)-4-propyl-5-chloro-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-(2-phenylethyl)-4-propyl-5-chloro-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-methyl-4-butyl-5-chloro-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 30 one;
 1-ethyl-4-butyl-5-chloro-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-propyl-4-butyl-5-chloro-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-isopropyl-4-butyl-5-chloro-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 35 one;
 1,4-dibutyl-5-chloro-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-secbutyl-4-butyl-5-chloro-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-isobutyl-4-butyl-5-chloro-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 40 one;
 1-tertbutyl-4-butyl-5-chloro-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-pentyl-4-butyl-5-chloro-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-isopentyl-4-butyl-5-chloro-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 45 one;
 1-cyclohexyl-4-butyl-5-chloro-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-cyclohexylmethyl-4-butyl-5-chloro-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-phenyl-4-butyl-5-chloro-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 50 one;
 1-phenylmethyl-4-butyl-5-chloro-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-(2-cyclohexylethyl)-4-butyl-5-chloro-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-(2-phenylethyl)-4-butyl-5-chloro-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 55 one;
 1-methyl-4-pentyl-5-chloro-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

- 1-ethyl-4-pentyl-5-chloro-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-propyl-4-pentyl-5-chloro-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-isopropyl-4-pentyl-5-chloro-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-butyl-4-pentyl-5-chloro-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-secbutyl-4-pentyl-5-chloro-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-isobutyl-4-pentyl-5-chloro-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-tertbutyl-4-pentyl-5-chloro-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1,4-dipentyl-5-chloro-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-isopentyl-4-pentyl-5-chloro-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-cyclohexyl-4-pentyl-5-chloro-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-cyclohexylmethyl-4-pentyl-5-chloro-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-phenyl-4-pentyl-5-chloro-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-phenylmethyl-4-pentyl-5-chloro-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-(2-cyclohexylethyl)-4-pentyl-5-chloro-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one; and
 1-(2-phenylethyl)-4-pentyl-5-chloro-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one.

A family of compounds of more particular interest within Formula VI consists of compounds and pharmaceutically-acceptable salts thereof as follows:

- 1-propyl-4-butyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1,4-dibutyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-pentyl-4-butyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-isopentyl-4-butyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-cyclohexylmethyl-4-butyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-phenylmethyl-4-butyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-(2-cyclohexylethyl)-4-butyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-(2-phenylethyl)-4-butyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-propyl-4-butyl-5-methyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1,4-dibutyl-5-methyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-pentyl-4-butyl-5-methyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-isopentyl-4-butyl-5-methyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-cyclohexylmethyl-4-butyl-5-methyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-phenylmethyl-4-butyl-5-methyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-(2-cyclohexylethyl)-4-butyl-5-methyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-(2-phenylethyl)-4-butyl-5-methyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-propyl-4-butyl-5-chloro-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1,4-dibutyl-5-chloro-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-pentyl-4-butyl-5-chloro-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-isopentyl-4-butyl-5-chloro-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

one;


1-cyclohexylmethyl-4-butyl-5-chloro-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

1-phenylmethyl-4-butyl-5-chloro-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;


1-(2-cyclohexylethyl)-4-butyl-5-chloro-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one; and

1-(2-phenylethyl)-4-butyl-5-chloro-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one.

Within Formula I there is sixth sub-class of compounds of high interest as represented by Formula VII:



wherein R¹ is selected from methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, 4-methylbutyl, n-pentyl, neopentyl, phenyl, benzyl, phenethyl, cyclohexyl, cyclohexylmethyl, cyclohexylethyl, cyclohexanoyl, 1-oxo-2-cyclohexylethyl, benzoyl, 1-oxo-2-phenethyl, 1-oxopropyl, 1-oxobutyl, 1-oxopentyl and 2-hydroxybutyl; wherein R⁰ is selected from hydrido, methyl, fluoro, chloro, monofluoromethyl, difluoromethyl, trifluoromethyl, formyl, carboxyl and dimethoxymethyl; wherein R² is selected from methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, n-pentyl, isopentyl, neopentyl, phenyl, benzyl, phenethyl, cyclohexyl, cyclohexylmethyl and cyclohexylethyl; wherein R⁵ is an acidic group selected from CO₂H and



or a tautomer thereof or a pharmaceutically-acceptable salt thereof.

A family of specific compounds of particular interest within Formula VII consists of compounds and pharmaceutically-acceptable salts thereof as follows:

1-methyl-4-propyl-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-ethyl-4-propyl-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1,4-dipropyl-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-isopropyl-4-propyl-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-butyl-4-propyl-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-secbutyl-4-propyl-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-isobutyl-4-propyl-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-tertbutyl-4-propyl-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-pentyl-4-propyl-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-isopentyl-4-propyl-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-cyclohexyl-4-propyl-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-cyclohexylmethyl-4-propyl-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-phenyl-4-propyl-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-phenylmethyl-4-propyl-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-(2-cyclohexylethyl)-4-propyl-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-(2-phenylethyl)-4-propyl-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

[illegible]

- imidazol-2-one;
 1-phenyl-4-propyl-5-methyl-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-phenylmethyl-4-propyl-5-methyl-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 5 1-(2-cyclohexylethyl)-4-propyl-5-methyl-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-(2-phenylethyl)-4-propyl-5-methyl-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 10 1,5-dimethyl-4-butyl-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-ethyl-4-butyl-5-methyl-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-propyl-4-butyl-5-methyl-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-isopropyl-4-butyl-5-methyl-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 15 one;
 1,4-dibutyl-5-methyl-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-secbutyl-4-butyl-5-methyl-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-isobutyl-4-butyl-5-methyl-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 20 one;
 1-tertbutyl-4-butyl-5-methyl-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-pentyl-4-butyl-5-methyl-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 25 1-isopentyl-4-butyl-5-methyl-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-cyclohexyl-4-butyl-5-methyl-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-cyclohexylmethyl-4-butyl-5-methyl-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 30 imidazol-2-one;
 1-phenyl-4-butyl-5-methyl-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-phenylmethyl-4-butyl-5-methyl-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 35 1-(2-cyclohexylethyl)-4-butyl-5-methyl-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-(2-phenylethyl)-4-butyl-5-methyl-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1,5-dimethyl-4-pentyl-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 40 1-ethyl-4-pentyl-5-methyl-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-propyl-4-pentyl-5-methyl-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-isopropyl-4-pentyl-5-methyl-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 45 one;
 1-butyl-4-pentyl-5-methyl-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-secbutyl-4-pentyl-5-methyl-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 50 1-isobutyl-4-pentyl-5-methyl-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-tertbutyl-4-pentyl-5-methyl-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1,4-dipentyl-5-methyl-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 55 1-isopentyl-4-pentyl-5-methyl-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-cyclohexyl-4-pentyl-5-methyl-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

- 1-cyclohexylmethyl-4-pentyl-5-methyl-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-phenyl-4-pentyl-5-methyl-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 5 1-phenylmethyl-4-pentyl-5-methyl-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-(2-cyclohexylethyl)-4-pentyl-5-methyl-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-(2-phenylethyl)-4-pentyl-5-methyl-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 10 1-methyl-4-propyl-5-chloro-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-ethyl-4-propyl-5-chloro-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1,4-dipropyl-5-chloro-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 15 1-isopropyl-4-propyl-5-chloro-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-butyl-4-propyl-5-chloro-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-secbutyl-4-propyl-5-chloro-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 20 1-isobutyl-4-propyl-5-chloro-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-tertbutyl-4-propyl-5-chloro-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-pentyl-4-propyl-5-chloro-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 25 1-isopentyl-4-propyl-5-chloro-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-cyclohexyl-4-propyl-5-chloro-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 30 1-cyclohexylmethyl-4-propyl-5-chloro-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-phenyl-4-propyl-5-chloro-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-phenylmethyl-4-propyl-5-chloro-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 35 1-(2-cyclohexylethyl)-4-propyl-5-chloro-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-(2-phenylethyl)-4-propyl-5-chloro-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 40 1-methyl-4-butyl-5-chloro-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-ethyl-4-butyl-5-chloro-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-propyl-4-butyl-5-chloro-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-isopropyl-4-butyl-5-chloro-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 45 1,4-dibutyl-5-chloro-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-secbutyl-4-butyl-5-chloro-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-isobutyl-4-butyl-5-chloro-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 50 1-tertbutyl-4-butyl-5-chloro-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-pentyl-4-butyl-5-chloro-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-isopentyl-4-butyl-5-chloro-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 55 1-cyclohexyl-4-butyl-5-chloro-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-cyclohexylmethyl-4-butyl-5-chloro-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-

- imidazol-2-one;
 1-phenyl-4-butyl-5-chloro-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-phenylmethyl-4-butyl-5-chloro-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 5 1-(2-cyclohexylethyl)-4-butyl-5-chloro-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-(2-phenylethyl)-4-butyl-5-chloro-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 10 1-methyl-4-pentyl-5-chloro-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-ethyl-4-pentyl-5-chloro-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-propyl-4-pentyl-5-chloro-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 15 1-isopropyl-4-pentyl-5-chloro-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-butyl-4-pentyl-5-chloro-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-secbutyl-4-pentyl-5-chloro-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 20 1-isobutyl-4-pentyl-5-chloro-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-tertbutyl-4-pentyl-5-chloro-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1,4-dipentyl-5-chloro-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 25 1-isopentyl-4-pentyl-5-chloro-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-cyclohexyl-4-pentyl-5-chloro-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-cyclohexylmethyl-4-pentyl-5-chloro-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 30 1-phenyl-4-pentyl-5-chloro-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-phenylmethyl-4-pentyl-5-chloro-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 35 1-(2-cyclohexylethyl)-4-pentyl-5-chloro-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one; and
 1-(2-phenylethyl)-4-pentyl-5-chloro-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one.

A family of compounds of more particular interest within Formula VII consists of compounds and pharmaceutically-acceptable salts as follows:

- 40 1-propyl-4-butyl-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1,4-dibutyl-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-pentyl-4-butyl-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-isopentyl-4-butyl-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 45 1-cyclohexylmethyl-4-butyl-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-phenylmethyl-4-butyl-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-(2-cyclohexylethyl)-4-butyl-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 50 1-(2-phenylethyl)-4-butyl-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-propyl-4-butyl-5-methyl-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1,4-dibutyl-5-methyl-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-pentyl-4-butyl-5-methyl-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 55 1-isopentyl-4-butyl-5-methyl-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-cyclohexylmethyl-4-butyl-5-methyl-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-

imidazol-2-one;

1-phenylmethyl-4-butyl-5-methyl-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

1-(2-cyclohexylethyl)-4-butyl-5-methyl-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

5 1-(2-phenylethyl)-4-butyl-5-methyl-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

1-propyl-4-butyl-5-chloro-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

1,4-dibutyl-5-chloro-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

10 1-pentyl-4-butyl-5-chloro-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

1-isopentyl-4-butyl-5-chloro-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

1-cyclohexylmethyl-4-butyl-5-chloro-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;


15 1-phenylmethyl-4-butyl-5-chloro-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

1-(2-cyclohexylethyl)-4-butyl-5-chloro-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one; and

20 1-(2-phenylethyl)-4-butyl-5-chloro-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one.

The term "hydrido" denotes a single hydrogen atom (H). This hydrido group may be attached, for example, to an oxygen atom to form a hydroxyl group; or, as another example, one hydrido group may be attached to a carbon atom to form a

25



group; or, as another example, two hydrido groups may be attached to a carbon atom to form a $-\text{CH}_2-$ group. Where the term "alkyl" is used, either alone or within other terms such as "haloalkyl" and "hydroxyalkyl", the term "alkyl" embraces linear or branched radicals having one to about twenty carbon atoms or, preferably, one to about twelve carbon atoms. More preferred alkyl radicals are "lower alkyl" radicals having one to about ten carbon atoms. Most preferred are lower alkyl radicals having one to about five carbon atoms. The term "cycloalkyl" embraces cyclic radicals having three to about ten ring carbon atoms, preferably three to about six carbon atoms, such as cyclopropyl, cyclobutyl, cyclopentyl and cyclohexyl. The term "haloalkyl" embraces radicals wherein any one or more of the alkyl carbon atoms is substituted with one or more halo groups, preferably selected from bromo, chloro and fluoro. Specifically embraced by the term "haloalkyl" are monohaloalkyl, dihaloalkyl and polyhaloalkyl groups. A monohaloalkyl group, for example, may have either a bromo, a chloro, or a fluoro atom within the group. Dihalalkyl and polyhaloalkyl groups may be substituted with two or more of the same halo groups, or may have a combination of different halo groups. A dihaloalkyl group, for example, may have two fluoro atoms, such as difluoromethyl and difluorobutyl groups, or two chloro atoms, such as a dichloromethyl group, or one fluoro atom and one chloro atom, such as a fluoro-chloromethyl group. Examples of a polyhaloalkyl are trifluoromethyl, 1,1-difluoroethyl, 2,2,2-trifluoroethyl, perfluoroethyl and 2,2,3,3-tetrafluoropropyl groups. The term "difluoroalkyl" embraces alkyl groups having two fluoro atoms substituted on any one or two of the alkyl group carbon atoms. The terms "alkylol" and "hydroxyalkyl" embrace linear or branched alkyl groups having one to about ten carbon atoms any one of which may be substituted with one or more hydroxyl groups. The term "alkenyl" embraces linear or branched radicals having two to about twenty carbon atoms, preferably three to about ten carbon atoms, and containing at least one carbon-carbon double bond, which carbon-carbon double bond may have either *cis* or *trans* geometry within the alkenyl moiety. The term "alkynyl" embraces linear or branched radicals having two to about twenty carbon atoms, preferably two to about ten carbon atoms, and containing at least one carbon-carbon triple bond. The term "cycloalkenyl" embraces cyclic radicals having three to about ten ring carbon atoms including one or more double bonds involving adjacent ring carbons. The terms "alkoxy" and "alkoxyalkyl" embrace linear or branched oxy-containing radicals each having alkyl portions of one to about ten carbon atoms, such as methoxy group. The term "alkoxyalkyl" also embraces alkyl radicals having two or more alkoxy groups attached to the alkyl radical, that is, to form monoalkoxyalkyl and dialkoxyalkyl groups. The "alkoxy" or "alkoxyalkyl" radicals may be further substituted with one or more halo atoms, such as fluoro, chloro or bromo, to provide haloalkoxy or haloalkoxyalkyl groups. The term "alkylthio" embraces radicals containing a linear or

branched alkyl group, of one to about ten carbon atoms attached to a divalent sulfur atom, such as a methythio group. Preferred aryl groups are those consisting of one, two, or three benzene rings. The term "aryl" embraces aromatic radicals such as phenyl, naphthyl and biphenyl. The term "aralkyl" embraces aryl-substituted alkyl radicals such as benzyl, diphenylmethyl, triphenylmethyl, phenylethyl, phenylbutyl and diphenylethyl. The terms "benzyl" and "phenylmethyl" are interchangeable. The terms "aryloxy" and "arylthio" denote radical respectively, aryl groups having an oxygen or sulfur atom through which the radical is attached to a nucleus, examples of which are phenoxy and phenylthio. The terms "sulfinyl" and "sulfonyl", whether used alone or linked to other terms, denotes respectively divalent radicals SO and SO₂. The term "aralkoxy", alone or within another term, embraces an aryl group attached to an alkoxy group to form, for example, benzyloxy. The term "acyl" whether used alone, or within a term such as acyloxy, denotes a radical provided by the residue after removal of hydroxyl from an organic acid, examples of such radical being acetyl and benzoyl. "Lower alkanoyl" is an example of a more preferred sub-class of acyl. The term "amido" denotes a radical consisting of nitrogen atom attached to a carbonyl group, which radical may be further substituted in the manner described herein. The amido radical can be attached to the nucleus of a compound of the invention through the carbonyl moiety or through the nitrogen atom of the amido radical. The term "alkenylalkyl" denotes a radical having a double-bond unsaturation site between two carbons, and which radical may consist of only two carbons or may be further substituted with alkyl groups which may optionally contain additional double-bond unsaturation. The term "heteroaryl" embraces aromatic ring systems containing one or two hetero atoms selected from oxygen, nitrogen and sulfur in a ring system having five or six ring members, examples of which are thienyl, furanyl, pyridinyl, thiazolyl, pyrimidyl and isoxazolyl. Such heteroaryl may be attached as a substituent through a carbon atom of the heteroaryl ring system, or may be attached through a carbon atom of a moiety substituted on a heteroaryl ring-member carbon atom, for example, through the methylene substituent of imidazolemethyl moiety. Also, such heteroaryl may be attached through a ring nitrogen atom as long as aromaticity of the heteroaryl moiety is preserved after attachment. For any of the foregoing defined radicals, preferred radicals are those containing from one to about ten carbon atoms.

Specific examples of alkyl groups are methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, n-pentyl, isopentyl, methylbutyl, dimethylbutyl and neopentyl. Typical alkenyl and alkynyl groups may have one unsaturated bond, such as an allyl group, or may have a plurality of unsaturated bonds, with such plurality of bonds either adjacent, such as allene-type structures, or in conjugation, or separated by several saturated carbons.


Compounds of Formula I have been found to inhibit the action of angiotensin II in mammals. Angiotensin II is a potent vasoconstrictor and participates in the formation of aldosterone which regulates sodium and water balance in mammals. Thus, compounds of Formula I are therapeutically useful in methods for treating hypertension by administering to a hypertensive patient a therapeutically-effective amount of a compound of Formula I. The phrase "hypertensive patient" means, in this context, a mammalian subject suffering from or afflicted by the effects of hypertension or susceptible to a hypertensive condition if not treated to prevent or control such hypertension.

Also included in the family of compounds of Formula I are isomeric forms including diastereoisomers, regioisomers and the pharmaceutically-acceptable salts thereof. The term "pharmaceutically-acceptable salts" embraces salts commonly used to form alkali metal salts and to form addition salts of free acids or free bases. The nature of the salt is not critical, provided that it is pharmaceutically-acceptable. Suitable pharmaceutically-acceptable acid addition salts of compounds of Formula I may be prepared from an inorganic acid or from an organic acid. Examples of such inorganic acids are hydrochloric, hydrobromic, hydroiodic, nitric, carbonic, sulfuric and phosphoric acid. Appropriate organic acids may be selected from aliphatic, cycloaliphatic, aromatic, araliphatic, heterocyclic, carboxylic and sulfonic classes of organic acids, example of which are formic, acetic, propionic, succinic, glycolic, gluconic, lactic, malic, tartaric, citric, ascorbic, glucuronic, maleic, fumaric, pyruvic, aspartic, glutamic, benzoic, anthranilic, p-hydroxybenzoic, salicylic, phenylacetic, mandelic, embonic (pamoic), methanesulfonic, ethanesulfonic, 2-hydroxyethanesulfonic, pantothenic, benzenesulfonic, toluenesulfonic, sulfanilic, mesylic, cyclohexylaminosulfonic, stearic, algalic, β -hydroxybutyric, malonic, galactaric and galacturonic acid. Suitable pharmaceutically-acceptable base addition salts of compounds of Formula I include metallic salts made from aluminium, calcium, lithium, magnesium, potassium, sodium and zinc or organic salts made from N,N'-dibenzylethylenediamine, chlorprocaine, choline, diethanolamine, ethylenediamine, meglumine (N-methylglucamine) and procaine. All of these salts may be prepared by conventional means from the corresponding compound of Formula I by reacting, for example, the appropriate acid or base with the compound of Formula I.

General Synthetic Procedures

The compounds of the invention can be synthesized according to the following procedures of Schemes I-XXIX, wherein the R substituents are as defined for Formula I, above, except where further noted.

Scheme I



Synthetic Scheme I shows the preparation of the boronic acid **1** where R^5 equals *N*-tertbutyl-*N*-methylcarboxamide. In step 1, benzoic acid is treated with thionyl chloride which is subsequently reacted with *N*-tertbutyl-*N*-methylamine to give the corresponding acid chloride which is subsequently reacted with *N*-tertbutyl-*N*-methylamine to give *N*-tertbutyl-*N*-methylbenzamide. In step 2, the amide is ortho-metalated and subsequently reacted with trimethyl borate. The free boronic acid **1** is produced on hydrolysis.

Scheme II

5

10


15

20

25

30

35



Synthetic Scheme II shows the preparation of the boronic acid **1** where R^5 equals N-triphenylmethyl-1H-tetrazole. In step 1, 2-bromobenzonitrile (Aldrich) is reacted with tributyltin azide to give the corresponding tetrazole. In step 2, the tetrazole is reacted with triphenylmethyl chloride in the presence of triethylamine to give the protected bromophenyltetrazole. In step 3, halogen-metal interchange with n-butyllithium generates the corresponding ortho-lithiated species which is reacted with trimethyl borate. The free boronic acid **1** is produced on hydrolysis.

45


50

55


Scheme III

5


10



15




20



25



30




35



40




45



50 Synthetic Scheme III shows the preparation of N-Boc-amino ketones 2 (or aldehydes when $\text{R}^0 = \text{H}$)
 from the corresponding N-Boc-amino acids 3. In step 1, the amino acid 3 is reacted with isobutyl
 chloroformate in the presence of triethylamine and subsequently with N,O-dimethylhydroxylamine to give
 the corresponding N-methoxy-N-methylamide 4. In step 2, the amide 4 is reacted with an organolithium
 55 reagent $\text{R}^0\text{-Li}$ (or lithium aluminum hydride (LAH) when $\text{R}^0 = \text{H}$) to give the desired ketone 2 (or aldehyde
 when $\text{R}^0 = \text{H}$).

Scheme IV


METHOD A:





Synthetic Scheme IV shows the preparation of imidazol-2-ones 5 from the corresponding amides 4 via Method A. In step 1, the protected amide 4 (prepared in Scheme III) is reacted with trifluoroacetic acid (TFA) to give the TFA salt 6 of the free amine. In step 2, the salt 6 is reacted with the appropriate isocyanate 7 in the presence of triethylamine to give the urea 8. In step 3, the urea 8 is reacted with an organolithium reagent $\text{R}^0\text{-Li}$ (or lithium aluminum hydride (LAH) when $\text{R}^0 = \text{H}$) and subsequently cyclized to the imidazole-2-one 5 on treatment with dilute acid during the work-up procedure.

Scheme V

METHOD B:




HCl, dioxane

 $\text{R}^1-\text{N}=\text{C}=\text{O}$, CHCl_3 

Synthetic Scheme V shows the preparation of imidazol-2-ones 5 from the corresponding N-Boc-protected amino ketones 2 (or aldehydes when $\text{R}^0 = \text{H}$) via Method B. In step 1, the carbonyl compound 2 (prepared in Scheme III) is reacted with anhydrous hydrogen chloride in dioxane to give the HCl salt 9. In step 2, the salt 9 is reacted with the appropriate isocyanate 7 in chloroform to give the imidazol-2-one 5 directly.

Scheme VI

METHOD C:



Synthetic Scheme VI shows the preparation of imidazol-2-ones 5 from the corresponding N-Boc-protected amino ketones 2 (or aldehydes when $R^0 = H$) via Method C. In step 1, the carbonyl compound 2 (prepared in Scheme III) is reacted with 2,2-dimethyl-1,3-propanediol to give the cyclic ketal 10. In step 2, the ketal 10 is reacted with TFA to give the TFA salt 11 of the free amine. In step 3, the salt 11 is reacted with the appropriate isocyanate 7 in the presence of triethylamine to give the urea ketal 12. In step 4, the urea ketal 12 is reacted with 6N hydrochloric acid at 60°C to give the desired imidazol-2-one 5 directly.

Scheme VII

5

10

15

20

25


30

35

40

45


50



Synthetic Scheme VII shows the preparation of 2-bromomethyl-5-bromopyridine (13) and 5-bromo-2-pyridinecarboxaldehyde (14) from 2-picoline (Aldrich). In step 1, 2-picoline is reacted with bromine in the presence of a large excess of aluminum chloride at elevated temperatures to give 5-bromo-2-picoline (15). In step 2a, 15 is reacted with NBS to give the 2-pyridinylmethyl bromide 13. In step 2b, the intermediate 15 is treated with potassium permanganate to give the corresponding picolinic acid 16. In step 3b, the acid 16

is first converted to its N-methoxy-N-methylamide and subsequently reduced with LAH to provide 5-bromo-2-pyridinecarboxaldehyde (14).


Scheme VIII



Synthetic Scheme VIII shows the preparation of 2-bromo-5-bromomethylpyridine (17) and 2-bromo-5-pyridinecarboxaldehyde (18) from 2-amino-5-picoline (Aldrich). In step 1, 2-amino-5-picoline is reacted with bromine in the presence of hydrobromic acid and sodium nitrite at 0°C to give 2-bromo-5-picoline (19). In step 2a, 19 is reacted with NBS to give the 3-pyridinylmethyl bromide 17. In step 2b, the intermediate 19 is

treated with potassium permanganate to give the corresponding nicotinic acid 20. In step 3b, the acid 20 is first converted to its N-methoxy-N-methylamide and subsequently reduced with LAH to provide 2-bromo-5-pyridinecarboxaldehyde (18).


Scheme IX



Synthetic Scheme IX shows the preparation of (4-bromobenzyl)imidazol-2-ones 21 from the TFA salt of the amino amide 6 (prepared in Scheme III). In step 1, the TFA salt 6 is allowed to react with the 4-bromobenzaldehyde in the presence of triethylamine and anhydrous magnesium sulfate to give the imine 22. In step 2, the imine 22 is allowed to react with sodium borohydride to give the substituted benzylamine 23. In step 3, the benzylamine 23 is allowed to react with the appropriate isocyanate 7 to give the substituted benzylurea 24. In step 4, the urea 24 is first allowed to react with an organolithium reagent R^0-Li (or lithium aluminum hydride (LAH) when $\text{R}^0 = \text{H}$) and subsequently with dilute aqueous acid to give the

desired 3-(4-bromobenzyl)imidazol-2-ones.

Scheme X



Synthetic Scheme X shows the preparation of 3-(5-bromo-2-pyridinylmethyl)imidazol-2-ones 25 from the TFA salt of the amino amide 6 (prepared in Scheme III). In step 1, the TFA salt 6 is allowed to react with the 5-bromo-2-pyridinaldehyde 14 (prepared in Scheme VII) in the presence of triethylamine and anhydrous magnesium sulfate to give the imine 26. In step 2, the imine 26 is allowed to react with sodium borohydride to give the substituted benzylamine 27. In step 3, the benzylamine 27 is allowed to react with the appropriate isocyanate 7 to give the substituted benzylurea 28. In step 4, the urea 28 is first allowed to react with an organolithium reagent $\text{R}^0\text{-Li}$ (or lithium aluminum hydride (LAH) when $\text{R}^0 = \text{H}$) and

subsequently with dilute aqueous acid to give the desired 3-(5-bromo-2-pyridinylmethyl)imidazol-2-ones 25.

Scheme XI

5

10

15

20

25

30


35

40

45

50

55



Synthetic Scheme XI shows the preparation of 3-(2-bromo-5-pyridinylmethyl)imidazol-2-ones 29 from the TFA salt of the amino amide 6 (prepared in Scheme III). In step 1, the TFA salt 6 is allowed to react with 2-bromo-5-pyridinaldehyde 18 (prepared in Scheme VIII) in the presence of triethylamine and anhydrous magnesium sulfate to give the imine 30. In step 2, the imine 30 is allowed to react with sodium borohydride to give the substituted benzylamine 31. In step 3, the benzylamine 31 is allowed to react with the appropriate isocyanate 7 to give the substituted benzylurea 32. In step 4, the urea 32 is first allowed to react with an organolithium reagent $\text{R}^0\text{-Li}$ (or lithium aluminum hydride (LAH) when $\text{R}^0 = \text{H}$) and subsequently with dilute aqueous acid to give the desired 3-(2-bromo-5-pyridinylmethyl)imidazol-2-ones 29.

Scheme XII

5

10

15


20

25

30

35

40




Synthetic Scheme XII shows the preparation of 3-(4-bromobenzyl)imidazol-2-ones 21, 3-(5-bromo-2-pyridinylmethyl)imidazol-2-ones 25, and 3-(2-bromo-5-pyridinylmethyl)imidazol-2-ones 29 from the parent imidazol-2-ones 9 (prepared in Scheme IV, Scheme V, or Scheme VI). The imidazol-2-one 9 is first treated with a base, such as potassium t-butoxide, and subsequently with the alkylating agent 4-bromobenzyl bromide, 13 (prepared in Scheme VII), and 17 (prepared in Scheme VIII) to give 3-(4-bromobenzyl)imidazol-2-ones 21, 3-(5-bromo-2-pyridinylmethyl)imidazol-2-ones 25, and 3-(2-bromo-5-pyridinylmethyl)imidazol-2-ones 29, respectively.

50


55

Scheme XIII




Synthetic Scheme XIII shows the preparation of 3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-ones **30** from the boronic acid **1** (prepared in Scheme I and Scheme II) and the bromoimidazol-2-one coupling reagent **25** (prepared in Scheme X and Scheme XII). The boronic acid **1** is reacted with the bromoimidazol-2-one coupling reagent **25** in the presence of a palladium zero catalyst via a Snieckus coupling [see M. J. Sharp and V. Snieckus, *Tetrahedron Lett.*, 5997(1985)] to give the angiotensin II antagonists **30** of this invention.

Scheme XIV



Synthetic Scheme XIV shows the preparation of 3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-ones **31** from the boronic acid **1** (prepared in Scheme I and Scheme II) and the bromoimidazol-2-one coupling reagent **29** (prepared in Scheme XI and Scheme XII). The boronic acid **1** is reacted with the bromoimidazol-2-one coupling reagent **29** in the presence of a palladium zero catalyst via a Snieckus coupling [see M. J. Sharp and V. Snieckus, *Tetrahedron Lett.*, 5997(1985)] to give the angiotensin II antagonists **31** of this invention.

Scheme XV



30 Synthetic Scheme XV shows the preparation of the imidazol-2-one boronic acid coupling reagents 32 from the corresponding 3-(4-bromobenzyl)imidazol-2-ones 21 (prepared in Scheme IX and Scheme XII). Halogen-metal interchange generates the corresponding lithiated species from 21 which is reacted with trimethyl borate. The free imidazol-2-one boronic acid coupling reagents 32 are produced on acid hydrolysis.

35

40

45

50

55

Scheme XVI

5

10

15

20


25

30

35

40

45



50

Synthetic Scheme XVI shows the preparation of the 4-bromopyridine coupling reagent **33** [$\text{R}^5 = \text{CON}(\text{CH}_3)\text{C}(\text{CH}_3)_3$] and the 2-bromopyridine coupling reagent **34** [$\text{R}^5 = \text{CON}(\text{CH}_3)\text{C}(\text{CH}_3)_3$] from nicotinic acid. In step 1, N-tertbutyl-N-methylnicotinamide is prepared from nicotinic acid and N-tertbutyl-N-methylamine. In step 2, ortho-metalation with sec-butyllithium gives a mixture of regioanions which are reacted with trimethylsilyl chloride; subsequent conversion to the corresponding bromides on treatment with bromine in acetic acid and separation of the regioisomers by chromatography provides **33** and **34**.

55

Scheme XVII

5

10

15

20

25


30

35

40

45

50



Synthetic Scheme XVII shows the preparation of the 3-bromopyridine coupling reagent 35 [$\text{R}^5 = \text{CON}(\text{CH}_3)\text{C}(\text{CH}_3)_3$] from isonicotinic acid. In step 1, N-tertbutyl-N-methylisonicotinamide is prepared from isonicotinoyl chloride and N-tertbutyl-N-methylamine. In step 2, reaction with sec-butyllithium gives the ortho-lithiated species which is reacted with trimethylsilyl chloride and subsequently converted to the corresponding bromide 35 on treatment with bromine in acetic acid.

Scheme XVIII

5

10

15

20

25


30

35

40


45

50



Synthetic Scheme XVIII shows the preparation of the 3-bromopyridine coupling reagent 36 [$\text{R}^5 = \text{CON}(\text{CH}_3)\text{C}(\text{CH}_3)_3$] from picolinic acid. In step 1, N-tertbutyl-N-methylpicolinamide is prepared from picolinoyl chloride and N-tertbutyl-N-methylamine. In step 2, reaction with sec-butyllithium gives the ortho-lithiated species which is reacted with trimethylsilyl chloride and subsequently converted to the corresponding bromide 36 on treatment with bromine in acetic acid.


Scheme XIX



Synthetic Scheme XIX shows the preparation of 3-(pyridinylbenzyl)imidazol-2-ones **37**, **38**, **39** and **40** from the common imidazol-2-one boronic acids **32** (Scheme XV) and the corresponding bromo coupling reagents **36** (Scheme XVIII), **33** (Scheme XVI), **35** (Scheme XVII), and **34** (Scheme XVI), respectively. The boronic acids **32** are reacted with the bromo coupling reagents **36**, **33**, **35** and **34** in the presence of a palladium zero catalyst via a Snieckus coupling [see M. J. Sharp and V. Snieckus, *Tetrahedron Lett.*, 5997 (1985)] to give the angiotensin II antagonists **37**, **38**, **39** and **40**, respectively, of this invention.


Scheme XX

5




Synthetic Scheme XX shows the preparation of carboxylic acid analogs 41 and 1H-tetrazole analogs 42 from analogs which have $\text{R}^5 = \text{CON}(\text{CH}_3)\text{C}(\text{CH}_3)_3$. In step 1, the N-tertbutyl-N-methylamide analog 43 is reacted with trifluoroacetic acid at reflux to give the N-methylamide 44. In step 2, the N-methylamide 44 is reacted with sodium nitrite in acetic anhydride/acetic acid at 0°C to give the corresponding N-methyl-N-nitrosoamide 45. In step 3, the N-methyl-N-nitrosoamide 45 is hydrolyzed in base to give the corresponding carboxylic acid angiotensin II antagonists of this invention. In step 4, the acid analog 41 is reacted with oxalyl chloride and subsequently with anhydrous ammonia to give the primary amide 46. In step 5, the amide 46 is reacted with triphenylphosphine in carbon tetrachloride at 50°C to give the corresponding nitrile 47. In step 6, the nitrile 47 is reacted with trimethyltin azide in xylene at reflux to provide the 1H-tetrazole angiotensin II antagonists of this invention.

Scheme XXI



Synthetic Scheme XXI shows the preparation of the organozinc reagent 48 from the appropriate benzoic acid analog 49. In step 1, the analog 49 is brominated with bromine in the presence of a suitable catalyst, e.g., iron, to give the 2-bromo analog 50. In step 2, the 2-bromo analog 50 was converted to the organolithium reagent 51 by reaction with *n*-butyllithium in THF at -78°C , a process known as halogen-metal interchange. Alternatively, the organolithium reagent 51 can be generated directly by the reaction of 49 with an alkyl lithium reagent in the presence or absence of a suitable complexing agent in THF at -78°C , e.g., *sec*-butyllithium/TMEDA (N,N,N',N'-tetramethylethylenediamine). In step 3, the organolithium reagent 51 was treated with anhydrous zinc chloride at -78°C and subsequently allowed to warm to ambient temperature. The organozinc reagent 48 was generated and used in situ.

Scheme XXII



Synthetic Scheme XXII shows the preparation of the 2-pyridinyl alkylating reagent 52 and the 3-pyridinyl alkylating reagent 53 from 15 (Scheme VII) and 19 (Scheme VIII), respectively. In step 1, 15 and 19 were coupled with 1 using Snieckus conditions (Scheme XIII) or 48 using Negishi conditions [see E. Negishi, A. O. King, and N. Okukado, *J. Org. Chem.*, 42, 1821 (1977)] to give 54 and 55, respectively. In step 2, the coupled biaryl compounds 54 and 55 were brominated using NBS/AIBN to give the 2-pyridinyl alkylating

reagent 52 and the 3-pyridinyl alkylating reagent 53, respectively.

Scheme XXIII

5

10

15

20

25





30

35

40

45


50



Synthetic Scheme XXIII shows the preparation of 3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-ones **30** ($R^5 = \text{CN}_4\text{H}$) and 3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-ones **31** ($R^5 = \text{CN}_4\text{H}$) from the parent imidazol-2-ones **9** (prepared in Scheme IV, Scheme V, or Scheme VI). The imidazol-2-one **9** was first treated with a base, such as potassium t-butoxide, and subsequently with the alkylating reagent 52 or 53 (Scheme XXII) to give 3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-


imidazol-2-ones 30 ($R^5 = CH_4H$) and 3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-ones 31 ($R^5 = CN_4H$), respectively.

Scheme XXIV



Synthetic Scheme XXIV shows the preparation of 4-methylphenylboronic acid (56) from 4-bromotoluene. In step 1, the Grignard reagent was generated by the reaction of 4-bromotoluene with metallic magnesium in ether at reflux. In step 2, a THF solution of trimethoxyborane was cooled to $-78^\circ C$ and slowly treated with the Grignard reagent. In step 3, the boronic ester was hydrolyzed with aqueous hydrochloric acid to give 4-methylphenylboronic acid (56).


Scheme XXV



Synthetic Scheme XXV shows the 8-step preparation of the alkylating reagent 2-(4-bromomethyl-phenyl)-3-cyanopyridine (**57**) from 2-amino-3-picoline (**58**) (Aldrich). In step 1, the aminopicoline **58** was converted to the bromopicoline **59** by reaction with bromine, concentrated hydrobromic acid, and sodium nitrite at 0°C . In step 2, the picoline **59** was oxidized with KMnO_4 to give the corresponding carboxylic acid **60**. In step 3, the acid **60** was reduced to the alcohol **61** with borane/THF. In step 4, the alcohol **61** was oxidized to the aldehyde **62** under Swern conditions or by using MnO_2 . In step 5, the aldehyde **62** was reacted with hydroxylamine to give the oxime **63**. In step 6, the oxime **63** was converted to 2-bromo-3-

cyanopyridine (64) with acetic anhydride at reflux. In step 7, the nitrile 64 was coupled with 4-methylphenylboronic acid (56) (Scheme XXIV) using Snieckus conditions (Scheme XIII) to give 3-cyano-2-(4-methylphenyl)pyridine (65). In step 8, 65 was brominated with NBS/AIBN in carbon tetrachloride at reflux to give the desired alkylating reagent 57.

Scheme XXVI



Synthetic Scheme XXVI shows the 6-step preparation of the alkylating reagent 3-(4-bromomethyl-phenyl)-4-cyanopyridine(66) from 4-picoline (67) (Aldrich). In step 1, 4-picoline was brominated with bromine in fuming sulfuric acid at high temperatures to give 3-bromo-4-picoline (68). In step 2, the picoline 68 was oxidized to the corresponding carboxylic acid 69 with KMnO_4 . In step 3, the acid 69 was first
5 converted to its acid chloride with oxalyl chloride and subsequently treated with condensed ammonia to give the amide 70. In step 4, the amide 70 was converted to 3-bromo-4-cyanopyridine (71) by treatment with P_2O_5 at high temperatures. In step 5, the nitrile 71 was coupled with 4-methylphenylboronic acid (56) (Scheme XXIV) using Snieckus conditions (Scheme XIII) to give 4-cyano-3-(4-methylphenyl)pyridine (72). In
10 step 6, 72 was brominated with NBS/AIBN in carbon tetrachloride at reflux to give the desired alkylating reagent 66.

15

20

25

30

35

40

45

50

55

Scheme XXVII

5

10

15

20

25


30

35

40

45


50



Synthetic Scheme XXVII shows the 5-step preparation of the alkylating reagent 4-(4-bromomethylphenyl)-3-cyanopyridine (73) from 4-bromopyridine (74) (Aldrich). In step 1, the ortho-bromo carbanion was generated with LDA in THF at -78°C and reacted with anhydrous DMF to give 4-bromo-3-carboxaldehyde 75. In step 2, the aldehyde 75 was reacted with hydroxylamine to give the oxime 76. In step 3, the oxime 76 was dehydrated with 1,1'-carbonyldiimidazole in methylene chloride at reflux to give 4-bromo-3-cyanopyridine (77). In step 4, the nitrile 77 was coupled with 4-methylphenylboronic acid (56) (Scheme

XXIV) using Snieckus conditions (Scheme XIII) to give 3-cyano-4-(4-methylphenyl)pyridine (78). In step 5, 78 was brominated with NBS/AIBN in carbon tetrachloride at reflux to give the desired alkylating reagent 73.


Scheme XXVIII



Synthetic Scheme XXVIII shows the 4-step preparation of the alkylating reagent 2-cyano-3-(4-bromomethylphenyl)pyridine (79) from 3-bromopyridine (80) (Aldrich). In step 1, the pyridine 80 was reacted with hydrogen peroxide in acetic acid at reflux to give the pyridine N-oxide 81. In step 2, the N-oxide 81

was converted to 3-bromo-2-cyanopyridine (82) by reaction with trimethylsilylcyanide and triethyl anine in acetonitrile at reflux. In step 3, the nitrile 82 was coupled with 4-methylphenylboronic acid (56) (Scheme XXIV) using Snieckus conditions (Scheme XIII) to give 2-cyano-3-(4-methylphenyl) pyridine (83). In step 4, 83 was brominated with NBS/AIBN in carbon tetrachloride at reflux to give the desired alkylating reagent 79.


Scheme XXIX



Synthetic Scheme XXIX shows the preparation of 1,4,5-trisubstituted-1,3-dihydro-3-[[4-(2-cyano-3-pyridinyl)phenyl]methyl]-2H-imidazol-2-ones **37** ($R^5 = \text{CN}$), 1,4,5-trisubstituted-1,3-dihydro-3-[[4-(3-cyano-4-pyridinyl)phenyl]methyl]-2H-imidazol-2-ones **38** ($R^5 = \text{CN}$), 1,4,5-trisubstituted-1,3-dihydro-3-[[4-(4-cyano-3-pyridinyl)phenyl]methyl]-2H-imidazol-2-ones **39** ($R^5 = \text{CN}$), and 1,4,5-trisubstituted-1,3-dihydro-3-[[4-(3-cyano-2-pyridinyl)phenyl]methyl]-2H-imidazol-2-ones **40** ($R^5 = \text{CN}$) from the parent 1,4,5-trisubstituted-1,3-dihydro-2H-imidazol-2-ones **9** (prepared in Scheme IV, Scheme V, or Scheme VI). The imidazol-2-one **9** was first treated with a base, such as potassium t-butoxide, and subsequently with the alkylating reagents **79** (Scheme XXVIII), **73** (Scheme XXVII), **66** (Scheme XXVI), and **57** (Scheme XXV) to give the alkylated products **37** ($R^5 = \text{CN}$), **38** ($R^5 = \text{CN}$), **39** ($R^5 = \text{CN}$), and **40** ($R^5 = \text{CN}$), respectively.

The following Examples contain detailed descriptions of the methods of preparation of compounds of Formula I. These detailed descriptions fall within the scope of, and serve to exemplify, the above described General Synthetic Procedures which form part of the invention. These detailed descriptions are presented for illustrative purposes only and are not intended as a restriction on the scope of the invention. All parts are by weight and temperatures are in degrees Centigrade, unless otherwise indicated.

Example 1



1,4-dibutyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one

Step 1: Preparation of N-t-Boc-L-norleucine-N-methoxy-N-methylamide.

Under nitrogen, a stirred solution of 400 g (1.73 mol) of N-t-Boc-norleucine (BACHEM) and 193 g (1.9 mol) of triethylamine (TEA) in 3000 mL of dichloromethane (DCM) at -50°C was treated with 258 g (1.9 mol) of isobutyl chloroformate. After 30 min, 203 g (2.08 mol) of solid N,O-dimethylhydroxylamine hydrochloride was added followed by 210.5 g (2.08 mol) of TEA at such a rate as to maintain the reaction temperature at -35°C . The reaction was stirred at -15°C for 1 h and then allowed to warm to ambient temperature and stir overnight. The reaction was washed with 1 M citric acid, NaHCO_3 (sat), and brine. The solution was dried (Na_2SO_4) and concentrated in vacuo to give 499.3 g of crude product as a pale green oil: NMR (CDCl_3) δ 0.88 (t, $J = 7$ Hz, 3H), 1.28-1.38 (m, 4H), 1.43 (s, 9H), 1.49-1.57 (m, 1H), 1.63-1.75 (m, 1H), 3.20 (s, 3H), 3.76 (s, 3H), 4.60-4.72 (m, 1H), 5.13 (d, $J = 8$ Hz, 1H).

Step 2A: Preparation of 1,4-dibutyl-1,3-dihydro-2H-imidazol-2-one: Method A.

Under nitrogen, a stirred solution of 449 g of crude N-t-Boc-L-norleucine-N-methoxy-N-methylamide from Step 1 in 1350 mL of methylene chloride at 0°C was treated with 1350 mL of trifluoroacetic acid (TFA). The reaction was allowed to warm to ambient temperature and stir. After 2 h, the reaction was concentrated in vacuo to give the TFA salt of L-norleucine-N-methoxy-N-methylamide as a viscous colorless oil: NMR (CDCl_3) δ 0.88 (t, $J = 7$ Hz, 3H), 1.30-1.40 (m, 4H), 1.88 (t, $J = 7$ Hz, 2H), 3.25 (s, 3H), 3.75 (s, 3H), 4.35-4.46 (m, 1H), 7.55-7.76 (br s, 3H). The TFA salt was cooled to 0°C and treated with aqueous sodium hydroxide until the pH was 10. The resulting solution was continuously extracted with ether for 24 h, continuously extracted with ether/DCM (3:1) for 24 h, and continuously extracted with ether/DCM (1:1) for 24 h.

h. The organic layer was dried (Na_2SO_4) and concentrated *in vacuo* to give 245 g (83% from N-t-Boc-norleucine) as the monohydrate. Under nitrogen, a solution of this material in 100 mL of chloroform at 0 °C was slowly treated with 208 g (2.1 mol) of neat n-butyliisocyanate over 15 min. The reaction was allowed to warm to ambient temperature and stir for 3 h. The reaction was concentrated *in vacuo*; purification by silica gel chromatography (Waters Prep-500A) using ethyl acetate gave 299.5 g as a colorless oil: NMR (CDCl_3) δ 0.83-0.95 (m, 6H), 1.25-1.47 (m, 8H), 1.48-1.60 (m, 1H), 1.63-1.75 (m, 1H), 3.02-3.26 (m, 2H), 3.21 (s, 3H), 3.84 (s, 3H), 4.82-4.93 (m, 1H). Under nitrogen, 299 g (1.09 mol) of this material was dissolved in 600 mL of anhydrous diethyl ether; the solution was cooled to -15 °C and slowly treated with 720 mL (720 mmol) of a 1.0 M solution of lithium aluminum hydride (LAH) in ether. The reaction was allowed to warm to ambient temperature and stir overnight prior to the addition of 50 mL of ethyl acetate. A solution of 285 g (2 mol) of potassium bisulfate in 1500 mL of water was added cautiously and the reaction stirred for 4 h. The reaction mixture was transferred to a separatory funnel and the phases separated. The aqueous phase was extracted 8 times with additional ether. The combined ether extracts were washed 3 times each with 3N hydrochloric acid, saturated sodium bicarbonate, and brine. The ether extracts were then dried (MgSO_4) and concentrated *in vacuo* to give 228 g (68% from N-t-Boc-norleucine) of 1,4-dibutyl-1,3-dihydro-2H-imidazol-2-one as a viscous colorless oil which solidified on storage in the refrigerator: NMR (CDCl_3) δ 0.91 (t, J=7 Hz, 3H), 0.93 (t, J=7 Hz, 3H), 1.28-1.41 (m, 4H), 1.47-1.65 (m, 4H), 2.36 (td, J=7 and 1 Hz, 2H), 3.55 (t, J=7 Hz, 2H), 5.84 (t, J=1 Hz, 1H), 9.78 (br s, 1H).

20 Step 2B: Preparation of 1,4-dibutyl-1,3-dihydro-2H-imidazol-2-one: Method B.

Under nitrogen, a stirred solution of 67.8 g (0.26 mol) of N-t-Boc-norleucine-N-methoxy-N-methylamide from Step 1 in 550 mL of anhydrous diethyl ether at 0 °C was treated with 145 mL (145 mmol) of a 1 M solution of lithium aluminum hydride (LAH) in ether over a 30 min period. The reaction was allowed to stir for an additional 30 min and then was quenched with the addition of 10 mL of ethyl acetate. The reaction was diluted with 1 L of cold water to which 63 g (0.46 mol) of potassium hydrogen sulfate had been added and the mixture stirred vigorously for 15 min. The phases were separated and the aqueous phase extracted 4 times with ether; the extracts were combined, washed 3 times with 3 N hydrochloric acid, once with saturated sodium bicarbonate, and once with brine, dried (Na_2SO_4) and concentrated *in vacuo* to give 47.6 g (84%) of N-t-Boc-L-norleucinal as a colorless waxy solid: NMR (CDCl_3) δ 0.91 (t, J=7 Hz, 3H), 1.34-1.41 (m, 4H), 1.46 (s, 9H), 1.55-1.64 (m, 1H), 1.80-1.95 (m, 1H), 4.16-4.30 (m, 1H), 5.07-5.15 (m, 1H), 9.59 (s, 1H). Under nitrogen, a stirred solution of 10.0 g (46.4 mmol) of this material in 10 mL of dioxane (anhydrous) at 0 °C was treated with 120 mL (480 mmol) of 4 N hydrogen chloride in dioxane over a 10 min period. The reaction was allowed to stir at 0 °C for an additional 20 min after the addition was complete and then concentrated *in vacuo*. The residue was dissolved in 200 mL of chloroform and treated with 77.4 g (0.78 mol) of butyl isocyanate. The mixture was stirred at ambient temperature for 24 h, stirred at 40 °C for 24 h, and concentrated *in vacuo*. Purification of the reddish colored residue by silica gel chromatography (Harrison Chromatotron) using ethyl acetate/2-propanol (95:5) gave 160 mg (1.7% from N-t-Boc-L-norleucinal) of 1,4-dibutyl-1,3-dihydro-2H-imidazol-2-one as an yellowish oil: NMR (CDCl_3) δ 0.91 (t, J=7 Hz, 3H), 0.93 (t, J=7 Hz, 3H), 1.28-1.41 (m, 4H), 1.47-1.65 (m, 4H), 2.36 (td, J=7 and 1 Hz, 2H), 3.55 (t, J=7 Hz, 2H), 5.84 (t, J=1 Hz, 1H), 9.78 (br s, 1H); MS (FAB) m/e (rel intensity) 197 (100), 153 (12), 141 (12), 125 (8), 111 (7).

45 Step 2C: Preparation of 1,4-dibutyl-1,3-dihydro-2H-imidazol-2-one: Method C.

Under nitrogen, a solution of 27.0 g (125 mmol) of N-t-Boc-L-norleucinal from step 2B, 39.1 g (376 mmol) of 2,2-dimethyl-1,3-propanediol, and 1.18 g (6.2 mmol) of p-toluenesulfonic acid monohydrate in 220 mL of benzene was stirred at reflux for 22 h over a Dean-Stark trap. The reaction was cooled, diluted with 300 mL of ethyl acetate, washed sequentially with saturated sodium bicarbonate and brine, dried (Na_2SO_4), and concentrated *in vacuo* to give 38.27 g of crude ketal which was a red oil: NMR (CDCl_3) δ 0.71 (s, 3H), 0.89 (t, J=7 Hz, 3H), 1.16 (s, 3H), 1.20-1.52 (m, 5H), 1.45 (s, 9H), 1.58-1.72 (m, 1H), 3.35-3.46 (m, 2H), 3.56-3.63 (m, 2H), 4.43 (s, 1H), 4.66 (d, J=9 Hz, 1H). Under nitrogen, a 38.2 g sample of the crude ketal was dissolved in 200 mL of methylene chloride; the solution was cooled to 0 °C and treated with 200 mL of trifluoroacetic acid. The reaction was allowed to warm to ambient temperature and stir for 2 h. Concentration *in vacuo* gave the crude TFA salt of the free amino ketal as a red viscous oil: NMR (CDCl_3) δ 0.75 (s, 3H), 0.90 (t, J=7 Hz, 3H), 1.13 (s, 3H), 1.27-1.42 (m, 4H), 1.60-1.82 (m, 2H), 3.27-3.38 (m, 1H), 3.42-3.52 (m, 2H), 3.63-3.72 (m, 2H), 4.58 (d, J=3 Hz, 1H), 7.1-7.5 (br s, 3H). Under nitrogen, the crude TFA salt was redissolved in 100 mL of methylene chloride and treated with 264 g (2.66 mol) of butyl isocyanate. The

reaction was stirred at ambient temperature for 17 h. The reaction was concentrated *in vacuo* to give the crude N-butyl urea of the ketal as a red oil. Purification of a small sample by silica gel chromatography (Harrison Chromatotron) using ethyl acetate/hexane (1:1) gave the pure N-butyl urea as a pale yellow oil: NMR (CDCl₃) δ 0.71 (s, 3H), 0.89 (t, J = 7 Hz, 3H), 0.92 (t, J = 7 Hz, 3H), 1.16 (s, 3H), 1.23-1.54 (m, 9H), 1.61-1.75 (m, 1H), 3.09-3.21 (m, 2H), 3.37-3.46 (m, 2H), 3.56-3.64 (m, 2H), 3.68-3.79 (m, 1H), 4.32-4.47 (br s, 1H), 4.42 (d, J = 3 Hz, 1H). Under nitrogen, a solution of the crude N-butyl urea ketal in 1 L of 1,4-dioxane was treated with 1 L of 6 N hydrochloric acid. The reaction was allowed to stir at ambient temperature for 16 h and then warmed to 60 °C and stirred for an additional 24 h. The reaction was concentrated *in vacuo*; the residue was treated with ethyl acetate and filtered. The ethyl acetate solution was dried (MgSO₄) and concentrated *in vacuo* to give the crude product. Purification by silica gel chromatography (Waters Prep-500A) using methylene chloride/2-propanol (95:5) gave 4.95 g (20% from N-t-Boc-L-norleucinal) of 1,4-dibutyl-1,3-dihydro-2H-imidazol-2-one as a pale yellow oil: NMR (CDCl₃) δ 0.91 (t, J = 7 Hz, 3H), 0.93 (t, J = 7 Hz, 3H), 1.28-1.41 (m, 4H), 1.47-1.65 (m, 4H), 2.36 (td, J = 7 and 1 Hz, 2H), 3.55 (t, J = 7 Hz, 2H), 5.84 (t, J = 1 Hz, 1H), 9.78 (br s, 1H).

Step 3: Preparation of 2-bromo-5-picoline.

A solution of 1500 mL (14 mol) of 48% hydrobromic acid was cooled to 10 °C and 300 g (2.8 mol) of 2-amino-5-picoline (Aldrich) was added slowly. The solution was maintained at or below 0 °C while 450 mL (8.8 mol) of bromine was added dropwise. After the bromine addition was complete, a solution of 500 g (7.3 mol) of sodium nitrite in 1000 mL of water was added slowly over 6 h. The reaction pH was adjusted by the careful addition of 1500 mL (56 mol) of 50% sodium hydroxide at such a rate that the temperature was maintained below 30 °C. The product precipitated from the nearly colorless reaction mixture; filtration gave 450 g (94%) of 2-bromo-5-picoline as a yellow powder: mp 38-40 °C; NMR 7.27 (s, 1H), 7.28 (s, 1H), 7.12 (br s, 1H).

Step 4: Preparation of 2-bromo-5-bromomethylpyridine.

A solution of 296.3 g (1.72 mol) of 2-bromo-5-picoline from step 3 in 6 L of carbon tetrachloride was treated with 306.5 g (1.72 mol) of N-bromosuccinimide (NBS) and 28.3 g (173 mmol) of azobisisobutyronitrile (AIBN). The reaction was stirred at reflux under nitrogen for 3 h, filtered, and concentrated *in vacuo* providing 476 g of crude 2-bromo-5-bromomethylpyridine as a brownish yellow solid (NMR indicates that this material is only 69% monobromomethyl product): NMR (CDCl₃) δ 4.42 (s, 2H), 7.48 (d, J = 9 Hz, 1H), 7.60 (dd, J = 9 and 3 Hz, 1H), 8.37 (d, J = 3 Hz, 1H).

Step 5: Preparation of 1,4-dibutyl-1,3-dihydro-3-[(6-bromo-3-pyridinyl)methyl]-2H-imidazol-2-one.

Under nitrogen, 138 mL (138 mmol) of 1.0 M potassium t-butoxide in THF was added to a stirred solution of 24.5 g (125 mmol) of 1,4-dibutylimidazol-2-one from step 2A in 370 mL of DMF at -30 °C at such a rate to maintain the solution temperature below -20 °C. After 10 min, 50 g [138 mmol (69% purity)] of 2-bromo-5-bromomethylpyridine from step 4 and 2.7 g (18.1 mmol) of sodium iodide were added. The reaction was stirred at 0 °C for 2 h and then allowed to slowly warm to ambient temperature overnight. The reaction was partitioned between ethyl acetate and water; the organic layer was washed twice with brine and the combined aqueous phases were back-extracted with ethyl acetate. The ethyl acetate extractions were combined, dried (MgSO₄), and concentrated *in vacuo*. Purification by silica gel chromatography (Waters Prep-500A) using hexane/ethyl acetate (25:75) gave 33 g (73%) of 1,4-dibutyl-1,3-dihydro-3-[(6-bromo-3-pyridinyl)methyl]-2H-imidazol-2-one as a reddish-brown waxy solid: NMR (CDCl₃) δ 0.89 (t, J = 7 Hz, 3H), 0.95 (t, J = 7 Hz, 3H), 1.25-1.51 (m, 6H), 1.57-1.71 (m, 2H), 2.22 (t, J = 8 Hz, 2H), 3.60 (t, J = 7 Hz, 2H), 4.80 (s, 2H), 5.90 (s, 1H), 7.43 (d, J = 9 Hz, 1H), 7.49 (dd, J = 9 and 2 Hz, 1H), 8.24 (d, J = 2 Hz, 1H).

Step 6: Preparation of 2-(N-triphenylmethyltetrazol-5-yl)phenylboronic acid.


A 64 g (350 mmol) sample of 2-bromobenzonitrile (Aldrich) was dissolved in 650 mL of xylene and treated with 22.75 g (350 mmol) of sodium azide and 95 mL (350 mmol) of tributyltin chloride at reflux for 48 h. The reaction was filtered; the filtrate was treated with 50 mL of anhydrous tetrahydrofuran (THF) and 20 g (550 mmol) of hydrogen chloride. The reaction was stirred for 2 h; filtration gave 59.6 g (76%) of 5-(2-bromophenyl)-1H-tetrazole: mp 178-180 °C; NMR (DMSO-d₆) δ 7.50-7.64 (m, 2H), 7.67-7.74 (m, 1H), 7.83-7.91 (m, 1H). A 41.8 g (187 mmol) sample of this material was dissolved in 650 mL of methylene chloride

and treated with 55.5 g (193 mmol) of triphenylmethyl chloride and 30 mL (220 mmol) of anhydrous triethylamine. The reaction was stirred overnight at reflux, cooled to ambient temperature, washed with water, dried (MgSO_4), and concentrated in *vacuo*. Recrystallization from toluene/hexane gave 80.7 g (92%) of N-triphenylmethyl-5-(2-bromophenyl)-1H-tetrazole: mp 160-162 °C; NMR (CDCl_3) δ 7.14-7.21 (m, 6H), 7.26-7.45 (m, 11H), 7.70 (dd, $J=8$ and 1.5 Hz, 1H), 7.89 (dd, $J=7.5$ and 2 Hz, 1H). A 34.05 g (73.0 mmol) sample of this material was dissolved in 1700 mL of THF under a nitrogen atmosphere and treated with 73 mmol of n-butyllithium in hexane. The reaction was allowed to stir for 17 min and then was treated with 24.9 mL (220 mmol) of trimethyl borate. The reaction was allowed to come to ambient temperature overnight while stirring, quenched with 10 mL of methanol, and concentrated in *vacuo*. The residue was dissolved in 1M NaOH and extracted with toluene to remove any unreacted starting material. The pH was adjusted to 6 with 6M HCl and the product extracted with toluene and dried (MgSO_4). Hexane was added and the solution kept in the freezer overnight. Filtration provided 31.3 g (99%) of 2-(N-triphenylmethyltetrazol-5-yl)-phenylboronic acid: NMR (CDCl_3) δ 7.13-7.21 (m, 7H), 7.33-7.42 (m, 8H), 7.49-7.55 (m, 2H), 8.15-8.19 (m, 1H), 8.21-8.26 (m, 1H).

Step 7: Preparation of 1,4-dibutyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one.

A solution of 21.7 g (50.2 mmol) of 2-(N-triphenylmethyltetrazol-5-yl)phenylboronic acid from step 6 in 80 mL of ethanol and 130 mL of toluene was added to a mixture of 5 g (4 mmol) of tetrakis-(triphenylphosphine) Pd(0), 16.75 g (45.8 mmol) of 1,4-dibutyl-1,3-dihydro-3-[(6-bromo-3-pyridinyl)methyl]-2H-imidazol-2-one from step 5, 225 mL of toluene, 100 mL of 2 M sodium carbonate, and 150 mL of ethanol. The reaction mixture was heated to reflux and vigorously stirred under nitrogen for 14 h. The pH was adjusted to 6 with acetic acid and the reaction concentrated in *vacuo*. The product was purified by silica gel chromatography (Waters Prep-500A) using 2-propanol/methylene chloride (0-50%) followed by reverse phase chromatography (Waters Deltaprep-3000) using acetonitrile/water (30-35:70-65) (0.05% TFA). The pure fractions (by analytical HPLC) were combined, the acetonitrile removed in *vacuo*, and the water extracted 4 times with chloroform. The extracts were combined, dried (MgSO_4), and concentrated in *vacuo*. Recrystallization from ethyl acetate gave 8.5 g (43%) of colorless 1,4-dibutyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one as a colorless solid: mp 170-171 °C; NMR (CDCl_3) δ 0.88 (t, $J=7$ Hz, 3H), 0.96 (t, $J=7$ Hz, 3H), 1.28-1.54 (m, 6H), 1.61-1.72 (m, 2H), 2.28 (t, $J=8$ Hz, 2H), 3.63 (t, $J=7$ Hz, 2H), 4.96 (s, 2H), 5.96 (s, 1H), 7.36 (d, $J=9$ Hz, 1H), 7.47-7.58 (m, 3H), 7.67 (dd, $J=9$ and 2 Hz, 1H), 8.06-8.13 (m, 1H), 8.52 (d, $J=2$ Hz, 1H); MS (FAB) m/e (rel intensity) 432 (100), 404 (18), 237 (14), 209 (92), 180 (82); HRMS. Calc'd for $\text{M}+\text{H}^+$: 432.2512. Found: 432.2554. Anal. Calc'd for $\text{C}_{24}\text{H}_{29}\text{N}_7\text{O}$: C, 66.80; H, 6.77; N, 22.72. Found: C, 66.59; H, 6.85; N, 22.57.

Example 2



1,4-dipropyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one


Step 1: Preparation of 1,4-dipropyl-1,3-dihydro-2H-imidazol-2-one.

Following General Synthetic Schemes III and IV, 1,4-dipropyl-1,3-dihydro-2H-imidazol-2-one was prepared: NMR (CDCl₃) δ 0.92 (t, J = 7 Hz, 3H), 0.93 (t, J = 7 Hz, 3H), 1.57 (m, J = 7 Hz, 2H), 1.67 (m, J = 7 Hz, 2H), 2.34 (td, J = 7 and 1 Hz, 2H), 3.52 (t, J = 7 Hz, 2H), 5.86 (t, J = 1 Hz, 1H), 10.06 (br s, 1H); MS (FAB) m/e (rel intensity) 169 (100), 153 (4), 139 (9), 127 (9); HRMS. Calc'd for M + H: 169.1341. Found 169.1346.

Step 2: Preparation of 1,4-dipropyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one.

Following General Synthetic Schemes XII and XIV or XXIII, the imidazol-2-one from Step 1 was converted to 1,4-dipropyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one as a colorless solid: NMR (CDCl₃) δ 0.94 (t, J = 7 Hz, 3H), 0.95 (t, J = 7 Hz, 3H), 1.53 (m, J = 7 Hz, 3H), 1.71 (m, J = 7 Hz, 2H), 2.27 (td, J = 7 and 1 Hz, 2H), 3.60 (t, J = 7 Hz, 2H), 4.95 (s, 2H), 5.97 (t, J = 1 Hz, 1H), 7.33 (d, J = 8 Hz, 1H), 7.47-7.57 (m, 3H), 7.64 (dd, J = 8 and 2 Hz, 1H), 8.04-8.11 (m, 1H), 8.52 (d, J = 2 Hz, 1H); MS (FAB) m/e (rel intensity) 404 (83), 376 (22), 347 (8), 237 (7), 209 (100), 180 (98); HRMS. Calc'd for M + H: 404.2199. Found: 404.2177.

Example 3



1-propyl-4-isobutyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one


Step 1: Preparation of 1-propyl-4-isobutyl-1,3-dihydro-2H-imidazol-2-one.

Following General Synthetic Schemes III and IV, 1-propyl-4-isobutyl-1,3-dihydro-2H-imidazol-2-one was prepared: NMR (CDCl₃) δ 0.91 (d, J = 7 Hz, 6H), 0.93 (t, J = 7 Hz, 3H), 1.60-1.72 (m, 2H), 1.83 (m, J = 7 Hz, 1H), 2.21 (d, J = 7 Hz, 2H), 3.52 (t, J = 7 Hz, 2H), 5.84-5.87 (m, 1H), 9.78 (br s, 1H); MS (FAB) m/e (rel intensity) 183 (100), 167 (4), 153 (2), 139 (15); HRMS. Calc'd for M + H: 183.1497. Found: 183.1492.

Step 2: Preparation of 1-propyl-4-isobutyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one.

Following General Synthetic Schemes XII and XIV or XXIII, the imidazol-2-one from Step 1 was converted to 1-propyl-4-isobutyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one as a colorless solid: NMR (CDCl₃) δ 0.90 (d, J = 7 Hz, 6H), 0.95 (t, J = 8 Hz, 3H), 1.62-1.78 (m, 3H), 2.16 (dd, J = 7 and 1 Hz, 2H), 3.61 (t, J = 7 Hz, 2H), 4.97 (s, 2H), 5.98 (s, 1H), 7.30 (d, J = 8 Hz, 1H), 7.45-7.55 (m, 3H), 7.60 (dd, J = 8 and 2 Hz, 1H), 8.01-8.07 (m, 1H), 8.49 (d, J = 2 Hz, 1H); MS (FAB) m/e (rel intensity) 418 (45), 390 (15), 375 (4), 361 (4), 237 (10), 209 (93), 195 (36), 180 (100); HRMS. Calc'd for M + H: 418.2355. Found: 418.2383.

Example 4



1-butyl-4-isobutyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one


Step 1: Preparation of 1-butyl-4-isobutyl-1,3-dihydro-2H-imidazol-2-one.

Following General Synthetic Schemes III and IV, 1-butyl-4-isobutyl-1,3-dihydro-2H-imidazol-2-one was prepared: NMR (CDCl₃) δ 0.91 (d, J = 7 Hz, 6H), 0.94 (t, J = 7 Hz, 3H), 1.28-1.41 (m, 2H), 1.57-1.68 (m, 2H), 1.75-1.87 (m, 1H), 2.21 (d, J = 7 Hz, 2H), 3.55 (t, J = 7 Hz, 2H), 5.86 (s, 1H), 8.86 (br s, 1H); MS (FAB) m/e (rel intensity) 197 (100), 181 (6), 173 (5), 167 (3), 153 (30), 141 (20), 125 (6), 111 (5); HRMS. Calc'd for M + H: 197.1654. Found: 197.1637.

Step 2: Preparation of 1-butyl-4-isobutyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one.

Following General Synthetic Schemes XII and XIV or XXIII, the imidazol-2-one from Step 1 was converted to 1-butyl-4-isobutyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one as a colorless solid: NMR (DMSO-d₆) δ 0.82 (d, J = 6 Hz, 6H), 0.88 (t, J = 7 Hz, 3H), 1.17-1.32 (m, 3H), 1.49-1.65 (m, 2H), 2.21 (d, J = 7 Hz, 2H), 3.52 (t, J = 7 Hz, 2H), 4.81 (s, 2H), 6.29 (s, 1H), 7.45 (d, J = 8 Hz, 1H), 7.58-7.79 (m, 5H), 8.25 (d, J = 2 Hz, 1H); MS (FAB) m/e (rel intensity) 432 (65), 404 (20), 375 (5), 237 (20), 209 (100), 194 (30), 180 (65); HRMS. Calc'd for M + H: 432.2512. Found: 432.2562.

Example 5




1-butyl-4-isopentyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one

Step 1: Preparation of 1-butyl-4-isopentyl-1,3-dihydro-2H-imidazol-2-one.

Following General Synthetic Schemes III and IV, 1-butyl-4-isopentyl-1,3-dihydro-2H-imidazol-2-one was prepared: NMR (CDCl₃) δ 0.90 (d, J=7 Hz, 6H), 0.93 (t, J=7 Hz, 3H), 1.26-1.67 (m, 7H), 2.35 (t, J=7 Hz, 2H), 3.54 (t, J=7 Hz, 2H), 5.83 (br s, 1H), 9.40 (br s, 1H); MS (FAB) m/e (rel intensity) 211 (100), 195 (8), 167 (15), 153 (25), 132 (8), 111 (20); HRMS. Calc'd for M+H: 211.1810. Found: 211.1802.

Step 2: Preparation of 1-butyl-4-isopentyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one.

Following General Synthetic Schemes XII and XIV or XXIII, the imidazol-2-one from Step 1 was converted to 1-butyl-4-isopentyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one as a colorless solid: NMR (CDCl₃) δ 0.87 (d, J=7 Hz, 6H), 0.95 (t, J=7 Hz, 3H), 1.30-1.43 (m, 4H), 1.50-1.72 (m, 3H), 2.28 (t, J=8 Hz, 2H), 3.62 (t, J=7 Hz, 2H), 4.97 (s, 2H), 5.95 (s, 1H), 7.32 (d, J=9 Hz, 1H), 7.46-7.56 (m, 3H), 7.69 (dd, J=9 and 2 Hz, 1H), 7.97-8.03 (m, 1H), 8.48 (br s, 1H); MS (FAB) m/e (rel intensity) 446 (100), 418 (25), 237 (15), 209 (75), 180 (65); HRMS. Calc'd for M+H: 446.2669. Found: 446.2654.


Example 6**1-butyl-4-pentyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one**Step 1: Preparation of 1-butyl-4-pentyl-1,3-dihydro-2H-imidazol-2-one.

Following General Synthetic Schemes III and IV, 1-butyl-4-pentyl-1,3-dihydro-2H-imidazol-2-one was prepared: NMR (CDCl₃) δ 0.88 (t, J=7 Hz, 3H), 0.92 (t, J=7 Hz, 3H), 1.24-1.43 (m, 6H), 1.47-1.70 (m, 4H), 2.35 (t, J=7 Hz, 2H), 3.55 (t, J=7 Hz, 2H), 5.82-5.87 (m, 1H), 9.60 (br s, 1H); MS (FAB) m/e (rel intensity) 211 (100), 155 (15); HRMS. Calc'd for M+H: 211.1810. Found: 211.1796.

Step 2: Preparation of 1-butyl-4-pentyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one.

Following General Synthetic Schemes XII and XIV or XXIII, the imidazol-2-one from Step 1 was converted to 1-butyl-4-pentyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one as a colorless solid: mp 176-177° C; NMR (CDCl₃) δ 0.87 (t, J=7 Hz, 3H), 0.96 (t, J=8 Hz, 3H), 1.24-1.43 (m, 6H), 1.45-1.56 (m, 2H), 1.61-1.72 (m, 2H), 2.29 (t, J=8 Hz, 2H), 3.63 (t, J=8 Hz, 2H), 4.96 (s, 2H), 5.97 (s, 1H), 7.45 (d, J=8 Hz, 1H), 7.50-7.62 (m, 3H), 7.79 (dd, J=8 and 2 Hz, 1H), 8.11-8.18 (m, 1H), 8.55 (d, J=2 Hz); MS (FAB) m/e (rel intensity) 446 (76), 418 (27), 403 (8), 390 (8), 389 (8), 237 (25), 209 (100), 194 (35), 180 (79); HRMS. Calc'd for M+H: 446.2668. Found: 446.2677.

Example 7



1-butyl-4-cyclohexyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one


Step 1: Preparation of 1-butyl-4-cyclohexyl-1,3-dihydro-2H-imidazol-2-one.

Following General Synthetic Schemes III and IV, 1-butyl-4-cyclohexyl-1,3-dihydro-2H-imidazol-2-one was prepared: NMR (CDCl₃) δ 0.93 (t, J = 7 Hz, 3H), 1.12-1.42 (m, 8H), 1.55-1.84 (m, 8H), 1.84-1.95 (m, 2H), 2.25-2.38 (m, 1H), 3.54 (t, J = 7 Hz, 2H), 5.79 (q, J = 1 Hz, 1H), 8.97 (br s, 1H); MS (FAB) m/e (rel intensity) 223 (100). HRMS. Calc'd for M + H: 223.1810. Found: 223.1811.

Step 2: Preparation of 1-butyl-4-cyclohexyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one.

Following General Synthetic Schemes XII and XIV or XXIII, the imidazol-2-one from Step 1 was converted to 1-butyl-4-cyclohexyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one as a colorless solid: NMR (CDCl₃) δ 0.96 (t, J = 7 Hz, 3H), 1.12-1.45 (m, 8H), 1.57-1.89 (m, 8H), 2.20-2.23 (m, 1H), 3.62 (t, J = 7 Hz, 2H), 5.02 (s, 2H), 5.96 (s, 1H), 7.53-7.68 (m, 4H), 8.03 (d, J = 8 Hz, 1H), 8.12-8.19 (m, 1H), 8.59 (d, J = 2 Hz, 1H); MS (FAB) m/e (rel intensity) 458 (70), 430 (20), 222 (20); HRMS. Calc'd for M + H: 458.2668. Found: 458.2676.

Example 8




1-butyl-4-cyclopentyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one

Step 1: Preparation of 1-butyl-4-cyclopentyl-1,3-dihydro-2H-imidazol-2-one.

Following General Synthetic Schemes III and IV, 1-butyl-4-cyclopentyl-1,3-dihydro-2H-imidazol-2-one was prepared: NMR (CDCl₃) δ0.94 (t, J = 7 Hz, 3H), 1.17-1.42 (m, 4H), 1.42-1.80 (m, 6H), 1.89-2.01 (m, 2H), 2.64-2.93 (m, 1H), 3.55 (t, J = 7 Hz, 2H), 5.84 (q, J = 1 Hz, 1H), 9.90 (br s, 1H); MS (FAB) m/e (rel intensity) 209 (100), 153 (10); HRMS. Calc'd for M + H: 209.1654. Found: 209.1725.

Step 2: Preparation of 1-butyl-4-cyclopentyl-1,3-dihydro-3-[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one.

Following General Synthetic Schemes XII and XIV or XXIII, the imidazol-2-one from Step 1 was converted to 1-butyl-4-cyclopentyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one as a colorless solid: mp 195-196° C (dec); NMR (CDCl₃) δ0.95 (t, J = 8 Hz, 3H), 1.30-1.49 (m, 4H), 1.53-1.77 (m, 6H), 1.83-1.95 (m, 2H), 2.61-2.74 (m, 1H), 3.62 (t, J = 8 Hz, 2H), 5.01 (s, 2H), 5.97 (d, J = 1 Hz, 1H), 7.37 (d, J = 8 Hz, 1H), 7.48-7.58 (m, 3H), 7.70 (dd, J = 8 and 2 Hz, 1H), 8.03-8.10 (m, 1H), 8.52 (d, J = 2 Hz, 1H); MS (FAB) m/e (rel intensity) 444 (57), 416 (17), 237 (20), 209 (100), 194 (34), 180 (70); HRMS. Calc'd for M + H: 444.2512. Found: 444.2487.


Example 9**1-butyl-4-(2-cyclopropylethyl)-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one**Step 1: Preparation of 1-butyl-4-(2-cyclopropylethyl)-1,3-dihydro-2H-imidazol-2-one.

Following General Synthetic Schemes III and IV, 1-butyl-4-(2-cyclopropylethyl)-1,3-dihydro-2H-imidazol-2-one was prepared: NMR (CDCl₃) δ0.00-0.07 (m, 2H), 0.37-0.45 (m, 2H), 0.61-0.76 (m, 1H), 0.92 (t, J = 7 Hz, 3H), 1.28-1.48 (m, 4H), 1.55-1.67 (m, 2H), 2.46 (t, J = 7 Hz, 2H), 3.54 (t, J = 7 Hz, 2H), 5.82-5.86 (m, 1H), 10.0 (br s, 1H); MS (FAB) m/e (rel intensity) 209 (100), 193 (4), 179 (3), 165 (8), 153 (30), 137 (4), 123 (4), 111 (10); HRMS. Calc'd for M + H: 209.1654. Found: 209.1721.

Step 2: Preparation of 1-butyl-4-(2-cyclopropylethyl)-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one.

Following General Synthetic Schemes XII and XIV or XXIII, the imidazol-2-one from Step 1 was converted to 1-butyl-4-(2-cyclopropylethyl)-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one as a colorless solid: NMR (CDCl₃) δ0.00-0.07 (m, 2H), 0.40-0.47 (m, 2H), 0.63-0.77 (m, 1H), 0.94 (t, J = 7 Hz, 3H), 1.03-1.45 (m, 4H), 1.65 (m, J = 7 Hz, 2H), 2.40 (t, J = 7 Hz, 2H), 3.62 (t, J = 7 Hz, 2H), 4.99 (s, 2H), 5.97 (s, 1H), 7.35 (d, J = 8 Hz, 1H), 7.48-7.56 (m, 3H), 7.74 (dd, J = 8 and 2 Hz, 1H), 7.97-8.04 (m, 1H), 8.50 (d, J = 1 Hz, 1H); MS (FAB) m/e (rel intensity) 444 (100), 416 (16), 401 (9), 388 (5), 237 (14), 209 (33), 194 (16), 180 (35); HRMS. Calc'd for M + H: 444.2512. Found: 444.2513.

Example 10



1-methyl-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one


Step 1: Preparation of 1-methyl-4-butyl-1,3-dihydro-2H-imidazol-2-one.

Following General Synthetic Schemes III and IV, 1-methyl-4-butyl-1,3-dihydro-2H-imidazol-2-one was prepared: NMR (CDCl₃) δ 0.89 (t, J = 7 Hz, 3H), 1.26-1.40 (m, 2H), 1.45-1.56 (m, 2H), 2.34 (t, J = 8 Hz, 2H), 3.18 (s, 3H), 5.79-5.83 (m, 1H), 10.18 (br s, 1H).

Step 2: Preparation of 1-methyl-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one.

Following General Synthetic Schemes XII and XIV or XXIII, the imidazol-2-one from Step 1 was converted to 1-methyl-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one as a colorless solid: mp 184-185° C (dec); NMR (CDCl₃) δ 0.88 (t, J = 8 Hz, 3H), 1.28-1.54 (m, 4H), 2.29 (t, J = 8 Hz, 2H), 3.28 (s, 3H), 4.97 (s, 1H), 5.95 (s, 1H), 7.38 (d, J = 8 Hz, 1H), 7.47-7.59 (m, 3H), 7.74 (dd, J = 8 and 2 Hz, 1H), 8.04-8.11 (m, 1H), 8.53 (d, J = 2 Hz, 1H); MS (FAB) m/e (rel intensity) 390 (100), 362 (12), 237 (17), 209 (46), 194 (18), 180 (40); HRMS. Calc'd for M + H: 390.2042. Found 390.2021.

Example 11




1-ethyl-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one

Step 1: Preparation of 1-ethyl-4-butyl-1,3-dihydro-2H-imidazol-2-one.

Following General Synthetic Schemes III and IV, 1-ethyl-4-butyl-1,3-dihydro-2H-imidazol-2-one was prepared: NMR (CDCl₃) δ 0.90 (t, J = 7 Hz, 3H), 1.24-1.41 (m, 2H), 1.27 (t, J = 8 Hz, 3H), 1.44-1.58 (m, 2H), 2.35 (t, J = 8 Hz, 2H), 3.59 (q, J = 8 Hz, 2H), 5.83-5.86 (m, 1H), 10.13 (br s, 1H); MS (FAB) m/e (rel intensity) 169 (100), 141 (4), 139 (5), 125 (17); HRMS. Calc'd for M + H: 169.1341. Found: 169.1365.

Step 2: Preparation of 1-ethyl-1-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one.

Following General Synthetic Schemes XII and XIV or XXIII, the imidazol-2-one from Step 1 was converted to 1-ethyl-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one as a colorless solid: mp 205° C (dec); NMR (CDCl₃) δ 0.89 (t, J = 7 Hz, 3H), 1.27-1.55 (m, 4H), 1.30 (t, J = 8 Hz, 3H), 2.31 (t, J = 8 Hz, 2H), 3.69 (q, J = 8 Hz, 2H), 4.98 (s, 2H), 6.00 (t, J = 1 Hz, 1H), 7.46 (d, J = 8 Hz, 1H), 7.52-7.62 (m, 3H), 7.84 (dd, J = 8 and 2 Hz, 1H), 8.08-8.15 (m, 1H), 8.55 (d, J = 2 Hz, 1H); MS (FAB) m/e (rel intensity) 404 (93), 376 (20), 237 (18), 209 (100), 194 (30), 180 (80), 168 (26); HRMS. Calc'd for M + H: 404.2199. Found: 404.2257.


Example 12**1-propyl-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one**Step 1: Preparation of 1-propyl-4-butyl-1,3-dihydro-2H-imidazol-2-one.

Following General Synthetic Schemes III and IV, 1-propyl-4-butyl-1,3-dihydro-2H-imidazol-2-one was prepared: NMR (CDCl₃) δ 0.87 (t, J = 7 Hz, 3H), 0.90 (t, J = 7 Hz, 3H), 1.25-1.39 (m, 2H), 1.44-1.56 (m, 2H), 1.58-1.61 (m, 2H), 2.34 (td, J = 8 and 1 Hz, 2H), 3.49 (t, J = 8 Hz, 2H), 5.82 (t, J = 1 Hz, 1H), 10.33 (br s, 1H); MS (FAB) m/e (rel intensity) 183 (100), 141 (13), 139 (15); HRMS. Calc'd for M + H: 183.1497. Found: 183.1497.

Steps 2: Preparation of 1-propyl-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one.

Following General Synthetic Schemes XII and XIV or XXIII, the imidazol-2-one from Step 1 was converted to 1-propyl-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one as a colorless solid: mp 164-165° C; NMR (CDCl₃) δ 0.88 (t, J = 7 Hz, 3H), 0.94 (t, J = 7 Hz, 3H), 1.29-1.55 (m, 4H), 1.63-1.77 (m, 2H), 2.30 (td, J = 8 and 1 Hz, 2H), 3.60 (t, J = 8 Hz, 2H), 4.97 (s, 1H), 5.98 (t, J = 1 Hz, 1H), 7.43 (d, J = 8 Hz, 1H), 7.50-7.61 (m, 3H), 7.80 (dd, J = 8 and 2 Hz, 1H), 8.05-8.12 (m, 1H), 8.53 (d, J = 2 Hz, 1H); MS (FAB) m/e (rel intensity) 418 (100), 390 (12), 209 (20), 180 (21); HRMS. Calc'd for M + H: 418.2355. Found: 418.2312.

Example 13



20 **1-isobutyl-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one**


Step 1: Preparation of 1-isobutyl-4-butyl-1,3-dihydro-2H-imidazol-2-one.

25 Following General Synthetic Schemes III and IV, 1-isobutyl-4-butyl-1,3-dihydro-2H-imidazol-2-one was prepared: NMR (CDCl₃) δ 0.90 (t, J = 7 Hz, 3H), 0.92 (d, J = 7 Hz, 6H), 1.27-1.41 (m, 2H), 1.46-1.58 (m, 2H), 1.97 (m, J = 7 Hz, 1H), 2.36 (td, J = 8 and 1 Hz, 2H), 3.36 (d, J = 8 Hz, 2H), 5.84 (t, J = 1 Hz, 1H), 9.86 (br s, 1H); MS (FAB) m/e (rel intensity) 197 (100), 141 (22); HRMS. Calc'd for M+H: 197.1654. Found: 197.1635.

30 Step 2: Preparation of 1-isobutyl-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one.

35 Following General Synthetic Schemes XII and XIV or XXIII, the imidazol-2-one from Step 1 was converted to 1-isobutyl-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one as a colorless solid: mp 160.0-161.5° C; NMR (CDCl₃) δ 0.87 (t, J = 7 Hz, 3H), 0.92 (d, J = 7 Hz, 6H), 1.27-1.53 (m, 4H), 1.99 (m, J = 7 Hz, 1H), 2.28 (t, J = 8 Hz, 2H), 3.43 (d, J = 7 Hz, 2H), 4.97 (s, 2H), 5.93 (s, 1H), 7.27 (d, J = 8 Hz, 1H), 7.43-7.52 (m, 3H), 7.65 (dd, J = 8 and 2 Hz, 1H), 7.91-7.98 (m, 1H), 8.45 (d, J = 2 Hz, 1H); MS (FAB) m/e (rel intensity) 432 (100), 404 (15), 209 (23), 180 (25); HRMS. Calc'd for M+H: 432.2512. Found: 432.2571.

40 **Example 14**




1-isopentyl-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one**Step 1: Preparation of 1-isopentyl-4-butyl-1,3-dihydro-2H-imidazol-2-one.**

Following General Synthetic Schemes III and IV, 1-isopentyl-4-butyl-1,3-dihydro-2H-imidazol-2-one was prepared: NMR (CDCl₃) δ 0.89 (t, J = 7 Hz, 3H), 0.92 (d, J = 7 Hz, 6H), 1.26-1.40 (m, 2H), 1.46-1.66 (m, 6H), 2.35 (td, J = 8 and 1 Hz, 2H), 3.55 (t, J = 8 Hz, 2H), 5.82 (t, J = 1 Hz, 1H), 10.41 (br s, 1H); MS (FAB) m/e (rel intensity) 211 (100), 141 (13), 111 (14); HRMS. Calc'd for M + H: 211.1810. Found: 211.1800.

Step 2: Preparation of 1-isopentyl-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one.

Following General Synthetic Schemes XII and XIV or XXIII, the imidazol-2-one from Step 1 was converted to 1-isopentyl-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one as a colorless solid: mp 168.5-170.0° C; NMR (CDCl₃) δ 0.88 (t, J = 7 Hz, 3H), 0.95 (d, J = 6 Hz, 6H), 1.27-1.67 (m, 7H), 2.38 (t, J = 8 Hz, 2H), 3.63 (t, J = 8 Hz, 2H), 4.95 (s, 2H), 5.96 (s, 1H), 7.32 (d, J = 8 Hz, 1H), 7.44-7.56 (m, 3H), 7.69 (dd, J = 8 and 1 Hz, 1H), 8.00 (td, J = 3 and 1 Hz, 1H), 8.47 (br s, 1H); MS (FAB) m/e (rel intensity) 446 (100), 418 (21), 403 (8), 237 (12), 209 (30), 194 (13), 180 (28); HRMS. Calc'd for M + H: 446.2662. Found: 446.2668.

Example 15**1-pentyl-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one****Step 1: Preparation of 1-pentyl-4-butyl-1,3-dihydro-2H-imidazol-2-one.**


Following General Synthetic Schemes III and IV, 1-pentyl-4-butyl-1,3-dihydro-2H-imidazol-2-one was prepared: NMR (CDCl₃) δ 0.87 (t, J = 7 Hz, 3H), 0.89 (t, J = 7 Hz, 3H), 1.21-1.39 (m, 6H), 1.42-1.69 (m, 4H), 2.34 (td, J = 8 and 1 Hz, 2H), 3.52 (t, J = 8 Hz, 2H), 5.82 (t, J = 1 Hz, 1H), 10.45 (br s, 1H); MS (FAB) m/e (rel intensity) 211 (100), 141 (19); HRMS. Calc'd for M + H: 211.1810. Found: 211.1792.

Step 2: Preparation of 1-pentyl-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one.

Following General Synthetic Schemes XII and XIV or XXIII, the imidazol-2-one from Step 1 was converted to 1-pentyl-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one as a colorless solid: mp 162-164° C; NMR (CDCl₃) δ 0.87 (t, J = 7 Hz, 3H), 0.89 (t, J = 7 Hz, 3H), 1.23-1.41 (m, 6H), 1.42-1.54 (m, 2H), 1.61-1.73 (m, 2H), 2.39 (t, J = 8 Hz, 2H), 3.61 (t, J = 8 Hz, 2H), 4.96 (s, 2H), 5.96 (s, 1H), 7.39 (d, J = 8 Hz, 1H), 7.48-7.58 (m, 3H), 7.76 (dd, J = 8 and 1 Hz, 1H), 8.02-8.09 (m, 1H), 8.51 (d, J = 1 Hz, 1H); MS (FAB) m/e (rel intensity) 446 (58), 418 (22), 237 (18), 209 (100), 194 (29), 180 (55);

HRMS. Calc'd for M + H: 446.2668. Found: 446.2683.

Example 16



1-isopropyl-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one


Step 1: Preparation of 1-isopropyl-4-butyl-1,3-dihydro-2H-imidazol-2-one.

Following General Synthetic Schemes III and IV, 1-isopropyl-4-butyl-1,3-dihydro-2H-imidazol-2-one was prepared: NMR (CDCl₃) δ 0.88 (t, J = 8 Hz, 3H), 1.24 (d, J = 7 Hz, 6H), 1.28-1.39 (m, 2H), 1.43-1.59 (m, 2H), 2.34 (td, J = 8 and 1 Hz, 2H), 4.34 (m, J = 7 Hz, 1H), 5.87 (t, J = 1 Hz, 1H), 10.39 (br s, 1H); MS (FAB) m/e (rel intensity) 183 (100), 141 (48), 139 (37); HRMS. Calc'd for M + H: 183.1497. Found: 183.1472.

Step 2: Preparation of 1-isopropyl-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one.

Following General Synthetic Schemes XII and XIV or XXIII, the imidazol-2-one from Step 1 was converted to 1-isopropyl-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one as a colorless solid: NMR (CDCl₃) δ 0.87 (t, J = 7 Hz, 3H), 1.30 (d, J = 7 Hz, 6H), 1.22-1.41 (m, 2H), 1.41-1.53 (m, 2H), 2.28 (t, J = 8 Hz, 2H), 4.42 (m, J = 7 Hz, 1H), 4.96 (s, 1H), 6.00 (s, 1H), 7.27 (d, J = 8 Hz, 1H), 7.43-7.52 (m, 3H), 7.65 (dd, J = 8 and 2 Hz, 1H), 7.89-7.96 (m, 1H), 8.44 (d, J = 2 Hz, 1H); MS (FAB) m/e (rel intensity) 418 (100), 390 (35), 375 (10), 361 (10), 237 (15), 209 (100), 193 (30), 180 (90); HRMS. Calc'd for M + H: 418.2355. Found: 418.2381.

Example 17



1-tertbutyl-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one


Step 1: Preparation of 1-tertbutyl-4-butyl-1,3-dihydro-2H-imidazol-2-one.

Following General Synthetic Schemes III and IV, 1-tertbutyl-4-butyl-1,3-dihydro-2H-imidazol-2-one was prepared: NMR (CDCl₃) δ 0.90 (t, J = 8 Hz, 3H), 1.27-1.38 (m, 2H), 1.45-1.59 (m, 2H), 1.51 (s, 9H), 2.32 (td, J = 8 and 1 Hz, 2H), 5.94 (t, J = 1 Hz, 1H), 9.86 (br s, 1H); MS (FAB) m/e (rel intensity) 197 (73), 141 (100); HRMS. Calc'd for M + H: 197.1654. Found: 197.1670.

Step 2: Preparation of 1-tertbutyl-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one.

Following General Synthetic Schemes XII and XIV, the imidazol-2-one from Step 1 was converted to 1-tertbutyl-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one as a colorless solid: NMR (CDCl₃) δ 0.88 (t, J = 7 Hz, 3H), 1.22-1.59 (m, 4H), 1.55 (s, 9H), 2.27 (t, J = 8 Hz, 2H), 4.90 (s, 2H), 6.06 (s, 1H), 7.37 (d, J = 8 Hz, 1H), 7.48-7.58 (m, 3H), 7.71 (dd, J = 8 and 2 Hz, 1H), 7.99-8.06 (m, 1H), 8.48 (d, J = 2 Hz, 1H); MS (FAB) m/e (rel intensity) 432 (100), 404 (20), 389 (5), 375 (15), 348 (10), 237 (15), 209 (95), 194 (25), 180 (55); HRMS. Calc'd for M + H: 432.2512. Found: 432.2578.

Example 18




1-neopentyl-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one

Step 1: Preparation of 1-neopentyl-4-butyl-1,3-dihydro-2H-imidazol-2-one.

Following General Synthetic Schemes III and IV, 1-neopentyl-4-butyl-1,3-dihydro-2H-imidazol-2-one was prepared: NMR (CDCl₃) δ 0.87 (t, J = 8 Hz, 3H), 0.92 (s, 9H), 1.23-1.37 (m, 2H), 1.43-1.55 (m, 2H), 2.31 (td, J = 8 and 1 Hz, 2H), 3.31 (s, 2H), 5.80 (t, J = 1 Hz, 1H), 10.62 (br s, 1H); MS (FAB) m/e (rel intensity) 211 (100), 141 (22); HRMS. Calc'd for M + H: 211.1810. Found: 211.1745.

Step 2: Preparation of 1-neopentyl-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one.

Following General Synthetic Schemes XII and XIV or XXIII, the imidazol-2-one from Step 1 was converted to 1-neopentyl-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one as a colorless solid: mp. 194.5-195.0 °C; NMR (CDCl₃) δ 0.89 (t, J = 7 Hz, 3H), 0.97 (s, 9H), 1.29-1.55 (m, 4H), 2.31 (t, J = 8 Hz, 2H), 3.41 (s, 2H), 4.98 (s, 2H), 5.96 (s, 1H), 7.43 (d, J = 8 Hz, 1H), 7.50-7.59 (m, 3H), 7.82 (dd, J = 8 and 2 Hz, 1H), 8.06-8.13 (m, 1H), 8.54 (d, J = 2 Hz, 1H); MS (FAB) m/e (rel intensity) 446 (100), 418 (17), 403 (6), 237 (11), 209 (27), 194 (11), 180 (27); HRMS. Calc'd for M + H: 446.2668. Found: 446.2647.


Example 19**1-(3,3-dimethylbutyl)-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one**Step 1: Preparation of 1-(3,3-dimethylbutyl)-4-butyl-1,3-dihydro-2H-imidazol-2-one.

Following General Synthetic Schemes III and IV, 1-(3,3-dimethylbutyl)-4-butyl-1,3-dihydro-2H-imidazol-2-one was prepared: NMR (CDCl₃) δ 0.92 (t, J = 8 Hz, 3H), 0.96 (s, 9H), 1.28-1.42 (m, 2H), 1.48-1.59 (m, 4H), 2.38 (td, J = 8 and 1 Hz, 2H), 3.54-3.62 (m, 2H), 5.58 (t, J = 1 Hz, 1H), 9.70 (br s, 1H); MS (FAB) m/e (rel intensity) 225 (100), 141 (13); HRMS. Calc'd for M + H: 225.1967. Found: 225.1897.

Step 2: Preparation of 1-(3,3-dimethylbutyl)-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one.

Following General Synthetic Schemes XII and XIV or XXIII, the imidazol-2-one from Step 1 was converted to 1-(3,3-dimethylbutyl)-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one as a colorless solid: NMR (CDCl₃) δ 0.89 (t, J = 8 Hz, 3H), 0.97 (s, 9H), 1.29-1.54 (m, 4H), 1.55-1.63 (m, 2H), 2.30 (t, J = 8 Hz, 2H), 3.60-3.68 (m, 2H), 4.95 (s, 2H), 5.97 (s, 1H), 7.38 (d, J = 8 Hz, 1H), 7.48-7.59 (m, 3H), 7.75 (dd, J = 8 and 2 Hz, 1H), 8.02-8.09 (m, 1H), 8.51 (d, J = 2 Hz, 1H); MS (FAB) m/e (rel intensity) 460 (47), 432 (23), 237 (18), 209 (100), 194 (32), 180 (74); HRMS. Calc'd for M + H: 460.2825. Found: 460.2821.

Example 20



1-(2-ethylbutyl)-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one


Step 1: Preparation of 1-(2-ethylbutyl)-4-butyl-1,3-dihydro-2H-imidazol-2-one.

Following General Synthetic Schemes III and IV, 1-(2-ethylbutyl)-4-butyl-1,3-dihydro-2H-imidazol-2-one was prepared: NMR (CDCl₃) δ 0.80-0.91 (m, 9H), 1.18-1.37 (m, 6H), 1.41-1.64 (m, 3H), 2.32 (td, J=8 and 1 Hz, 2H), 3.41 (d, J=8 Hz, 2H), 5.72 (t, J=1 Hz, 1H), 10.60 (br s, 1H); MS (FAB) m/e (rel intensity) 225 (100), 141 (22); HRMS. Calc'd for M+H: 225.1967. Found: 225.1971.

Step 2: Preparation of 1-(2-ethylbutyl)-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one.

Following General Synthetic Schemes XII and XIV or XXIII, the imidazol-2-one from Step 1 was converted to 1-(2-ethylbutyl)-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one as a colorless solid: mp 174-176° C (dec); NMR (CDCl₃) δ 0.85-0.96 (m, 9H), 1.27-1.40 (m, 6H), 1.41-1.55 (m, 2H), 1.59-1.70 (m, 1H), 2.30 (t, J=8 Hz, 2H), 3.53 (d, J=8 Hz, 2H), 4.97 (s, 2H), 5.94 (s, 1H), 7.39 (d, J=8 Hz, 1H), 7.48-7.58 (m, 3H), 7.75 (dd, J=8 and 2 Hz, 1H), 8.03-8.09 (m, 1H), 8.51 (d, J=2 Hz, 1H); MS (FAB) m/e (rel intensity) 460 (73), 432 (25), 237 (23), 209 (100), 194 (29), 180 (61); HRMS. Calc'd for M+H: 460.2825. Found: 460.2862.

Example 21




1-cyclohexyl-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-oneStep 1: Preparation of 1-cyclohexyl-4-butyl-1,3-dihydro-2H-imidazol-2-one.

Following General Synthetic Schemes III and IV, 1-cyclohexyl-4-butyl-1,3-dihydro-2H-imidazol-2-one was prepared: NMR (CDCl₃) δ 0.91 (t, J = 7 Hz, 3H), 1.05-1.24 (m, 2H), 1.26-1.46 (m, 4H), 1.47-1.58 (m, 2H), 1.63-1.76 (m, 2H), 1.78-1.96 (m, 4H), 2.36 (td, J = 8 and 1 Hz, 2H), 3.87-4.00 (m, 1H), 5.89 (t, J = 1 Hz, 1H), 9.82 (br s, 1H); MS (FAB) m/e (rel intensity) 223 (100), 141 (53); HRMS. Calc'd for M + H: 223.1810. Found: 223.1738.

Step 2: Preparation of 1-cyclohexyl-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one.

Following General Synthetic Schemes XII and XIV or XXIII, the imidazol-2-one from Step 1 was converted to 1-cyclohexyl-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one as a colorless solid: mp 202-203° C (dec); NMR (CDCl₃) δ 0.89 (t, J = 8 Hz, 3H), 1.08-1.24 (m, 2H), 1.26-1.54 (m, 7H), 1.66-1.76 (m, 1H), 1.77-2.00 (m, 4H), 2.28 (t, J = 8 Hz, 2H), 3.95-4.07 (m, 1H), 4.97 (s, 2H), 6.00 (s, 1H), 7.32 (d, J = 8 Hz, 1H), 7.45-7.55 (m, 3H), 7.69 (dd, J = 8 and 2 Hz, 1H), 7.96-8.04 (m, 1H), 8.47 (d, J = 2 Hz, 1H); MS (FAB) m/e (rel intensity) 458 (56), 430 (19), 237 (19), 208 (100), 194 (36), 180 (76); HRMS. Calc'd for M + H: 458.2668. Found: 458.2732.

Example 22**1-(2-cyclopropylethyl)-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one**Step 1: Preparation of 1-(2-cyclopropylethyl)-4-butyl-1,3-dihydro-2H-imidazol-2-one.


Following General Synthetic Schemes III and IV or XXIII, 1-(2-cyclopropylethyl)-4-butyl-1,3-dihydro-2H-imidazol-2-one was prepared: NMR (CDCl₃) δ 0.00-0.06 (m, 2H), 0.38-0.46 (m, 2H), 0.59-0.73 (m, 1H), 0.89 (t, J = 8 Hz, 3H), 1.16-1.40 (m, 4H), 1.44-1.57 (m, 4H), 2.34 (td, J = 8 and 1 Hz, 2H), 3.62 (t, J = 7 Hz, 2H), 5.84 (t, J = 1 Hz, 1H), 10.38 (br s, 1H); MS (FAB) m/e (rel intensity) 209 (100), 153 (43), 141 (16); HRMS. Calc's for M + H: 209.1654. Found: 209.1695.

Step 2: Preparation of 1-(2-cyclopropylethyl)-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one.

Following General Synthetic Schemes XII and XIV or XXIII, the imidazol-2-one from Step 1 was converted to 1-(2-cyclopropylethyl)-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-

2H-imidazol-2-one as a colorless solid: mp 182-183° C; NMR (CDCl₃) δ 0.04-0.11 (m, 2H), 0.43-0.51 (m, 2H), 0.60-0.76 (m, 1H), 0.91 (t, J=7 Hz, 3H), 1.30-1.44 (m, 2H), 1.45-1.64 (m, 4H), 2.36 (t, J=7 Hz, 2H), 3.72 (t, J=7 Hz, 2H), 5.04 (s, 2H), 6.01 (s, 1H), 7.57-7.68 (m, 4H), 8.11 (d, J=7 Hz, 1H), 8.15-8.20 (m, 1H), 8.64 (s, 1H); MS (FAB) m/e (rel intensity) 237 (24), 209 (53), 192 (18), 180 (43); HRMS. Calc'd for M+H: 444.2512. Found: 444.2509.

Example 23



1-cyclopentyl-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one


Step 1: Preparation of 1-cyclopentyl-4-butyl-1,3-dihydro-2H-imidazol-2-one.

Following General Synthetic Schemes III and IV, 1-cyclopentyl-4-butyl-1,3-dihydro-2H-imidazol-2-one was prepared: NMR (CDCl₃) δ 0.91 (t, J=8 Hz, 3H), 1.29-1.42 (m, 2H), 1.48-1.85 (m, 10H), 1.98-2.12 (m, 2H), 2.47 (td, J=8 and 1 Hz, 2H), 4.44-4.57 (m, 1H), 5.88 (t, J=1 Hz, 1H), 10.22 (br s, 1H); MS (FAB) m/e (rel intensity) 209 (100), 141 (43); HRMS. Calc'd for M+H: 209.1654. Found: 209.1656.

Step 2: Preparation of 1-cyclopentyl-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one.

Following General Synthetic Schemes XII and XIV or XXIII, the imidazol-2-one from Step 1 was converted to 1-cyclopentyl-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one as a colorless solid: NMR (CDCl₃) δ 0.88 (t, J=7 Hz, 3H), 1.30-1.40 (m, 2H), 1.41-1.52 (m, 2H), 1.56-1.86 (m, 6H), 2.02-2.14 (m, 2H), 2.27 (t, J=7 Hz, 2H), 4.55 (m, J=7 Hz, 1H), 4.97 (s, 2H), 5.98 (s, 1H), 7.27 (d, J=8 Hz, 1H), 7.44-7.54 (m, 3H), 7.62 (dd, J=8 and 2 Hz, 1H), 7.98-8.05 (m, 1H), 8.48 (d, J=2 Hz, 1H); MS (FAB) m/e (rel intensity) 444 (100), 416 (12), 401 (6), 387 (4), 233 (13), 209 (22), 194 (12), 180 (25); HRMS. Calc'd for M+H: 444.2512. Found: 444.2518.

Example 24



1-cyclopentylmethyl-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one


Step 1: Preparation of 1-cyclopentylmethyl-4-butyl-1,3-dihydro-2H-imidazol-2-one.

Following General Synthetic Schemes III and IV, 1-cyclopentylmethyl-4-butyl-1,3-dihydro-2H-imidazol-2-one was prepared: NMR (CDCl₃) δ 0.89 (t, J=8 Hz, 3H), 1.16-1.39 (m, 4H), 1.44-1.77 (m, 8H), 2.21 (m, J=8 Hz, 1H), 2.45 (t, J=8 Hz, 2H), 3.46 (d, J=8 Hz, 2H), 5.81-5.85 (m, 1H), 10.33 (br s, 1H); MS (FAB) m/e (rel intensity) 223 (100), 141 (30); HRMS. Calc'd for M+H: 223.1810. Found: 223.1767.

Step 2: Preparation of 1-cyclopentylmethyl-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one.

Following General Synthetic Schemes XII and XIV or XXIII, the imidazol-2-one from Step 1 was converted to 1-cyclopentylmethyl-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one as a colorless solid: mp 204° C (dec); NMR (CDCl₃) δ 0.88 (t, J=7 Hz, 3H), 1.19-1.70 (m, 12H), 2.19-2.34 (m, 3H), 3.55 (d, J=8 Hz, 2H), 4.98 (s, 2H), 5.98 (t, J=1 Hz, 1H), 7.44 (d, J=8 Hz, 1H), 7.50-7.61 (m, 3H), 7.81 (dd, J=8 and 2 Hz, 1H), 8.07-8.13 (m, 1H), 8.53 (d, J=2 Hz, 1H); MS (FAB) m/e (rel intensity) 458 (65), 430 (22), 237 (24), 209 (100), 194 (30), 180 (65); HRMS. Calc'd for M+H: 458.2668. Found: 458.2699.

Example 25



1-[2-(2-thienyl)ethyl]-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-oneStep 1: Preparation of 1-[2-(2-thienyl)ethyl]-4-butyl-1,3-dihydro-2H-imidazol-2-one.

5

Following General Synthetic Schemes III and IV, 1-[2-(2-thienyl)ethyl]-4-butyl-1,3-dihydro-2H-imidazol-2-one was prepared: NMR (CDCl₃) δ 0.90 (t, J = 7 Hz, 3H), 1.26-1.40 (m, 2H), 1.44-1.56 (m, 2H), 2.32 (t, J = 7 Hz, 2H), 3.16 (t, J = 7 Hz, 2H), 3.82 (t, J = 7 Hz, 2H), 5.69-5.75 (m, 1H), 6.80-6.83 (m, 1H), 6.90-6.95 (m, 1H), 7.12-7.17 (m, 1H), 10.02 (br s, 1H); MS (FAB) m/e (rel intensity) 251 (90), 111 (100); HRMS. Calc'd for M + H: 251.1218. Found: 251.1209.

10

Step 2: Preparation of 1-[2-(2-thienyl)ethyl]-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one.

15

Following General Synthetic Schemes XII and XIV or XXIII, the imidazol-2-one from Step 1 was converted to 1-[2-(2-thienyl)ethyl]-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one as a colorless solid: mp 137-138° C; NMR (CDCl₃) δ 0.86 (t, J = 7 Hz, 3H), 1.23-1.36 (m, 2H), 1.37-1.51 (m, 2H), 2.26 (td, J = 8 and 1 Hz, 2H), 3.23 (t, J = 8 Hz, 2H), 3.93 (t, J = 8 Hz, 2H), 4.93 (s, 2H), 5.84 (t, J = 1 Hz, 1H), 6.84 (d, J = 8 Hz, 1H), 6.91-6.96 (m, 1H), 7.17 (dd, J = 8 and 2 Hz, 1H), 7.49 (d, J = 8 Hz, 1H), 7.53-7.64 (m, 3H), 7.69 (dd, J = 8 and 2 Hz, 1H), 8.25-8.30 (m, 1H), 8.57 (d, J = 2 Hz, 1H); HRMS. Calc'd for M + H: 486.2076. Found: 486.2092.

20


Example 26

25

30

35

40

**1-(2-phenylethyl)-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one**

45

Step 1: Preparation of 1-(2-phenylethyl)-4-butyl-1,3-dihydro-2-imidazol-2-one.

50

Following General Synthetic Schemes III and IV, 1-(2-phenylethyl)-4-butyl-1,3-dihydro-2-imidazol-2-one was prepared: NMR (CDCl₃) δ 0.90 (t, J = 7 Hz, 3H), 1.31 (m, J = 7 Hz, 2H), 1.49 (m, J = 7 Hz, 2H), 2.32 (td, J = 7 and 1 Hz, 2H), 2.94 (t, J = 7 Hz, 2H), 3.80 (t, J = 7 Hz, 2H), 5.66 (t, J = 1 Hz, 1H), 7.15-7.33 (m, 5H), 9.89 (br s, 1H); MS (FAB) m/e (rel intensity) 245 (100), 153 (10), 105 (50).


55

Step 2: Preparation of 1-(2-phenylethyl)-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one.

Following General Synthetic Schemes XII and XIV or XXIII, the imidazol-2-one from Step 1 was converted to 1-(2-phenylethyl)-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-

imidazol-2-one as a colorless solid: NMR (CDCl₃) δ 0.85 (t, J=7 Hz, 3H), 1.22-1.47 (m, 4H), 2.24 (t, J=7 Hz, 2H), 2.99 (t, J=7 Hz, 2H), 3.90 (t, J=7 Hz, 2H), 4.93 (s, 2H), 5.80 (s, 1H), 7.16-7.47 (m, 5H), 7.48-7.59 (m, 4H), 8.05-8.12 (m, 1H), 8.47 (s, 1H); MS (FAB) m/e (rel intensity) 480 (100), 452 (20), 437 (5), 237 (18), 209 (8), 180 (81); HRMS. Calc'd for M + H: 480.2512. Found: 480.2561.

Example 27



1-(2-cyclopentylethyl)-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one


Step 1: Preparation of 1-(2-cyclopentylethyl)-4-butyl-1,3-dihydro-2H-imidazol-2-one.

Following General Synthetic Schemes III and IV, 1-(2-cyclopentylethyl)-4-butyl-1,3-dihydro-2H-imidazol-2-one was prepared: NMR (CDCl₃) δ 0.88 (t, J=7 Hz, 3H), 1.00-1.16 (m, 2H), 1.25-1.39 (m, 2H), 1.41-1.67 (m, 8H), 1.68-1.84 (m, 3H), 2.33 (t, J=8 Hz, 2H), 3.54 (t, J=8 Hz, 2H), 5.79-5.83 (m, 1H), 10.58 (br s, 1H); MS (FAB) m/e (rel intensity) 273 (100), 141 (8); HRMS. Calc'd for M + H: 237.1967. Found: 237.1898.

Step 2: Preparation of 1-(2-cyclopentylethyl)-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one.

Following General Synthetic Scheme XII and XIV or XXIII, the imidazol-2-one from Step 1 was converted to 1-(2-cyclopentylethyl)-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one as a colorless solid: NMR (CDCl₃) δ 0.88 (t, J=7 Hz, 3H), 1.06-1.20 (m, 2H), 1.23-1.87 (m, 13H), 2.23 (t, J=7 Hz, 2H), 3.63 (t, J=7 Hz, 2H), 4.95 (s, 2H), 5.96 (s, 1H), 7.28 (d, J=8 Hz, 1H), 7.44-7.54 (m, 3H), 7.62 (dd, J=8 and 2 Hz, 1H), 7.97-8.04 (m, 1H), 8.48 (d, J=2 Hz, 1H); MS (FAB) m/e (rel intensity) 472 (100), 444 (27), 429 (5), 416 (3), 237 (19), 209 (65), 195 (19); HRMS. Calc'd for M + H: 472.2825. Found: 472.2881.

Example 28



1-[2-(cyclopenten-1-yl)ethyl]-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one


Step 1: Preparation of 1-[2-(cyclopenten-1-yl)ethyl]-4-butyl-1,3-dihydro-2H-imidazol-2-one.

Following General Synthetic Schemes III and IV, 1-[2-(cyclopenten-1-yl)ethyl]-4-butyl-1,3-dihydro-2H-imidazol-2-one was prepared: NMR (CDCl₃) δ 0.90 (t, J = 7 Hz, 3H), 1.22-1.41 (m, 2H), 1.44-1.58 (m, 2H), 1.77-1.91 (m, 2H), 2.18-2.46 (m, 8H), 3.68 (t, J = 8 Hz, 2H), 5.36-5.43 (m, 1H), 5.82-5.87 (m, 1H), 9.23 (br s, 1H); MS (FAB) m/e (rel intensity) 235 (90), 141 (100); HRMS. Calc'd for M + H: 235.1810. Found: 235.1804.

Step 2: Preparation of 1-[2-(cyclopenten-1-yl)ethyl]-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one.

Following General Synthetic Schemes XII and XIV or XXIII, the imidazol-2-one from Step 1 was converted to 1-[2-(cyclopenten-1-yl)ethyl]-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one as a colorless solid: NMR (CDCl₃) δ 0.88 (t, J = 7 Hz, 3H), 1.21-1.39 (m, 2H), 1.39-1.53 (m, 2H), 1.86 (m, J = 7 Hz, 2H), 2.28 (t, J = 7 Hz, 6H), 2.45 (t, J = 7 Hz, 2H), 3.76 (t, J = 7 Hz, 2H), 4.95 (s, 2H), 5.41 (s, 1H), 5.96 (s, 1H), 7.45 (d, J = 8 Hz, 1H), 7.49-7.63 (m, 3H), 7.76 (d, J = 8 Hz, 1H), 8.12-8.20 (m, 1H), 8.55 (s, 1H); MS (FAB) m/e (rel intensity) 470 (30), 442 (10); HRMS. Calc'd for M + H: 470.2668. Found: 470.2599.

Example 29



1-(2-adamantyl)-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one


Step 1: Preparation of 1-(2-adamantyl)-4-butyl-1,3-dihydro-2H-imidazol-2-one.

Following General Synthetic Schemes III and IV, 1-(2-adamantyl)-4-butyl-1,3-dihydro-2H-imidazol-2-one was prepared: NMR (CDCl₃) δ 0.91 (t, J = 7 Hz, 3H), 1.27-1.42 (m, 2H), 1.47-1.59 (m, 2H), 1.63 (br s, 1H), 1.67 (br s, 1H), 1.78 (br s, 2H), 1.84-2.02 (m, 8H), 2.37 (t, J = 8 Hz, 2H), 2.42 (br s, 2H), 4.09 (s, 1H), 6.18-6.22 (m, 1H), 9.53 (br s, 1H); MS (FAB) m/e (rel intensity) 275 (100), 141 (30), 135 (30); HRMS. Calc'd for M + H: 275.2123. Found: 275.2065.

Step 2: Preparation of 1-(2-adamantyl)-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one.

Following General Synthetic Schemes XII and XIV or XXIII, the imidazol-2-one from Step 1 was converted to 1-(2-adamantyl)-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one as a colorless solid: NMR (CDCl₃) δ 0.91 (t, J = 7 Hz, 3H), 1.32-1.45 (m, 2H), 1.47-1.58 (m, 2H), 1.62-2.00 (m, 12H), 2.34 (t, J = 8 Hz, 2H), 2.47 (br s, 2H), 4.15 (s, 1H), 4.96 (s, 2H), 6.35 (s, 1H), 7.51-7.58 (m, 2H), 7.61-7.68 (m, 2H), 7.93 (dd, J = 8 and 2 Hz, 1H), 8.13 (dd, J = 6 and 4 Hz, 1H), 8.56 (d, J = 2 Hz, 1H); MS (FAB) m/e (rel intensity) 510 (100), 482 (10), 467 (5), 237 (10), 209 (10); HRMS. Calc'd for M + H: 510.2981. Found: 510.3038.

Example 30



1-(1-adamantyl)-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one


Step 1: Preparation of 1-(1-adamantyl)-4-butyl-1,3-dihydro-2H-imidazol-2-one.

Following General Synthetic Schemes III and IV, 1-(1-adamantyl)-4-butyl-1,3-dihydro-2H-imidazol-2-one was prepared: NMR (CDCl₃) δ 0.92 (t, J = 8 Hz, 3H), 1.29-1.43 (m, 2H), 1.47-1.59 (m, 2H), 1.72 (br s, 1H), 2.16 (br s, 3H), 2.21 (br s, 6H), 2.36 (t, J = 8 Hz, 2H), 6.01 (br s, 1H), 9.43 (br s, 1H); MS (FAB) m/e (rel intensity) 275 (100), 141 (6), 135 (83); HRMS. Calc'd for M + H: 275.2123. Found: 275.2057.

Step 2: Preparation of 1-(1-adamantyl)-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one.

Following General Synthetic Schemes XII and XIV or XXIII, the imidazol-2-one from Step 1 was converted to 1-(1-adamantyl)-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one as a colorless solid: NMR (CDCl₃) δ 0.89 (t, J = 7 Hz, 3H), 1.30-1.42 (m, 2H), 1.42-1.54 (m, 2H), 1.64-1.71 (m, 6H), 2.07-2.34 (m, 11H), 4.90 (s, 2H), 6.08 (s, 1H), 7.47-7.66 (m, 4H), 7.78 (dd, J = 8 and 2 Hz, 1H), 8.20-8.25 (m, 1H), 8.57 (d, J = 2 Hz, 1H); MS (FAB) m/e (rel intensity) 510 (60), 482 (5), 467 (5), 237 (5); HRMS. Calc'd for M + H: 510.2981. Found: 510.3033.

Example 31




1-phenyl-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-oneStep 1: Preparation of 1-phenyl-4-butyl-1,3-dihydro-2H-imidazol-2-one.

Following General Synthetic Schemes III and IV, 1-phenyl-4-butyl-1,3-dihydro-2H-imidazol-2-one was prepared: NMR (CDCl₃) δ 0.93 (t, J = 7 Hz, 3H), 1.39 (m, J = 7 Hz, 2H), 1.59 (m, J = 7 Hz, 2H), 2.42 (td, J = 8 and 1 Hz, 2H), 6.26 (t, J = 1 Hz, 1H), 7.21 (t, J = 7 Hz, 1H), 7.41 (t, J = 8 Hz, 2H), 7.57-7.62 (m, 2H), 10.28 (br s, 1H); MS (FAB) m/e (rel intensity) 217⁺ (100), 201 (2), 187 (3), 173 (11); HRMS. Calc'd for M + H: 217.1341. Found: 217.1309.

Step 2: Preparation of 1-phenyl-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one.

Following General Synthetic Schemes XII and XIV or XXIII, the imidazol-2-one from Step 1 was converted to 1-phenyl-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one as a colorless solid: NMR (CDCl₃) δ 0.91 (t, J = 7 Hz, 3H), 1.40 (m, J = 7 Hz, 2H), 1.55 (m, J = 7 Hz, 2H), 2.39 (t, J = 7 Hz, 2H), 5.05 (s, 2H), 6.38 (s, 1H), 7.20-7.26 (m, 1H), 7.34-7.56 (m, 6H), 7.59-7.66 (m, 2H), 7.85 (dd, J = 8 and 2 Hz, 1H), 7.97-8.03 (m, 1H), 8.58 (d, J = 2 Hz, 1H); MS (FAB) m/e (rel intensity) 452 (51), 424 (23), 409 (7), 237 (13), 217 (13), 209 (100), 208 (67), 194 (38), 193 (38); HRMS. Calc'd for M + H: 452.2199. Found: 452.2248.

Example 32**1-(2-chlorophenyl)-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one**Step 1: Preparation of 1-(2-chlorophenyl)-4-butyl-1,3-dihydro-2H-imidazol-2-one.


Following General Synthetic Schemes III and IV, 1-(2-chlorophenyl)-4-butyl-1,3-dihydro-2H-imidazol-2-one was prepared: NMR (CDCl₃) δ 0.90 (t, J = 7 Hz, 3H), 1.36 (m, J = 7 Hz, 2H), 1.54 (m, J = 7 Hz, 2H), 2.37 (t, J = 7 Hz, 2H), 6.06 (br s, 1H), 7.23-7.38 (m, 2H), 7.41-7.53 (m, 2H), 10.29 (br s, 1H); MS (FAB) m/e (rel intensity) 252 (36), 250 (100).

Step 2: Preparation of 1-(2-chlorophenyl)-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one.

Following General Synthetic Schemes XII and XIV or XXIII, the imidazol-2-one from Step 1 was converted to 1-(2-chlorophenyl)-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-

imidazol-2-one as a colorless solid: mp 81-83° C; NMR (CDCl₃) δ 0.90 (t, J=7 Hz, 3H), 1.39 (m, J=7 Hz, 2H), 1.54 (m, J=7 Hz, 2H), 2.37 (td, J=8 and 1 Hz, 2H), 5.03 (s, 2H), 6.19 (t, J=1 Hz, 1H), 7.31-7.37 (m, 2H), 7.46-7.63 (m, 6H), 7.88 (dd, J=8 and 2 Hz, 1H), 8.15-8.21 (m, 1H), 8.65 (d, J=2 Hz, 1H); MS (FAB) m/e (rel intensity) 488 (15), 486 (37), 460 (5), 458 (8); HRMS. Calc'd for M+H: 486.1809. Found: 486.1781.

Example 33



1-(2-methylphenyl)-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one


Step 1: Preparation of 1-(2-methylphenyl)-4-butyl-1,3-dihydro-2H-imidazol-2-one.

Following General Synthetic Schemes III and IV, 1-(2-methylphenyl)-4-butyl-1,3-dihydro-2H-imidazol-2-one was prepared: NMR (CDCl₃) δ 0.92 (t, J=7 Hz, 3H), 1.37 (m, J=7 Hz, 2H), 1.55 (m, J=7 Hz, 2H), 2.28 (s, 3H), 2.39 (td, J=7 Hz, 2H), 5.96-5.99 (m, 1H), 7.21-7.31 (m, 4H), 9.88 (br s, 1H); MS (FAB) m/e (rel intensity) 231 (100), 201 (2), 175 (2), 155 (3), 127 (4), 106 (3); HRMS. Calc'd for M+H: 231.1497. Found: 231.1448.

Step 2: Preparation of 1-(2-methylphenyl)-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one.

Following General Synthetic Schemes XII and XIV or XXIII, the imidazol-2-one from Step 1 was converted to 1-(2-methylphenyl)-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one as a colorless solid: NMR (CDCl₃) δ 0.89 (t, J=8 Hz, 3H), 1.31-1.45 (m, 2H), 1.46-1.59 (m, 2H), 2.26 (s, 3H), 2.37 (td, J=8 and 1 Hz, 2H), 5.05 (s, 2H), 6.10 (t, J=1 Hz, 1H), 7.22-7.31 (m, 4H), 7.37 (d, J=8 Hz, 1H), 7.48-7.59 (m, 3H), 7.81 (dd, J=8 and 2 Hz, 1H), 7.98-8.05 (m, 1H), 8.57 (d, J=2 Hz, 1H); MS (FAB) m/e (rel intensity) 466 (64), 438 (23), 237 (25), 231 (33), 209 (100), 194 (37), 193 (32), 180 (83); HRMS. Calc'd for M+H: 466.2355. Found: 466.2397.

Example 34



1-(2-methoxyphenyl)-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one


Step 1: Preparation of 1-(2-methoxyphenyl)-4-butyl-1,3-dihydro-2H-imidazol-2-one.

Following General Synthetic Schemes III and IV, 1-(2-methoxyphenyl)-4-butyl-1,3-dihydro-2H-imidazol-2-one was prepared: NMR (CDCl₃) δ 0.91 (t, J = 7 Hz, 3H), 1.37 (m, J = 7 Hz, 2H), 1.56 (m, J = 7 Hz, 2H), 2.40 (td, J = 8 and 1 Hz, 2H), 3.83 (s, 3H), 6.08-6.12 (m, 1H), 6.96-7.04 (m, 2H), 7.29 (td, J = 8 and 2 Hz, 1H), 7.43 (dd, J = 8 and 2 Hz, 1H), 9.94 (s, 1H); MS (FAB) m/e (rel intensity) 247 (100), 217 (3), 203 (7); HRMS. Calc'd for M + H: 247.1447. Found: 247.1470.

Step 2: Preparation of 1-(2-methoxyphenyl)-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one.

Following General Synthetic Schemes XII and XIV or XXIII, the imidazol-2-one from Step 1 was converted to 1-(2-methoxyphenyl)-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one as a colorless solid: NMR (CDCl₃) δ 0.91 (t, J = 7 Hz, 3H), 1.40 (m, J = 7 Hz, 2H), 1.53 (m, J = 7 Hz, 2H), 2.40 (t, J = 7 Hz, 2H), 3.83 (s, 3H), 5.06 (s, 2H), 6.20 (s, 1H), 6.96-7.03 (m, 2H), 7.27-7.34 (m, 3H), 7.42 (dd, J = 8 and 2 Hz, 1H), 7.53-7.65 (m, 4H), 8.09-8.17 (m, 2H), 8.66 (s, 1H); MS (FAB) m/e (rel intensity) 482 (67), 454 (20), 439 (5), 247 (40), 237 (22), 209 (100), 208 (66), 194 (38), 193 (31); HRMS. Calc'd for M + H: 482.2304. Found: 482.2324.

Example 35



1-(2-isopropylphenyl)-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one


Step 1: Preparation of 1-(2-isopropylphenyl)-4-butyl-1,3-dihydro-2H-imidazol-2-one.

Following General Synthetic Schemes III and IV, 1-(2-isopropylphenyl)-4-butyl-1,3-dihydro-2H-imidazol-2-one was prepared: NMR (CDCl₃) δ 0.89 (t, J = 7 Hz, 3H), 1.21 (d, J = 7 Hz, 6H), 1.30-1.44 (m, 2H), 1.48-1.60 (m, 2H), 2.38 (t, J = 7 Hz, 2H), 3.05 (m, J = 7 Hz, 1H), 5.93-5.96 (m, 1H), 7.17-7.42 (m, 4H), 10.65 (br s, 1H); MS (FAB) m/e (rel intensity) 259 (100); HRMS. Calc'd for M + H: 259.1810. Found: 259.1799.

Step 2: Preparation of 1-(2-isopropylphenyl)-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one.

Following General Synthetic Schemes XII and XIV or XXIII, the imidazol-2-one from Step 1 was converted to 1-(2-isopropylphenyl)-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one as a colorless solid: NMR (CDCl₃) δ 0.89 (t, J = 7 Hz, 3H), 1.21 (d, J = 7 Hz, 6H), 1.37 (m, J = 7 Hz, 2H), 1.52 (m, J = 7 Hz, 2H), 2.36 (t, J = 7 Hz, 2H), 2.98 (m, J = 7 Hz, 1H), 5.07 (s, 2H), 6.07 (s, 1H), 7.09-7.25 (m, 2H), 7.30-7.43 (m, 3H), 7.46-7.55 (m, 3H), 7.81 (dd, J = 8 and 1 Hz, 1H), 7.92-7.99 (m, 1H), 8.53 (d, J = 1 Hz, 1H); MS (FAB) m/e (rel intensity) 494 (75), 466 (26), 451 (4), 259 (14), 237 (37), 209 (100), 208 (61), 194 (37), 193 (32); HRMS. Calc'd for M + H: 494.2668. Found: 494.2651.

Example 36



1-(2,6-difluorophenyl)-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one


Step 1: Preparation of 1-(2,6-difluorophenyl)-4-butyl-1,3-dihydro-2H-imidazol-2-one.

Following General Synthetic Schemes III and IV, 1-(2,6-difluorophenyl)-4-butyl-1,3-dihydro-2H-imidazol-2-one was prepared: NMR (CDCl₃) δ 0.93 (t, J = 7 Hz, 3H), 1.31-1.46 (m, 2H), 1.50-1.62 (m, 2H), 2.41 (td, J = 8 and 1 Hz, 2H), 6.05 (t, J = 1 Hz, 1H), 6.98-7.08 (m, 2H), 7.27-7.39 (m, 1H), 9.18 (br s, 1H).

Step 2: Preparation of 1-(2,6-difluorophenyl)-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one.

Following General Synthetic Schemes XII and XIV or Scheme XXIII, the imidazol-2-one from Step 1 was converted to 1-(2,6-difluorophenyl)-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one as a colorless solid: mp 164.5-165.5° C (dec); NMR (CDCl₃) δ 0.90 (t, J = 8 Hz, 3H), 1.32-1.46 (m, 2H), 1.48-1.60 (m, 2H), 2.37 (td, J = 8 and 1 Hz, 2H), 5.04 (s, 2H), 6.14 (t, J = 1 Hz, 1H), 7.05 (t, J = 8 Hz, 2H), 7.29-7.38 (m, 1H), 7.42 (d, J = 8 Hz, 1H), 7.48-7.58 (m, 3H), 7.81 (dd, J = 8 and 2 Hz, 1H), 8.04-8.11 (m, 1H), 8.60 (d, J = 2 Hz, 1H); MS (FAB) m/e (rel intensity) 488 (83), 460 (26), 253 (10), 237 (25), 209 (100), 194 (37), 193 (35), 180 (80); HRMS. Calc'd for M + H: 488.2010. Found: 488.1951.

Example 37



1-(2,6-dichlorophenyl)-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one


Step 1: Preparation of 1-(2,6-dichlorophenyl)-4-butyl-1,3-dihydro-2H-imidazol-2-one.

Following General Synthetic Schemes III and IV, 1-(2,6-dichlorophenyl)-4-butyl-1,3-dihydro-2H-imidazol-2-one was prepared: NMR (CDCl₃) δ 0.93 (t, J = 7 Hz, 3H), 1.32-1.46 (m, 2H), 1.51-1.62 (m, 2H), 2.43 (td, J = 8 and 1 Hz, 2H), 5.94 (t, J = 1 Hz, 1H), 7.27-7.32 (m, 1H), 7.40-7.46 (m, 2H), 9.33 (br s, 1H).

Step 2: Preparation of 1-(2,6-dichlorophenyl)-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one.

Following General Synthetic Scheme XII and XIV or Scheme XXIII, the imidazol-2-one from Step 1 was converted to 1-(2,6-dichlorophenyl)-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one as a colorless solid: mp 212.0-212.5° C; NMR (CDCl₃) δ 0.88 (t, J = 8 Hz, 3H), 1.30-1.44 (m, 2H), 1.47-1.59 (m, 2H), 2.34 (td, J = 8 and 1 Hz, 2H), 5.10 (s, 2H), 6.04 (t, J = 1 Hz, 1H), 7.28-7.35 (m, 2H), 7.42-7.55 (m, 5H), 7.76 (dd, J = 8 and 2 Hz, 1H), 7.97-8.04 (m, 1H), 8.53 (d, J = 2 Hz, 1H); MS (FAB) m/e (rel intensity) 522 (48), 520 (85), 494 (8), 492 (12), 287 (4), 285 (8), 237 (51), 209 (93), 194 (35), 193 (31), 180 (95); HRMS. Calc'd for M + H: 520.1419. Found: 420.1443.

Example 38



1-(2,6-dimethylphenyl)-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one


Step 1: Preparation of 1-(2,6-dimethylphenyl)-4-butyl-1,3-dihydro-2H-imidazol-2-one.

Following General Synthetic Schemes III and IV, 1-(2,6-dimethylphenyl)-4-butyl-1,3-dihydro-2H-imidazol-2-one was prepared: NMR (CDCl₃) δ 0.91 (t, J = 7 Hz, 3H), 1.29-1.32 (m, 2H), 1.49-1.61 (m, 2H), 2.18 (s, 6H), 2.41 (t, J = 7 Hz, 2H), 5.83-5.86 (m, 1H), 7.07-7.21 (m, 3H), 9.66 (br s, 1H); MS (FAB) m/e (rel intensity) 245 (100), 215 (4), 118 (9); HRMS. Calc'd for M + H: 245.1654. Found: 245.1668.

Step 2: Preparation of 1-(2,6-dimethylphenyl)-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one.


Following General Synthetic Schemes XII and XIV or Scheme XXIII, the imidazol-2-one from Step 1 was converted to 1-(2,6-dimethylphenyl)-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one as a colorless solid: NMR (CDCl₃) δ 0.91 (t, J = 8 Hz, 3H), 1.33-1.47 (m, 2H), 1.49-1.61 (m, 2H), 2.09 (s, 6H), 2.42 (td, J = 8 and 1 Hz, 1H), 5.09 (s, 2H), 6.04 (t, J = 1 Hz, 1H), 7.06-7.11 (m, 2H), 7.14-7.21 (m, 1H), 7.52-7.61 (m, 2H), 7.62-7.69 (m, 2H), 7.96-8.07 (m, 2H), 8.57 (d, J = 2 Hz, 1H); MS (FAB) m/e (rel intensity) 480 (53), 452 (22), 245 (23), 237 (25), 209 (98), 194 (45), 193 (42), 180 (100); HRMS. Calc'd for M + H: 480.2512. Found: 480.2551.

Example 39




1,4-dibutyl-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one

Following General Synthetic Schemes XII and XIII, 1,4-dibutyl-1,3-dihydro-2H-imidazol-2-one (prepared in Step 2A of Example 1) was converted to 1,4-dibutyl-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one as a colorless solid: NMR (CDCl₃) δ 0.85 (t, J = 7 Hz, 3H), 0.88 (t, J = 7 Hz, 3H), 1.11-1.35 (m, 4H), 1.42 (m, J = 7 Hz, 2H), 1.57 (m, J = 7 Hz, 2H), 2.27 (t, J = 7 Hz, 2H), 3.57 (t, J = 7 Hz, 2H), 4.78 (s, 2H), 5.93 (s, 1H), 6.86 (d, J = 8 Hz, 1H), 7.35 (dd, J = 8 and 2 Hz, 1H), 7.42 (d, J = 8 Hz, 1H), 7.52-7.56 (m, 1H), 7.58-7.62 (m, 1H), 7.89 (d, J = 8 Hz, 1H), 8.27 (d, J = 2 Hz, 1H); MS (FAB) m/e (rel intensity) 425 (66), 404 (15), 389 (15), 237 (6), 208 (82), 193 (43), 180 (100); HRMS. Calc'd for M+L: 438.2594. Found: 438.2645. Anal. Calc'd for C₂₄H₂₉N₇O: C, 66.80; H, 6.77; N, 22.72. Found: C, 67.15; H, 6.83; N, 22.94.


Example 40**1,4-dibutyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one**

Following General Synthetic Schemes XXVIII, XXIV, and XX, 1,4-dibutyl-1,3-dihydro-2H-imidazol-2-one (prepared in Step 2A of Example 1) was converted to 1,4-dibutyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one as a colorless solid: mp 178.0-179.5° C; NMR (CDCl₃) δ 0.85 (t, J = 7 Hz, 3H), 0.88 (t, J = 7 Hz, 3H), 1.23-1.49 (m, 6H), 1.50-1.63 (m, 2H), 2.20 (t, J = 7 Hz, 2H), 3.50 (t, J = 7 Hz, 2H), 4.70 (s, 2H), 5.94 (s, 1H), 7.02 (d, J = 8 Hz, 2H), 7.22 (d, J = 8 Hz, 2H), 7.56 (dd, J = 5 and 3 Hz, 1H), 8.31 (d, J = 5 Hz, 1H), 8.77 (d, J = 3 Hz, 1H); MS (FAB) m/e (rel intensity) 432 (21), 389 (4), 237 (8), 209 (28), 208 (100), 194 (20), 180 (14), 153 (7); HRMS. Calc'd for M+H: 432.2512. Found: 432.2558.


Example 41

1,4-dibutyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one

Following General Synthetic Schemes XXVIII, XXV, and XX, 1,4-dibutyl-1,3-dihydro-2H-imidazol-2-one (prepared in Step 2A of Example 1) was converted to 1,4-dibutyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one as a colorless solid: mp 128-130° C; NMR (CDCl₃) δ 0.86 (t, J=7 Hz, 3H), 0.89 (t, J=7 Hz, 3H), 1.21-1.36 (m, 4H), 1.36-1.49 (m, 2H), 2.21 (t, J=7 Hz, 2H), 4.74 (s, 2H), 5.96 (s, 1H), 7.04 (d, J=8 Hz, 2H), 7.11 (d, J=8 Hz, 2H), 7.90 (d, J=5 Hz, 1H), 8.64 (s, 1H), 8.66 (d, J=5 Hz, 1H); MS (FAB) m/e (rel Intensity) 432 (20), 389 (10), 208 (100), 197 (30), 193 (30); HRMS. Calc'd for M+H: 432.2512. Found: 432.2508.

Example 42**1,4-dibutyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one**

Following General Synthetic Schemes XXVIII, XXVI, and XX, 1,4-dibutyl-1,3-dihydro-2H-imidazol-2-one (prepared in Step 2A of Example 1) was converted to 1,4-dibutyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one as a colorless solid: mp 176-178° C (dec); NMR (CDCl₃) δ 0.86 (t, J=7 Hz, 3H), 0.90 (t, J=7 Hz, 3H), 1.22-1.36 (m, 4H), 1.37-1.49 (m, 2H), 1.53-1.66 (m, 2H), 2.22 (t, J=8 Hz, 2H), 3.57 (t, J=8 Hz, 2H), 4.72 (s, 2H), 5.96 (s, 1H), 7.01 (d, J=9 Hz, 2H), 7.09 (d, J=9 Hz, 2H), 7.43 (d, J=5 Hz, 1H), 8.71 (d, J=5 Hz, 1H), 8.87 (s, 1H); MS (FAB) m/e (rel intensity) 432 (22), 404 (3), 389 (5), 237 (22), 208 (100), 193 (34), 180 (75), 167 (18), 153 (22), 141 (10), 128 (6), 111 (12); HRMS. Calc'd. for M+H: 432.2512. Found: 432.2512.

Example 43

1,4-dibutyl-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one

Following General Synthetic Schemes XXVIII, XXVII, and XX, 1,4-dibutyl-1,3-dihydro-2H-imidazol-2-one (prepared in Step 2A of Example 1) was converted to 1,4-dibutyl-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one as a colorless solid: mp 178.0-179.5° C; NMR (CDCl₃) δ 0.88 (t, J = 7 Hz, 3H), 0.93 (t, J = 7 Hz, 3H), 1.25-1.54 (m, 6H), 1.59-1.72 (m, 2H), 2.28 (t, J = 7 Hz, 2H), 3.64 (t, J = 7 Hz, 2H), 4.85 (s, 2H), 5.97 (s, 1H), 7.05 (d, J = 9 Hz, 2H), 7.10 (d, J = 9 Hz, 2H), 7.55 (dd, J = 5 and 3 Hz, 1H), 7.80 (d, J = 5 Hz, 1H), 8.68 (d, J = 3 Hz, 1H); MS (FAB) m/e (rel intensity) 432 (27), 389 (5), 237 (4), 209 (29), 208 (100), 194 (21), 180 (45), 153 (10); HRMS. Calc'd for M + H: 432.2512. Found: 432.2563.

BIOLOGICAL EVALUATION

Assay A: Angiotensin II Binding Activity

Compounds of the invention were tested for ability to bind to the smooth muscle angiotensin II receptor using a rat uterine membrane preparation. Angiotensin II (All) was purchased from Peninsula Labs. ¹²⁵I-angiotensin II (specific activity of 2200 Ci/mmol) was purchased from Du Pont-New England Nuclear. Other chemicals were obtained from Sigma Chemical Co. This assay was carried out according to the method of Douglas et al [Endocrinology, 106, 120-124 (1980)]. Rat uterine membranes were prepared from fresh tissue. All procedures were carried out at 4° C. Uteri were stripped of fat and homogenised in phosphate-buffered saline at pH 7.4 containing 5 mM EDTA. The homogenate was centrifuged at 1500 x g for 20 min., and the supernatant was recentrifuged at 100,000 x g for 60 min. The pellet was resuspended in buffer consisting of 2 mM EDTA and 50 mM Tris-HCl (pH 7.5) to a final protein concentration of 4 mg/ml. Assay tubes were charged with 0.25 ml of a solution containing 5 mM MgCl₂, 2 mM EDTA, 0.5% bovine serum albumin, 50 mM Tris-HCl, pH 7.5 and ¹²⁵I-All (approximately 10⁵ cpm) in the absence or in the presence of unlabelled ligand. The reaction was initiated by the addition of membrane protein and the mixture was incubated at 25° C for 60 min. The incubation was terminated with ice-cold 50 mM Tris-HCl (pH 7.5) and the mixture was filtered to separate membrane-bound labelled peptide from the free ligand. The incubation tube and filter were washed with ice-cold buffer. Filters were assayed for radioactivity in a Micromedic gamma counter. Nonspecific binding was defined as binding in the presence of 10 μ M of unlabelled All. Specific binding was calculated as total binding minus nonspecific binding. The receptor binding affinity of an All antagonist compound was indicated by the concentration (IC₅₀) of the tested All antagonist which gives 50% displacement of the total specifically bound ¹²⁵I-All from the high affinity All receptor. Binding data were analyzed by a nonlinear least-squares curve fitting program. Results are reported in Table I.

Assay B: In Vitro Vascular Smooth Muscle-Response for All

The compounds of the invention were tested for antagonist activity in rabbit aortic rings. Male New Zealand white rabbits (2-2.5 kg) were sacrificed using an overdose of pentobarbital and exsanguinated via the carotid arteries. The thoracic aorta was removed, cleaned of adherent fat and connective tissue and then cut into 3-mm ring segments. The endothelium was removed from the rings by gently sliding a rolled-up piece of filter paper into the vessel lumen. The rings were then mounted in a water-jacketed tissue bath, maintained at 37° C, between moveable and fixed ends of a stainless steel wire with the moveable end attached to an FT03 Grass transducer coupled to a Model 7D Grass Polygraph for recording isometric force responses. The bath was filled with 20 ml of oxygenated (95% oxygen/5% carbon dioxide) Krebs solution of the following composition (mM): 130 NaCl, 15 NaHCO₃, 15 KCl, 1.2 NaH₂PO₄, 1.2 MgSO₄, 2.5 CaCl₂, and 11.4 glucose. The preparations were equilibrated for one hour before approximately one gram of passive tension was placed on the rings. Angiotensin II concentration-response curves were then recorded (3 X 10⁻¹⁰ to 1 X 10⁻⁵ M). Each concentration of All was allowed to elicit its maximal contraction, and then All was washed out repeatedly for 30 minutes before rechallenging with a higher concentration of All. Aorta rings were exposed to the test antagonist at 10⁻⁵ M for 5 minutes before challenging with All. Adjacent segments of the same aorta ring were used for all concentration-response curves in the presence or absence of the test antagonist. The effectiveness of the test compound was expressed in terms of pA₂ values and were calculated according to H.O. Schild [Br. J. Pharmacol. Chemother., 2,189-206 (1947)]. The pA₂ value is the concentration of the antagonist which increases the EC₅₀ value for All by a factor of two. Each test antagonist was evaluated in aorta rings from two rabbits. Results are reported in Table I.

Assay C: In Vivo Intra gastric Pressor Assay Response for All Antagonists

Male Sprague-Dawley rats weighing 225-300 grams were anesthetized with methohexital (30 mg/kg, i.p.) and catheters were implanted into the femoral artery and vein. The catheters were tunneled subcutaneously to exit dorsally, posterior to the head and between the scapulae. The catheters were filled with heparin (1000 units/ml of saline). The rats were returned to their cage and allowed regular rat chow and water ad libitum. After full recovery from surgery (3-4 days), rats were placed in Lucite holders and the arterial line was connected to a pressure transducer. Arterial pressure was recorded on a Gould polygraph (mmHg). Angiotensin II was administered as a 30 ng/kg bolus via the venous catheter delivered in a 50 μ l volume with a 0.2 ml saline flush. The pressor response in mm Hg was measured by the difference from pre-injection arterial pressure to the maximum pressure achieved. The AII injection was repeated every 10 minutes until three consecutive injections yielded responses within 4 mmHg of each other. These three responses were then averaged and represented the control response to AII. The test compound was suspended in 0.5% methylcellulose in water and was administered by gavage. The volume administered was 2 ml/kg body weight. The standard dose was 3 mg/kg. Angiotensin II bolus injections were given at 30, 45, 60, 75, 120, 150, and 180 minutes after gavage. The pressor response to AII was measured at each time point. The rats were then returned to their cage for future testing. A minimum of 3 days was allowed between tests. Percent inhibition was calculated for each time point following gavage by the following formula: $[(\text{Control Response} - \text{Response at time point}) / \text{Control Response}] \times 100$. Results are shown in Table I.

20

25

30

35

40

45

50

55

TABLE I
In Vitro and In Vivo Angiotensin II
Activity of Compounds of the Invention

Test Compound Example #	1 Assay A IC ₅₀ (nM)	2 Assay B pA ₂	3 Assay C Dose: 3 mg/kg (i.g.)	
			Inhibition (%)	Duration (min.)
1	6.5	8.82/8.53	50	> 180
2	38	8.13/7.40	25	180
3	770	7.46/6.95		NA
4	140	7.72/7.09		NA
5	29	8.64/8.23		NA
6	10	7.87/7.89	10	180
7	81	7.75/7.76	10	180
8	140	NA		NA
9	11	9.27/8.87	10	180
10	47	7.64/7.35		NA
11	34	8.44/8.03		NA
12	31	7.68/8.26		NA
13	14	8.03/8.60		NA
14	7.6	8.76/8.64	35	> 180
15	10	8.79/8.85	60	> 180
16	20	8.42/8.77	45	> 180
17	17	8.78/8.63	10	180
18	12	8.79/8.64	65	> 180
19	9.2	8.43/8.36	50	> 180
20	16	9.17/8.86	75	> 180
21	20	9.14/9.15	40	> 180
22	5.4	8.75/8.89	30	> 180
23	99	9.04/8.60		NA
24	22	9.19/8.69	50	> 180
25	5.0	9.41/9.16	25	> 180
26	3.6	8.36/8.44	15	180
27	18	8.74/8.67	35	> 180
28	23	8.85/8.25	15	180
29	51	NA		NA
30	65	NA		NA
31	45	NA		NA
32	5.4	8.80/9.04	50	> 180

TABLE I (CONTINUED)
In Vitro and In Vivo Angiotensin II
Activity of Compounds of the Invention

Test Compound Example #	¹ Assay A IC ₅₀ (nM)	² Assay B pA ₂	³ Assay C Dose: 3mg/kg (i.g.)	
			Inhibition (%)	Duration (min.)
33	9.4	NA	65	> 180
34	9.0	NA		NA
35	14	NA		NA
36	7.0	NA	75	120
37	4.8	NA	25	> 180
38	5.0	NA		NA
39	14	7.45/7.87	20	> 180
40	91	NA		NA
41	160	NA		NA
42	93	NA		NA
43	89	7.55/7.67		NA

¹Assay A: Angiotensin II Binding Activity

²Assay B: In Vitro Vascular Smooth Muscle Response

³Assay C: In Vivo Pressor Response (all test compounds administered intragastrically at 3 mg/kg).

*NA = Not Assayed

Also embraced within this invention is a class of pharmaceutical compositions comprising one or more compounds of Formula I in association with one or more non-toxic, pharmaceutically acceptable carriers and/or diluents and/or adjuvants (collectively referred to herein as "carrier" materials) and, if desired, other active ingredients. The compounds of the present invention may be administered by any suitable route, preferably in the form of a pharmaceutical composition adapted to such a route, and in a dose effective for the treatment intended. Therapeutically effective doses of the compounds of the present invention required to prevent or arrest the progress of the medical condition are readily ascertained by one of ordinary skill in the art. The compounds and composition may, for example, be administered intravascularly, intraperitoneally, subcutaneously, intramuscularly or topically.

For oral administration, the pharmaceutical composition may be in the form of, for example, a tablet, capsule, suspension or liquid. The pharmaceutical composition is preferably made in the form of a dosage unit containing a particular amount of the active ingredient. Examples of such dosage units are tablets or capsules. These may with advantage contain an amount of active ingredient from about 1 to 250 mg, preferably from about 25 to 150 mg. A suitable daily dose for a mammal may vary widely depending on the condition of the patient and other factors. However, a dose of from about 0.1 to 3000 mg/kg body weight, particularly from about 1 to 100 mg/kg body weight, may be appropriate.

The active ingredient may also be administered by injection as a composition wherein, for example, saline, dextrose or water may be used as a suitable carrier. A suitable daily dose is from about 0.1 to 100 mg/kg body weight injected per day in multiple doses depending on the disease being treated. A preferred daily dose would be from about 1 to 30 mg/kg body weight. Compounds indicated for prophylactic therapy will preferably be administered in a daily dose generally in a range from about 0.1 mg to about 100 mg per kilogram of body weight per day. A more preferred dosage will be a range from about 1 mg to about 100 mg per kilogram of body weight. Most preferred is a dosage in a range from about 1 to about 50 mg per

kilogram of body weight per day. A suitable dose can be administered, in multiple sub-doses per day. These sub-doses may be administered in unit dosage forms. Typically, a dose or sub-dose may contain from about 1 mg to about 100 mg of active compound per unit dosage form. A more preferred dosage will contain from about 2 mg to about 50 mg of active compound per unit dosage form. Most preferred is a dosage form containing from about 3 mg to about 25 mg of active compound per unit dose.


The dosage regimen for treating a disease condition with the compounds and/or compositions of this invention is selected in accordance with a variety of factors, including the type, age, weight, sex and medical condition of the patient, the severity of the disease, the route of administration, and the particular compound employed, and thus may vary widely.

For therapeutic purposes, the compounds of this invention are ordinarily combined with one or more adjuvants appropriate to the indicated route of administration. If administered per os, the compounds may be admixed with lactose, sucrose, starch powder, cellulose esters of alkanolic acids, cellulose alkyl esters, talc, stearic acid, magnesium stearate, magnesium oxide, sodium and calcium salts of phosphoric and sulfuric acids, gelatin, acacia gum, sodium alginate, polyvinylpyrrolidone, and/or polyvinyl alcohol, and then tableted or encapsulated for convenient administration. Such capsules or tablets may contain a controlled-release formulation as may be provided in a dispersion of active compound in hydroxypropylmethyl cellulose. Formulations for parenteral administration may be in the form of aqueous or non-aqueous isotonic sterile injection solutions or suspensions. These solutions and suspensions may be prepared from sterile powders or granules having one or more of the carriers or diluents mentioned for use in the formulations for oral administration. The compounds may be dissolved in water, polyethylene glycol, propylene glycol, ethanol, corn oil, cottonseed oil, peanut oil, sesame oil, benzyl alcohol, sodium chloride, and/or various buffers. Other adjuvants and modes of administration are well and widely known in the pharmaceutical art.


Although this invention has been described with respect to specific embodiments, the details of these embodiments are not to be construed as limitations.

Claims

1. A compound of Formula I




wherein A is selected from



wherein m is a number selected from one to four, inclusive;

wherein R¹ is selected from alkyl, hydroxyalkyl, haloalkyl, cycloalkyl, cycloalkylalkyl, formyl, thienylal-


kyl, phenylalkyl, polycycloalkyl, polycycloalkylalkyl, phenyl, halophenyl, alkylphenyl, alkoxyphenyl, cycloalkenyl, cycloalkenylalkyl, aroyl, alkoxyalkyl, alkylcarbonyl, alkylcarbonylalkyl, cycloalkylcarbonyl, cycloalkylalkylcarbonyl, aralkylcarbonyl, alkoxyalkyl, alkenyl, cycloalkenyl, aralkoxycarbonyl, alkynyl, alkylthiocarbonyl, alkylthiothiocarbonyl, arylthiocarbonyl, arylthiothiocarbonyl, aralkylthiocarbonyl, alkylthiocarbonyl, alkylsulfinyl, alkylsulfonyl, aralkylsulfinyl, aralkylsulfonyl, arylsulfinyl, arylsulfonyl, heteroaryl having one or more ring atoms selected from oxygen, sulfur and nitrogen atoms, and amido radicals of the formula




wherein X is oxygen atom or sulfur atom;

wherein each of R¹² and R¹³ is independently selected from hydrido, alkyl, cycloalkyl, cyano, amino, monoalkylamino, dialkylamino, hydroxyalkyl, cycloalkylalkyl, alkoxyalkyl, aralkyl and aryl, and wherein R¹² and R¹³ taken together may form a heterocyclic group having five to seven ring members including the nitrogen atom of said amido radical and which heterocyclic group may further contain one or more hetero atoms as ring members selected from oxygen, nitrogen and sulfur atoms and which heterocyclic group may be saturated or partially unsaturated; wherein each of R¹² and R¹³ taken together may form an aromatic heterocyclic group having five ring members including the nitrogen atom of said amido radical and which aromatic heterocyclic group may further contain one or more additional nitrogen atoms;

wherein each of R⁰ and R² through R¹¹ is independently selected from hydrido, alkyl, hydroxyalkyl, formyl, halo, haloalkyl, cycloalkyl, cycloalkylalkyl, cycloalkylhaloalkyl, cycloalkylcarbonyl, alkoxy, thienylalkyl, phenylalkyl, polycycloalkyl, polycycloalkylalkyl, phenyl, halophenyl, alkylphenyl, alkoxyphenyl, cycloalkenyl, cycloalkenylalkyl, aralkylhaloalkyl, aroyl, aryloxy, aryloxyalkyl, aralkoxy, alkoxyalkyl, alkylcarbonyl, alkylcarbonylalkyl, alkoxyalkyl, alkenyl, cycloalkenyl, alkynyl, cyano, nitro, carboxyl, carboxyalkyl, alkylcarbonyloxy, alkylcarbonyloxyalkyl, alkoxyalkyl, aralkoxyalkyl, aralkylcarbonyloxyalkyl, mercaptocarbonyl, mercaptothiocarbonyl, mercaptoalkyl, alkoxyalkyl, alkylthio, cycloalkylthio, cycloalkylalkylthio, alkylthiocarbonyl, alkylcarbonylthio, alkylthiocarbonyloxy, alkylthiocarbonylthio, alkylthiothiocarbonyl, alkylthiothiocarbonylthio, arylthio, arylthiocarbonyl, arylcarbonylthio, arylthiocarbonyloxy, arylthiocarbonylthio, arylthiothiocarbonyl, arylthiothiocarbonylthio, aralkylthio, aralkylthiocarbonyl, aralkylcarbonylthio, aralkylthiocarbonyloxy, aralkylthiocarbonylthio, alkylthiocarbonyl, aralkylthiocarbonylthio, mercapto, alkylsulfinyl, alkylsulfonyl, aralkylsulfinyl, aralkylsulfonyl, arylsulfinyl, arylsulfonyl, phthalimido, phthalimidoalkyl, heteroaryl, heteroarylalkyl, cycloheteroalkyl, cycloheteroalkylalkyl and cycloheteroalkylcarbonylalkyl wherein each of said heteroaryl- and cyclohetero-containing groups has one or more ring atoms selected from oxygen, sulfur and nitrogen atoms, and wherein each of R⁰ and R² through R¹¹ may be further independently selected from amino and amido radicals of the formula



wherein X is oxygen atom or sulfur atom;




wherein X is oxygen atom or sulfur atom;

wherein each of R^{35} , R^{36} , R^{37} , R^{38} , R^{39} and R^{40} is independently selected from hydrido, alkyl, cycloalkyl, cyano, hydroxyalkyl, cycloalkylalkyl, alkoxyalkyl, haloalkylsulfinyl, haloalkylsulfonyl, aralkyl and aryl, and wherein each of R^{28} and R^{29} taken together and each of R^{30} and R^{31} taken together may form a heterocyclic group having five to seven ring members including the nitrogen atom of said amino or amido radical, which heterocyclic group may further contain one or more hetero atoms as ring members selected from oxygen, nitrogen and sulfur atoms and which heterocyclic group may be saturated or partially unsaturated; wherein each of R^{28} and R^{29} taken together and each of R^{33} and R^{34} taken together may form an aromatic heterocyclic group having five ring members including the nitrogen atom of said amino or amido radical and which aromatic heterocyclic group may further contain one or more additional nitrogen atoms; or a tautomer thereof or a pharmaceutically-acceptable salt thereof.

2. Compound of Claim 1 wherein m is one;


wherein R^1 is selected from alkyl, hydroxyalkyl, haloalkyl, cycloalkyl, cycloalkylalkyl, thienylalkyl, phenylalkyl, polycycloalkyl, polycycloalkylalkyl, phenyl, halophenyl, alkylphenyl, alkoxyphenyl, cycloalkenyl, cycloalkenylalkyl, aroyl, alkoxyalkyl, alkylcarbonyl, cycloalkylcarbonyl, cycloalkylalkylcarbonyl, aralkylcarbonyl, alkoxycarbonyl, alkenyl, cycloalkenyl, alkynyl, cycloalkynyl, alkylthiocarbonyl, arylthiocarbonyl, arylthiothiocarbonyl, aralkylthiocarbonyl, alkylsulfonyl, aralkylsulfonyl, arylsulfonyl, heteroaryl having one or more ring atoms selected from oxygen, sulfur and nitrogen atoms, and amido radicals of the formula



wherein X is oxygen atom or sulfur atom;

wherein each of R^{12} and R^{13} is independently selected from hydrido, alkyl, cycloalkyl, cyano, amino, monoalkylamino, dialkylamino, hydroxyalkyl, cycloalkylalkyl, alkoxyalkyl, aralkyl and aryl;

wherein each of R^0 and R^2 is independently selected from hydrido, alkyl, hydroxyalkyl, formyl, halo, haloalkyl, cycloalkyl, cycloalkylalkyl, cycloalkylhaloalkyl, cycloalkylcarbonyl, alkoxy, thienylalkyl, phenylalkyl, polycycloalkyl, polycycloalkylalkyl, phenyl, halophenyl, alkylphenyl, alkoxyphenyl, cycloalkenyl, cycloalkenylalkyl, aralkylhaloalkyl, aroyl, aryloxy, aryloxyalkyl, aralkoxy, alkoxyalkyl, alkylcarbonyl, alkoxycarbonyl, alkenyl, cycloalkenyl, alkynyl, cycloalkynyl, cyano, nitro, carboxyl, carboxyalkyl, alkylcarbonyloxy, alkylcarbonyloxyalkyl, alkoxycarbonylalkyl, aralkoxycarbonylalkyl, aralkylcarbonyloxyalkyl, mercaptocarbonyl, mercaptothiocarbonyl, mercaptoalkyl, alkoxycarbonyloxy, alkylthio, cycloalkylthio, cycloalkylalkylthio, alkylthiocarbonyl, alkylcarbonylthio, alkylthiocarbonyloxy, alkylthiocarbonylthio, alkylthiothiocarbonyl, alkylthiothiocarbonylthio, arylthio, arylthiocarbonyl, arylcarbonylthio, arylthiocarbonyloxy, arylthiocarbonylthio, arylthiothiocarbonyl, arylthiothiocarbonylthio, aralkylthio, aralkylthiocarbonyl, aralkylcarbonylthio, aralkylthiocarbonyloxy, aralkylthiocarbonylthio, aralkylthiocarbonyl, aralkylthiocarbonylthio, mercapto, alkylsulfinyl, alkylsulfonyl, aralkylsulfinyl, aralkylsulfonyl, arylsulfinyl, arylsulfonyl, phthalimido, phthalimidoalkyl, heteroaryl, heteroarylalkyl, cycloheteroalkyl, cycloheteroalkylalkyl and cycloheteroalkylcarbonylalkyl wherein each of said heteroaryl- and cycloheteroalkyl-containing groups has one or more hetero ring atoms selected from oxygen, sulfur and nitrogen atoms, and wherein each of R^0 and R^2 through R^{11} may be further independently selected from amino and amido radicals of the formula




wherein X is selected from oxygen atom or sulfur atom;

wherein each n is a number independently selected from zero to six, inclusive;

wherein each of R¹⁴ through R²⁶ is independently selected from hydrido, alkyl, cycloalkyl, cyano, amino, monoalkylamino, dialkylamino, hydroxyalkyl, cycloalkylalkyl, alkoxyalkyl, aralkyl and aryl;

wherein each of R³ through R¹¹ is independently selected from hydrido, hydroxy, alkyl, hydroxyalkyl, halo, haloalkyl, cycloalkyl, cycloalkylalkyl, alkoxy, aralkyl, aryl, aroyl, aryloxy, aralkoxy, alkoxyalkyl, alkylcarbonyl, alkoxy carbonyl, alkenyl, cycloalkenyl, alkynyl, cycloalkynyl, cyano, nitro, carboxyl, alkylcarbonyloxy, mercaptocarbonyl, mercaptothiocarbonyl, alkoxy carbonyloxy, alkylthio, alkylthiocarbonyl, alkylcarbonylthio, alkylthiocarbonyloxy, alkylthiocarbonylthio, alkylthiothiocarbonyl, arylthio, arylthiocarbonyl, arylcarbonylthio, arylthiocarbonyloxy, arylthiothiocarbonyl, aralkylthio, aralkylthiocarbonyl, aralkylcarbonylthio, aralkylthiocarbonyloxy, aralkylthiocarbonylthio, aralkylthiocarbonyl, aralkylthiocarbonylthio, mercapto, alkylsulfonyl, aralkylsulfonyl and arylsulfonyl, and amino and amido radicals of the formula



wherein X is oxygen atom or sulfur atom;


wherein each of R¹⁴, R¹⁵, R¹⁶, R¹⁷, R¹⁸ and R¹⁹ is independently selected from hydrido, alkyl, cycloalkyl, cyano, hydroxyalkyl, cycloalkylalkyl, alkoxyalkyl, aralkyl and aryl;

and wherein each of R⁰ and R³ through R¹¹ may be further independently selected from acidic moieties of the formula


-Y_nA

wherein n is a number selected from zero through three, inclusive; wherein A is an acidic group selected from acids containing one or more atoms selected from oxygen, sulfur, phosphorus and nitrogen atoms, and wherein said acidic group is selected to contain at least one acidic hydrogen atom, and the amide, ester and salt derivatives of said acidic moieties; wherein Y is a spacer group independently selected from one or more of alkyl, cycloalkyl, cycloalkylalkyl, alkenyl, alkynyl, aryl, aralkyl and heteroaryl having one or more ring atoms selected from oxygen, sulfur and nitrogen atoms;

and wherein any of the foregoing R⁰ through R²⁶, Y and A groups having a substitutable position may be substituted by one or more groups independently selected from hydroxy, alkyl, alkenyl, aralkyl, hydroxyalkyl, halo, haloalkyl, oxo, alkoxy, aryloxy, aralkoxy, alkoxyalkyl, alkylcarbonyl, alkoxy carbonyl, carboxyl, cyano, nitro, alkylsulfonyl, haloalkylsulfonyl, aryl, aralkyl, mercaptocarbonyl, alkylthio and alkylthiocarbonyl, and amino and amido radicals of the formula



wherein X is oxygen atom or sulfur atom; wherein each of R²⁷ through R³¹ is independently selected from hydrido, alkyl, cycloalkyl, cycloalkylalkyl, aralkyl, aryl, and DR³² and




wherein D is selected from oxygen atom and sulfur atom, and R³² is selected from hydrido, alkyl, cycloalkyl, cycloalkylalkyl, aralkyl and aryl; wherein each of R²⁷, R²⁸, R²⁹, R³⁰, R³¹, R³³ and R³⁴ is independently selected from hydrido, alkyl, cycloalkyl, cyano, hydroxyalkyl, haloalkyl, cycloalkylalkyl, alkoxyalkyl, alkanoyl, alkoxycarbonyl, carboxyl, haloalkylsulfinyl, haloalkylsulfonyl, aralkyl and aryl;

or a tautomer thereof or a pharmaceutically-acceptable salt thereof.


3. Compound of Claim 2 wherein m is one;

wherein R¹ is selected from alkyl, hydroxyalkyl, haloalkyl, cycloalkyl, cycloalkylalkyl, thienylalkyl, phenylalkyl, polycycloalkyl, polycycloalkylalkyl, phenyl, halophenyl, alkylphenyl, alkoxyphenyl, cycloalkenyl, cycloalkenylalkyl, aroyl, alkoxyalkyl, alkylcarbonyl, cycloalkylcarbonyl, cycloalkylalkylcarbonyl, aralkylcarbonyl, alkoxycarbonyl, alkenyl, cycloalkenyl, alkynyl, cycloalkynyl, alkylsulfonyl, aralkylsulfonyl and arylsulfonyl, and amido radicals of the formula



wherein each of R¹² and R¹³ is independently selected from hydrido, alkyl, cycloalkyl, cyano, amino, monoalkylamino, dialkylamino, hydroxyalkyl, cycloalkylalkyl, alkoxyalkyl, aralkyl and aryl;

wherein each of R⁰ and R² is independently selected from hydrido, alkyl, hydroxyalkyl, formyl, halo, haloalkyl, cycloalkyl, cycloalkylalkyl, cycloalkylhaloalkyl, cycloalkylcarbonyl, alkoxy, thienylalkyl, phenylalkyl, polycycloalkyl, polycycloalkylalkyl, phenyl, halophenyl, alkylphenyl, alkoxyphenyl, cycloalkenyl, cycloalkenylalkyl, aralkylhaloalkyl, aroyl, aryloxy, aryloxyalkyl, aralkoxy, alkoxyalkyl, alkylcarbonyl, alkoxycarbonyl, alkenyl, cycloalkenyl, alkynyl, cycloalkynyl, cyano, nitro, carboxyl, carboxyalkyl, alkylcarbonyloxy, mercaptocarbonyl, alkoxycarbonyloxy, alkylcarbonyloxyalkyl, alkoxycarbonylalkyl, aralkoxycarbonylalkyl, aralkylcarbonyloxyalkyl, alkylthio, cycloalkylthio, cycloalkylalkylthio, arylthio, aralkylthio, aralkylthiocarbonylthio, mercapto, alkylsulfinyl, alkylsulfonyl, aralkylsulfinyl, aralkylsulfonyl, arylsulfinyl, arylsulfonyl, phthalimido, phthalimidoalkyl, heteroaryl, heteroarylalkyl, cycloheteroalkyl, cycloheteroalkylalkyl and cycloheteroalkylcarbonylalkyl wherein each of said heteroaryl- and cycloheteroalkyl-containing groups has one or more hetero ring atoms selected from oxygen, sulfur and nitrogen atoms, and wherein each of R⁰ and R² may be further independently selected from amino and amido radicals of the formula




wherein X is selected from oxygen atom or sulfur atom;

wherein each n is a number independently selected from zero to six, inclusive;

wherein each of R¹⁴ through R²⁶ is independently selected from hydrido, alkyl, cycloalkyl, cyano, amino, monoalkylamino, dialkylamino, hydroxyalkyl, cycloalkylalkyl, alkoxyalkyl, aralkyl and aryl;

wherein each of R³ through R¹¹ is independently selected from hydrido, hydroxy, alkyl, hydroxyalkyl, halo, haloalkyl, cycloalkyl, alkoxy, aralkyl, aryl, aroyl, aryloxy, aralkoxy, alkoxyalkyl, alkylcarbonyl, alkoxy carbonyl, alkenyl, cycloalkenyl, alkynyl, cyano, nitro, carboxyl, alkylcarbonyloxy, mercaptocarbonyl, alkoxy carbonyloxy, alkylthio, arylthio, aralkylthio, mercapto, alkylsulfonyl, aralkylsulfonyl and arylsulfonyl, and amino and amido radicals of the formula





wherein each of R¹⁴, R¹⁵, R¹⁶, R¹⁷, R¹⁸ and R¹⁹ is independently selected from hydrido, alkyl, cycloalkyl, cyano, amino, monoalkylamino, dialkylamino, hydroxyalkyl, cycloalkylalkyl, alkoxyalkyl, aralkyl and aryl;


and wherein each of R⁰ and R³ through R¹¹ may be further independently selected from acidic moieties of the formula




wherein n is a number selected from zero through three, inclusive;

wherein A is selected from carboxylic acid and bioisosteres of carboxylic acid selected from





wherein each W is independently selected from oxygen atom, sulfur atom and NR^{39} ; wherein each of R^{35} , R^{36} , R^{37} , R^{38} and R^{39} is independently selected from hydrido, alkyl, haloalkyl, haloalkylsulfonyl, haloalkylcarbonyl, cycloalkyl, cycloalkylalkyl, aryl and aralkyl; wherein each of R^{35} , R^{36} , R^{37} and R^{39} may be further independently selected from amino radical of the formula




wherein each of R^{40} and R^{41} is independently selected from hydrido, alkyl, cycloalkyl, hydroxyalkyl, haloalkyl, cycloalkylalkyl, alkoxyalkyl, aralkyl and aryl, and wherein R^{40} and R^{41} taken together may form a heterocyclic group having five to seven ring members including the nitrogen atom of said amino radical, which heterocyclic group may further contain one or more hetero atoms as ring members selected from oxygen, nitrogen and sulfur atoms and which heterocyclic group may be saturated or partially unsaturated; wherein R^{40} and R^{41} taken together may form an aromatic heterocyclic group having five ring members including the nitrogen atom of said amino radical and which aromatic heterocyclic group may further contain one or more hetero atoms as ring atoms selected from oxygen, nitrogen and sulfur atoms; wherein each of R^{36} and R^{37} may be further independently selected from hydroxy, alkoxy, alkylthio, aryloxy, arylthio, aralkylthio and aralkoxy; and the amide, ester and salt derivatives of said acidic groups;


wherein said bioisostere of carboxylic acid may be further selected from heterocyclic acidic groups consisting of heterocyclic rings of four to about nine ring members, which heterocyclic ring contains at least one hetero atom selected from oxygen, sulfur and nitrogen atoms, which heterocyclic ring may be saturated, fully unsaturated or partially unsaturated, and which heterocyclic ring may be attached at a single position selected from R^3 through R^{11} or may be attached at any two adjacent positions selected from R^3 through R^{11} so as to form a fused-ring system with one of the phenyl rings of Formula I; and the amide, ester and salt derivatives of said heterocyclic acidic groups;

wherein Y is a spacer group independently selected from one or more of alkyl, cycloalkyl, cycloalkylalkyl, alkenyl, aryl and aralkyl;

and wherein any of the foregoing R^0 through R^{26} and R^{35} through R^{41} , Y and A groups having a substitutable position may be substituted by one or more groups independently selected from hydroxy, alkyl, alkenyl, aralkyl, hydroxyalkyl, halo, oxo, haloalkyl, alkoxy, aryloxy, aralkoxy, alkoxyalkyl, alkylcarbonyl, alkoxy carbonyl, carboxyl, cyano, nitro, alkylsulfonyl, haloalkylsulfonyl, aryl, aralkyl, mercaptocarbonyl, alkylthio and alkylthiocarbonyl, and amino and amido radicals of the formula



wherein X is selected from oxygen atom and sulfur atom; wherein R^{27} is selected from hydrido, alkyl, cycloalkyl, cycloalkylalkyl, aralkyl, aryl and DR^{32} and




wherein D is selected from oxygen atom and sulfur atom; wherein R³² is selected from hydrido, alkyl, cycloalkyl, cycloalkylalkyl, aralkyl and aryl;

wherein each of R²⁷, R²⁸, R²⁹, R³⁰, R³¹, R³³ and R³⁴ is independently selected from hydrido, alkyl, cycloalkyl, cyano, hydroxyalkyl, haloalkyl, cycloalkylalkyl, alkoxyalkyl, alkanoyl, alkoxy carbonyl, carboxyl, haloalkylsulfinyl, haloalkylsulfonyl, aralkyl and aryl;


or a tautomer thereof or a pharmaceutically-acceptable salt thereof.

4. Compound of Claim 3 wherein m is one;
 wherein R¹ is selected from alkyl, hydroxyalkyl, haloalkyl, cycloalkyl, cycloalkylalkyl, thienylalkyl, phenylalkyl, polycycloalkyl, polycycloalkylalkyl, phenyl, halophenyl, alkylphenyl, alkoxyphenyl, cycloalkenyl, cycloalkenylalkyl, aroyl, alkoxyalkyl, alkylcarbonyl, cycloalkylcarbonyl, cycloalkylalkylcarbonyl, aralkylcarbonyl, alkenyl, cycloalkenyl, alkynyl, mercaptocarbonyl, alkylsulfonyl, aralkylsulfonyl, arylsulfonyl and amido radicals of the formula



wherein each of R¹² and R¹³ is independently selected from hydrido, alkyl, cycloalkyl, cyano, amino, monoalkylamino, dialkylamino, hydroxyalkyl, cycloalkylalkyl, alkoxyalkyl, aralkyl and aryl;

wherein each of R⁰ and R² is independently selected from hydrido, alkyl, hydroxyalkyl, formyl, halo, haloalkyl, cycloalkyl, cycloalkylalkyl, cycloalkylhaloalkyl, cycloalkylcarbonyl, alkoxy, thienylalkyl, phenylalkyl, polycycloalkyl, polycycloalkylalkyl, phenyl, halophenyl, alkylphenyl, alkoxyphenyl, cycloalkenyl, cycloalkenylalkyl, aralkylhaloalkyl, aroyl, aryloxy, aryloxyalkyl, aralkoxy, alkoxyalkyl, alkylcarbonyl, alkoxy carbonyl, alkenyl, cycloalkenyl, alkynyl, cycloalkynyl, cyano, nitro, carboxyl, carboxyalkyl, alkylcarbonyloxy, alkylcarbonyloxyalkyl, alkoxy carbonylalkyl, aralkoxy carbonylalkyl, aralkylcarbonyloxyalkyl, mercaptocarbonyl, mercaptoalkyl, alkoxy carbonyloxy, alkylthio, cycloalkylthio, cycloalkylalkylthio, arylthio, aralkylthio, mercapto, alkylsulfinyl, alkylsulfonyl, aralkylsulfinyl, aralkylsulfonyl, arylsulfinyl, arylsulfonyl, phthalimido, phthalimidoalkyl, heteroaryl, heteroarylalkyl, cycloheteroalkyl, cycloheteroalkylalkyl and cycloheteroalkylcarbonylalkyl wherein each of said heteroaryl- and cycloheteroalkyl-containing groups has one or more hetero ring atoms selected from oxygen, sulfur and nitrogen atoms, and wherein each of R² through R¹¹ may be further independently selected from amino and amido radicals of the formula



wherein X is selected from oxygen atom and sulfur atom;

wherein each n is a number independently selected from zero to six, inclusive;

5 wherein each of R¹⁴ through R²⁶ is independently selected from hydrido, alkyl, cycloalkyl, cyano, amino, monoalkylamino, dialkylamino, hydroxyalkyl, cycloalkylalkyl, alkoxyalkyl, aralkyl and aryl;


wherein each of R³ through R¹¹ is independently selected from hydrido, hydroxy, alkyl, hydroxyalkyl, halo, haloalkyl, cycloalkyl, alkoxy, aralkyl, aryl, aroyl, aryloxy, aralkoxy, alkoxyalkyl, alkylcarbonyl, alkoxyalkyl, alkenyl, cycloalkenyl, alkynyl, cyano, nitro, carboxyl, alkylthio, aralkylthio and mercapto;

and wherein each of R⁰ and R³ through R¹¹ may be further independently selected from acidic moieties of the formula

15 -Y_nA

wherein n is a number selected from zero through three, inclusive; wherein A is selected from carboxylic acid and bioisosteres of carboxylic acid selected from


20 -OH, -SH,



25

30 -S-WH,


30



35

wherein each W is independently selected from oxygen atom, sulfur atom and NR³⁹; wherein each of R³⁵, R³⁸ and R³⁹ is independently selected from hydrido, alkyl, haloalkyl, haloalkylsulfonyl, haloalkylcarbonyl, cycloalkyl, cycloalkylalkyl, aryl and aralkyl; wherein each of R³⁵ and R³⁹ may be further independently selected from amino radical of the formula

40



45

wherein each of R⁴⁰ and R⁴¹ is independently selected from hydrido, alkyl, cycloalkyl, hydroxyalkyl, haloalkyl, cycloalkylalkyl, alkoxyalkyl, aralkyl and aryl, and wherein R⁴⁰ and R⁴¹ taken together may form a heterocyclic group having five to seven ring members including the nitrogen atom of said amino radical, which heterocyclic group may further contain one or more hetero atoms as ring members selected from oxygen, nitrogen and sulfur atoms, and which heterocyclic group may be saturated or partially unsaturated; wherein R⁴⁰ and R⁴¹ taken together may form an aromatic heterocyclic group having five ring members including the nitrogen atom of said amino radical and which aromatic heterocyclic group may further contain one or more hetero atoms as ring atoms selected from oxygen, nitrogen and sulfur atoms; and the amide, ester and salt derivatives of said acidic groups; wherein said bioisostere of carboxylic acid may be further selected from heterocyclic acidic groups consisting of

55

heterocyclic rings of four to about nine ring members, which ring contains at least one hetero atom, selected from oxygen, sulfur and nitrogen atoms, which heterocyclic ring may be saturated, fully unsaturated or partially unsaturated, and which heterocyclic ring may be attached at a single position selected from R³ through R¹¹ or may be attached at any two adjacent positions selected from R³ through R¹¹ so as to form a fused-ring system with one of the phenyl rings of Formula I; and the amide, ester and salt derivatives of said heterocyclic acidic groups;


wherein Y is a spacer group independently selected from one or more of alkyl, cycloalkyl, cycloalkylalkyl, alkenyl, aryl and aralkyl;

wherein each of R⁰ through R²⁶, R³⁵ and R³⁸ through R⁴¹, Y and A independently may be substituted at any substitutable position with one or more groups selected from alkyl, hydroxy, halo, oxo, haloalkyl, alkoxycarbonyl, cyano, nitro, alkylsulfonyl, haloalkylsulfonyl, aryl, aralkyl, alkoxy, aryloxy and aralkoxy;

or a tautomer thereof or a pharmaceutically-acceptable salt thereof.


5. Compound of Claim 4 wherein m is one;

wherein R¹ is selected from alkyl, hydroxyalkyl, haloalkyl, cycloalkyl, cycloalkylalkyl, thienylalkyl, phenylalkyl, polycycloalkyl, polycycloalkylalkyl, phenyl, halophenyl, alkylphenyl, alkoxyphenyl, cycloalkenyl, cycloalkenylalkyl, aroyl, alkoxyalkyl, alkylcarbonyl, cycloalkylcarbonyl, cycloalkylalkylcarbonyl, aralkylcarbonyl, alkenyl, alkynyl, alkylsulfonyl, aralkylsulfonyl, arylsulfonyl and amido radicals of the formula



wherein each of R¹² and R¹³ is independently selected from hydrido, alkyl, cycloalkyl, cyano, amino, hydroxyalkyl, alkoxyalkyl, phenalkyl and phenyl;

wherein each of R⁰ and R² is independently selected from hydrido, alkyl, hydroxyalkyl, formyl, halo, haloalkyl, cycloalkyl, cycloalkylalkyl, cycloalkylhaloalkyl, cycloalkylcarbonyl, alkoxy, thienylalkyl, phenylalkyl, polycycloalkyl, polycycloalkylalkyl, phenyl, halophenyl, alkylphenyl, alkoxyphenyl, cycloalkenyl, cycloalkenylalkyl, aralkylhaloalkyl, benzoyl, phenoxy, phenoxyalkyl, phenalkyloxy, phenylthio, phenalkylthio, aralkoxy, alkoxyalkyl, alkylcarbonyl, alkoxycarbonyl, alkenyl, cycloalkenyl, alkynyl, cyano, nitro, carboxyl, carboxyalkyl, alkylcarbonyloxy, alkylcarbonyloxyalkyl, alkoxycarbonylalkyl, aralkoxycarbonylalkyl, aralkylcarbonyloxyalkyl, mercaptocarbonyl, mercaptoalkyl, alkoxycarbonyloxy, alkylthio, cycloalkylthio, cycloalkylalkylthio, phthalimido, phthalimidoalkyl, heteroaryl, heteroarylalkyl, cycloheteroalkyl, cycloheteroalkylalkyl and cycloheteroalkylcarbonylalkyl wherein each of said heteroaryl- and cycloheteroalkyl-containing groups has one or more hetero ring atoms selected from oxygen, sulfur and nitrogen atoms, and wherein each of R⁰ and R² through R¹¹ may be further independently selected from amino and amido radicals of the formula



wherein X is selected from oxygen atom and sulfur atom;

wherein each n is a number independently selected from zero to six, inclusive;

5 wherein each of R¹⁴ through R²⁶ is independently selected from hydrido, alkyl, cycloalkyl, cyano, amino, hydroxyalkyl, alkoxyalkyl, phenalkyl and phenyl;


wherein each of R³ through R¹¹ is independently selected from hydrido, hydroxy, alkyl, hydroxyalkyl, halo, haloalkyl, cycloalkyl, alkoxy, phenalkyl, phenyl, benzoyl, phenoxy, phenalkyloxy, alkoxyalkyl, 10 alkylcarbonyl, alkoxycarbonyl, alkenyl, cyano, nitro, carboxyl, alkylthio and mercapto;

and wherein each of R⁰ and R³ through R¹¹ may be further independently selected from acidic moieties of the formula

15 -Y_nA

wherein n is a number selected from zero through two, inclusive; wherein A is selected from carboxylic acid and bioisosteres of carboxylic acid selected from


20 -OH, -SH,



25

-S-WH,


30



35

wherein each W is independently selected from oxygen atom, sulfur atom and NR³⁹; wherein each of R³⁵, R³⁸ and R³⁹ is independently selected from hydrido, alkyl, haloalkyl, haloalkylsulfonyl, haloalkylcarbonyl, cycloalkyl, phenyl and benzyl; wherein each of R³⁵ and R³⁹ may be further independently 40 selected from amino radical of the formula

40



45

wherein each of R⁴⁰ and R⁴¹ is independently selected from hydrido, alkyl, cycloalkyl, hydroxyalkyl, haloalkyl, alkoxyalkyl, benzyl and phenyl; and the amide, ester and salt derivatives of said acidic groups;

50

wherein said bioisostere of carboxylic acid may be further selected from heterocyclic acidic groups consisting of heterocyclic rings of four to about nine ring members, which ring contains at least one hetero atom, selected from oxygen, sulfur and nitrogen atoms, which heterocyclic ring may be saturated, fully unsaturated or partially unsaturated, and which heterocyclic ring may be attached at a 55 single position selected from R³ through R¹¹ or may be attached at any two adjacent positions selected from R³ through R¹¹ so as to form a fused-ring system with one of the phenyl rings of Formula I; and the amide, ester and salt derivatives of said heterocyclic acidic groups;

wherein Y is a spacer group independently selected from one or more of alkyl, cycloalkyl, cycloalkylalkyl, alkenyl, phenyl, phenalkyl and aralkyl;


wherein each of R⁰ through R²⁶, R³⁵ and R³⁸ through R⁴¹, Y and A and independently may be substituted at any substitutable position with one or more groups selected from alkyl, cycloalkyl, cycloalkylalkyl, hydroxy, halo, oxo, haloalkyl, alkoxycarbonyl, cyano, nitro, alkylsulfonyl, haloalkylsulfonyl, aryl, aralkyl, alkoxy, aryloxy and aralkoxy;

or a tautomer thereof or a pharmaceutically-acceptable salt thereof.

6. Compound of Claim 5 wherein m is one;

wherein R¹ is selected from alkyl, hydroxyalkyl, haloalkyl, cycloalkyl, cycloalkylalkyl, thienylalkyl, phenylalkyl, polycycloalkyl, polycycloalkylalkyl, phenyl, halophenyl, alkylphenyl, alkoxyphenyl, cycloalkenyl, cycloalkenylalkyl, benzoyl, alkoxyalkyl, alkylcarbonyl, alkoxy carbonyl, cycloalkylcarbonyl, cycloalkylalkylcarbonyl, aralkylcarbonyl, alkenyl and alkynyl;

where each of R⁰ and R² is independently selected from hydrido, alkyl, aminoalkyl, hydroxyalkyl, formyl, halo, haloalkyl, cycloalkyl, cycloalkylalkyl, cycloalkylhaloalkyl, cycloalkylcarbonyl, alkoxy, thienylalkyl, phenylalkyl, polycycloalkyl, polycycloalkylalkyl, phenyl, halophenyl, alkylphenyl, alkoxyphenyl, cycloalkenyl, cycloalkenylalkyl, aralkylhaloalkyl, benzoyl, phenoxy, phenoxyalkyl, phenalkyloxy, phenylthio, phenalkylthio, aralkoxy, alkoxyalkyl, acetyl, alkoxycarbonyl, alkenyl, cycloalkenyl, alkynyl, cyano, nitro, carboxyl, carboxyalkyl, alkylcarbonyl, alkylcarbonyloxy, mercaptoalkyl, mercaptocarbonyl, alkoxycarbonyloxy, alkylcarbonyloxyalkyl, alkoxycarbonylalkyl, aralkoxycarbonylalkyl, aralkylcarbonyloxyalkyl, phthalimido, phthalimidoalkyl, imidazoalkyl, tetrazole, tetrazolealkyl, alkylthio, cycloalkylthio, cycloalkylalkylthio, and amino and amido radicals of the formula




wherein X is selected from oxygen atom and sulfur atom;


wherein each n is a number independently selected from zero to six, inclusive;

wherein each of R¹⁴ through R²⁶ is independently selected from hydrido, alkyl, cycloalkyl, cyano, amino, hydroxyalkyl, alkoxyalkyl, phenalkyl and phenyl;

wherein each of R³ through R¹¹ is independently selected from hydrido, hydroxy, alkyl, hydroxyalkyl, halo, haloalkyl, alkoxy, phenyl, benzoyl, phenoxy, alkoxyalkyl, acetyl, alkoxycarbonyl, alkenyl, cyano, nitro, carboxyl, alkylthio and mercapto;

and wherein each of R⁰ and R³ through R¹¹ may be further independently selected from acidic moieties consisting of CO₂H, CO₂CH₃, SH, CH₂SH, C₂H₄SH, PO₃H₂, NHSO₂CF₃, NHSO₂C₆F₅, SO₃H, CONHNH₂, CONHNHSO₂CF₃, CONHOCH₃, CONHOC₂H₅, CONHCF₃, OH, CH₂OH, C₂H₄OH, OPO₃H₂, OSO₃H.






and the esters, amides and salts of said acidic moieties; or a tautomer thereof or a pharmaceutically-acceptable salt thereof.

7. Compound of Claim 6 wherein m is one;

35 wherein R¹ is selected from methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, n-pentyl, isopentyl, neopentyl, phenyl, benzyl, phenethyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cyclopropylmethyl, cyclopropylethyl, cyclobutylmethyl, cyclobutylethyl, cyclopentylmethyl, cyclopentylethyl, cyclohexylmethyl, cyclohexylethyl, cyclohexanoyl, methylbutyl, ethylbutyl, dimethylbutyl, thienylmethyl, thienylethyl, thienylpropyl, cyclopentenylmethyl, cyclopentenylethyl, cyclopentenylpropyl, cyclohexenyl, cyclohexenylmethyl, cyclohexenylethyl, adamantyl, adamantylmethyl, adamantylethyl, phenyl, chlorophenyl, dichlorophenyl, fluorophenyl, difluorophenyl, methoxyphenyl, ethoxyphenyl, methylphenyl, ethylphenyl, propylphenyl, isopropylphenyl, dimethylphenyl, diethylphenyl, benzyl, 1-oxo-2-cyclohexylethyl, benzoyl, 1-oxo-2-phenethyl, 1-oxoethyl, 1-oxopropyl, 1-oxobutyl, 1-oxopentyl, 2-butanyl, 3-butanyl, 2-butenyl, 3-butenyl and 2-hydroxybutyl; wherein R⁰ is selected from hydrido, C₄H₉-(n), CH₃CH₂CH=CH, C₃H₇(n), SC₃H₇,




C₂H₅, C₅H₁₁(n), C₆H₁₃(n), SC₄H₉,




5 $\text{CH}_3\text{CH}=\text{CH}$, $\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}=\text{CH}$ -, amino, aminomethyl, aminoethyl, aminopropyl, CH_2OH , $\text{CH}_2\text{OCOCH}_3$, CH_2Cl , CH_2OCH_3 , $\text{CH}_2\text{OCH}(\text{CH}_3)_2$, CHO , $\text{CH}_2\text{CO}_2\text{H}$, $\text{CH}(\text{CH}_3)\text{CO}_2\text{H}$, NO_2 , Cl ,

10




15



20


$-\text{CO}_2\text{CH}_3$, $-\text{CONH}_2$, $-\text{CONHCH}_3$, $\text{CON}(\text{CH}_3)_2$, $-\text{CH}_2-\text{NHCO}_2\text{C}_2\text{H}_5$,

25




30 $-\text{CH}_2\text{NHCO}_2\text{CH}_3$, $-\text{CH}_2\text{NHCO}_2\text{C}_3\text{H}_7$, $-\text{CH}_2\text{NHCO}_2\text{CH}_2(\text{CH}_3)_2$, $-\text{CH}_2\text{NHCO}_2\text{C}_4\text{H}_9$, CH_2NHCO_2 -adamantyl, $-\text{CH}_2\text{NHCO}_2$ -(1-naphthyl), $-\text{CH}_2\text{NHCONHCH}_3$, $-\text{CH}_2\text{NHCONHC}_2\text{H}_5$, $-\text{CH}_2\text{NHCONHC}_3\text{H}_7$, $-\text{CH}_2\text{NHCONHC}_4\text{H}_9$, $-\text{CH}_2\text{NHCONHCH}(\text{CH}_3)_2$, $-\text{CH}_2\text{NHCONH}(1\text{-naphthyl})$, $-\text{CH}_2\text{NHCONH}(1\text{-adamantyl})$,

35




$-\text{CH}_2\text{CH}_2\text{CH}_2\text{CO}_2\text{H}$, $-\text{CH}_2\text{CH}_2\text{F}$, $-\text{CH}_2\text{OCONHCH}_3$, $-\text{CH}_2\text{OCSNHCH}_3$, $-\text{CH}_2\text{NHCOC}_3\text{H}_7$, $-\text{CH}_2\text{CH}_2\text{CH}_2\text{F}$, $-\text{CH}_2\text{ONO}_2$,

40



45

$-\text{CH}_2\text{SH}$,




50

55 Cl , NO_2 , CF_3 , CH_2OH , Br , F , I , methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, n-pentyl, isopentyl, neopentyl, phenyl, cyclohexyl, cyclohexylmethyl, carboxyl, formyl, 1-oxoethyl, 1-oxopropyl, 1-oxobutyl, 1-oxopentyl, dimethoxymethyl, 1,1-dimethoxypropyl, 1,1-dimethoxypentyl, hydroxyalkyl, halo, 1-oxo-2-phenylethyl, 1-oxo-2-cyclohexylethyl, 1,1-difluoro-2-phenylethyl, monofluoromethyl, 1,1-difluoro-2-cyclohexylethyl, 2-cyclohexylethyl, 1,1-difluoro-3-cyclohexylpropyl, 1,1-dimethoxybutyl, 1,1-difluoroethyl, 1,1-difluoropropyl, 1,1-difluorobutyl, 1,1-difluoropentyl, benzyl, 2-phenylethyl, 1,1-difluoro-3-phenylpropyl, cyclohexylmethyl, cyclohexylethyl, cyclohexanoyl, 1-butenyl, 2-butenyl, 3-butenyl, 1-butyne, 2-butyne, 3-butyne, difluoromethyl, CO_2H , SH , PO_3H_2 , SO_3H ,

CONHNH₂, CONHNHSO₂CF₃, OH,

5



10


15

20

wherein each of R⁴² and R⁴³ is independently selected from chloro, cyano, nitro, trifluoromethyl, methoxycarbonyl and trifluoromethylsulfonyl; wherein R² is selected from methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, n-pentyl, isopentyl, neopentyl, phenyl, benzyl, phenethyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cyclopropylmethyl, cyclopropylethyl, cyclobutylmethyl, cyclobutylethyl, cyclopentylmethyl, cyclopentylethyl, cyclohexylmethyl, cyclohexylethyl, methylbutyl, ethylbutyl, dimethylbutyl, thienylmethyl, thienylethyl, thienylpropyl, cyclopentenylmethyl, cyclopentenylethyl, cyclopentenylpropyl, cyclohexenyl, cyclohexenylmethyl, cyclohexenylethyl, adamantyl, adamantylmethyl, adamantylethyl, phenyl, chlorophenyl, dichlorophenyl, fluorophenyl, difluorophenyl, methoxyphenyl, ethoxyphenyl, methylphenyl, ethylphenyl, propylphenyl, isopropylphenyl, dimethylphenyl, diethylphenyl, benzyl, propylthio, butylthio and hydroxyalkyl; wherein each of R³ through R¹¹ is hydrido with the proviso that at least one of R⁵, R⁶, R⁸ and R⁹ is an acidic group selected from CO₂H, SH, PO₃H₂, SO₃H, CONHNH₂, CONHNHSO₂CF₃, OH,

25

30



35

wherein each of R⁴² and R⁴³ is independently selected from chloro, cyano, nitro, trifluoromethyl, methoxycarbonyl and trifluoromethylsulfonyl;

or a tautomer thereof or a pharmaceutically-acceptable salt thereof.

8. Compound of Claim 7 wherein m is one;

40


45

50

55


wherein R¹ is selected from methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, 4-methylbutyl, n-pentyl, isopentyl, neopentyl, phenyl, benzyl, phenethyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cyclopropylmethyl, cyclopropylethyl, cyclobutylmethyl, cyclobutylethyl, cyclopentylmethyl, cyclopentylethyl, cyclohexylmethyl, cyclohexylethyl, cyclohexanoyl, methylbutyl, ethylbutyl, dimethylbutyl, thienylmethyl, thienylethyl, thienylpropyl, cyclopentenylmethyl, cyclopentenylethyl, cyclopentenylpropyl, cyclohexenyl, cyclohexenylmethyl, cyclohexenylethyl, adamantyl, adamantylmethyl, adamantylethyl, phenyl, chlorophenyl, dichlorophenyl, fluorophenyl, difluorophenyl, methoxyphenyl, ethoxyphenyl, methylphenyl, ethylphenyl, propylphenyl, isopropylphenyl, dimethylphenyl, diethylphenyl, benzyl, 1-oxo-2-cyclohexylethyl, benzoyl, 1-oxo-2-phenethyl, 1-oxoethyl, 1-oxopropyl, 1-oxobutyl, 1-oxopentyl, 2-butenyl, 3-butenyl, 2-butyryl, 3-butyryl and 2-hydroxybutyl; wherein R⁰ is selected from hydrido, methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, 4-methylbutyl, tert-butyl, n-pentyl, neopentyl, 1-oxoethyl, 1-oxopropyl, 1-oxobutyl, 1-oxopentyl, 1-oxo-2-phenylethyl, 1-oxo-2-cyclohexylethyl, 1,1-difluoro-2-phenylethyl, 1,1-difluoro-2-cyclohexylethyl, 2-cyclohexylethyl, 1,1-difluoro-3-cyclohexylpropyl, fluoro, chloro, monofluoromethyl, difluoromethyl, trifluoromethyl, formyl, carboxyl, dimethoxymethyl, 1,1-dimethoxybutyl, 1,1-difluoroethyl, 1,1-difluoropropyl, 1,1-difluorobutyl, 1,1-difluoropentyl, benzyl, 2-phenylethyl, 1,1-difluoro-3-phenylpropyl, cyclohexylmethyl, cyclohexylethyl, cyclohexanoyl, 1-butenyl, 2-butenyl, 3-butenyl, 1-butyryl, 2-butyryl, 3-butyryl, propylthio, butylthio, CO₂H, SH, PO₃H₂, SO₃H, CONHNH₂, CONHNHSO₂CF₃, OH,

5



10 wherein each of R⁴² and R⁴³ is independently selected from chloro, cyano, nitro, trifluoromethyl, methoxycarbonyl and trifluoromethylsulfonyl; wherein R² is selected from methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, n-pentyl, isopentyl, neopentyl, phenyl, benzyl, phenethyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cyclopropylmethyl, cyclopropylethyl, cyclobutylmethyl, cyclobutylethyl, cyclopentylmethyl, cyclopentylethyl, cyclohexylmethyl, cyclohexylethyl, methylbutyl, ethylbutyl, dimethylbutyl, thienylmethyl, thienylethyl, thienylpropyl, cyclopentenylmethyl, cyclopentenylethyl, cyclopentenylpropyl, cyclohexenyl, cyclohexenylmethyl, cyclohexenylethyl, adamantyl, adamantylmethyl, adamantylethyl, phenyl, chlorophenyl, dichlorophenyl, fluorophenyl, difluorophenyl, methoxyphenyl, ethoxyphenyl, methylphenyl, ethylphenyl, propylphenyl, isopropylphenyl, dimethylphenyl, diethylphenyl, benzyl, propylthio, butylthio and hydroxyalkyl; wherein each of R³ through R¹¹ is hydrido with the proviso that at least one of R⁵, R⁶, R⁸ and R⁹ is an acidic group selected from CO₂H, SH, PO₃H₂, SO₃H, CONHNH₂, CONHNHSO₂CF₃, OH,

25



30

wherein each of R⁴² and R⁴³ is independently selected from chloro, cyano, nitro, trifluoromethyl, methoxycarbonyl and trifluoromethylsulfonyl;


35

or a tautomer thereof or a pharmaceutically-acceptable salt thereof.

9. Compound of Claim 8 wherein m is one;

wherein R¹ is selected from methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tertbutyl, 4-methylbutyl, n-pentyl, isopentyl, neopentyl, phenyl, benzyl, phenethyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cyclopropylmethyl, cyclopropylethyl, cyclobutylmethyl, cyclobutylethyl, cyclopentylmethyl, cyclopentylethyl, cyclohexylmethyl, cyclohexylethyl, cyclohexanoyl, methylbutyl, ethylbutyl, dimethylbutyl, thienylmethyl, thienylethyl, thienylpropyl, cyclopentenylmethyl, cyclopentenylethyl, cyclopentenylpropyl, cyclohexenyl, cyclohexenylmethyl, cyclohexenylethyl, adamantyl, adamantylmethyl, adamantylethyl, phenyl, chlorophenyl, dichlorophenyl, fluorophenyl, difluorophenyl, methoxyphenyl, ethoxyphenyl, methylphenyl, ethylphenyl, propylphenyl, isopropylphenyl, dimethylphenyl, diethylphenyl, benzyl, 1-oxo-2-cyclohexylethyl, benzoyl, 1-oxo-2-phenethyl, 1-oxopropyl, 1-oxobutyl, 1-oxopentyl and 2-hydroxybutyl; wherein R⁹ is selected from hydrido, methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, 4-methylbutyl, n-pentyl, 1-oxo-2-phenylethyl, 1-oxo-2-cyclohexylethyl, fluoro, chloro, monofluoromethyl, difluoromethyl, trifluoromethyl, formyl, carboxyl, dimethoxymethyl, 1,1-difluoro-2-phenylethyl, 1,1-difluoro-2-cyclohexylethyl, 2-cyclohexylethyl, 1,1-difluoro-3-cyclohexylpropyl, dimethoxymethyl, 1,1-dimethoxybutyl, 1,1-difluoroethyl, 1,1-difluoropropyl, 1,1-difluorobutyl, 1,1-difluoropentyl, benzyl, 2-phenylethyl, 1,1-difluoro-3-phenylpropyl, cyclohexylmethyl, cyclohexylethyl, cyclohexanoyl, 1-butenyl, 2-butenyl, 3-butenyl, 1-butyryl, 2-butyryl, 3-butyryl, propylthio and butylthio; wherein R² is selected from methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, n-pentyl, isopentyl, neopentyl, phenyl, benzyl, phenethyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cyclopropylmethyl, cyclopropylethyl, cyclobutylmethyl, cyclobutylethyl, cyclopentylmethyl, cyclopentylethyl, cyclohexylmethyl, cyclohexylethyl, methylbutyl, ethylbutyl, dimethylbutyl, thienylmethyl, thienylethyl, thienylpropyl, cyclopentenylmethyl, cyclopentenylethyl,

cyclopentenylpropyl, cyclohexenyl, cyclohexenylmethyl, cyclohexenylethyl, adamantyl, adamantylmethyl, adamantylethyl, phenyl, chlorophenyl, dichlorophenyl, fluorophenyl, difluorophenyl, methoxyphenyl, ethoxyphenyl, methylphenyl, ethylphenyl, propylphenyl, isopropylphenyl, dimethylphenyl, diethylphenyl, benzyl, propylthio, butylthio and hydroxyalkyl; wherein each of R^3 , R^4 , R^6 , R^7 , R^8 , R^{10} and R^{11} is hydrido; wherein one of R^5 and R^9 is hydrido and the other of R^5 and R^9 is an acidic group selected from COOH , SH , PO_3H_2 , SO_3H , CONHNH_2 , $\text{CONHNHSO}_2\text{CF}_3$, OH ,




wherein each of R^{42} and R^{43} is independently selected from chloro, cyano, nitro, trifluoromethyl, methoxycarbonyl and trifluoromethylsulfonyl;

or a tautomer thereof or a pharmaceutically-acceptable salt thereof.

10. Compound of Claim 9 wherein m is one;

wherein R^1 is selected from methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, 4-methylbutyl, n-pentyl, isopentyl, neopentyl, phenyl, benzyl, phenethyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cyclopropylmethyl, cyclopropylethyl, cyclobutylmethyl, cyclobutylethyl, cyclopentylmethyl, cyclopentylethyl, cyclohexylmethyl, cyclohexylethyl, cyclohexanoyl, methylbutyl, ethylbutyl, dimethylbutyl, thienylmethyl, thienylethyl, thienylpropyl, cyclopentenylmethyl, cyclopentenylethyl, cyclopentenylpropyl, cyclohexenyl, cyclohexenylmethyl, cyclohexenylethyl, adamantyl, adamantylmethyl, adamantylethyl, phenyl, chlorophenyl, dichlorophenyl, fluorophenyl, difluorophenyl, methoxyphenyl, ethoxyphenyl, methylphenyl, ethylphenyl, propylphenyl, isopropylphenyl, dimethylphenyl, diethylphenyl, benzyl, 1-oxo-2-cyclohexylethyl, benzoyl, 1-oxo-2-phenethyl, 1-oxopropyl, 1-oxobutyl, 1-oxopentyl and 2-hydroxybutyl; wherein R^0 is selected from hydrido, methyl, fluoro, chloro, monofluoromethyl, difluoromethyl, trifluoromethyl, formyl, carboxyl and dimethoxymethyl; wherein R^2 is selected from methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, n-pentyl, isopentyl, neopentyl, phenyl, benzyl, phenethyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cyclopropylmethyl, cyclopropylethyl, cyclobutylmethyl, cyclobutylethyl, cyclopentylmethyl, cyclopentylethyl, cyclohexylmethyl, cyclohexylethyl, methylbutyl, ethylbutyl, dimethylbutyl, thienylmethyl, thienylethyl, thienylpropyl, cyclopentenylmethyl, cyclopentenylethyl, cyclopentenylpropyl, cyclohexenyl, cyclohexenylmethyl, cyclohexenylethyl, adamantyl, adamantylmethyl, adamantylethyl, phenyl, chlorophenyl, dichlorophenyl, fluorophenyl, difluorophenyl, methoxyphenyl, ethoxyphenyl, methylphenyl, ethylphenyl, propylphenyl, isopropylphenyl, dimethylphenyl, diethylphenyl, benzyl, propylthio, butylthio and hydroxyalkyl; wherein each of R^3 , R^4 , R^6 , R^7 , R^8 , R^{10} and R^{11} is hydrido; with the proviso that at least one of R^5 and R^9 must be selected from COOH , SH , PO_3H_2 , SO_3H , CONHNH_2 , $\text{CONHNHSO}_2\text{CF}_3$, OH ,




wherein each of R^{42} and R^{43} is independently selected from chloro, cyano, nitro, trifluoromethyl, methoxycarbonyl and trifluoromethylsulfonyl;

or a tautomer thereof or a pharmaceutically-acceptable salt thereof.

11. Compound of Claim 10 wherein m is one;


wherein R^1 is selected from methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, 4-methylbutyl, n-pentyl, isopentyl, neopentyl, phenyl, benzyl, phenethyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cyclopropylmethyl, cyclopropylethyl, cyclobutylmethyl, cyclobutylethyl,

cyclopentylmethyl, cyclopentylethyl, cyclohexylmethyl, cyclohexylethyl, cyclohexanoyl, methylbutyl, ethylbutyl, dimethylbutyl, thienylmethyl, thienylethyl, thienylpropyl, cyclopentenylmethyl, cyclopentenylethyl, cyclopentenylpropyl, cyclohexenyl, cyclohexenylmethyl, cyclohexenylethyl, adamantyl, adamantylmethyl, adamantylethyl, phenyl, chlorophenyl, dichlorophenyl, fluorophenyl, difluorophenyl, methoxyphenyl, ethoxyphenyl, methylphenyl, ethylphenyl, propylphenyl, isopropylphenyl, dimethylphenyl, diethylphenyl, benzyl, 1-oxo-2-cyclohexylethyl, benzoyl, 1-oxo-2-phenethyl, 1-oxopropyl, 1-oxobutyl, 1-oxopentyl and 2-hydroxybutyl; wherein R^0 is selected from hydrido, methyl, fluoro, chloro, monofluoromethyl, difluoromethyl, trifluoromethyl, formyl, carboxyl and dimethoxymethyl; wherein R^2 is selected from methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, n-pentyl, isopentyl, neopentyl, phenyl, benzyl, phenethyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cyclopropylmethyl, cyclopropylethyl, cyclobutylmethyl, cyclobutylethyl, cyclopentylmethyl, cyclopentylethyl, cyclohexylmethyl, cyclohexylethyl, methylbutyl, ethylbutyl, dimethylbutyl, thienylmethyl, thienylethyl, thienylpropyl, cyclopentenylmethyl, cyclopentenylethyl, cyclopentenylpropyl, cyclohexenyl, cyclohexenylmethyl, cyclohexenylethyl, adamantyl, adamantylmethyl, adamantylethyl, phenyl, chlorophenyl, dichlorophenyl, fluorophenyl, difluorophenyl, methoxyphenyl, ethoxyphenyl, methylphenyl, ethylphenyl, propylphenyl, isopropylphenyl, dimethylphenyl, diethylphenyl, benzyl, propylthio, butylthio and hydroxyalkyl; wherein each of R^3 , R^4 , R^5 , R^7 , R^8 , R^{10} and R^{11} is hydrido; wherein one of R^5 and R^9 is hydrido and the other of R^5 and R^9 is an acidic group selected from CO_2H and




or a tautomer thereof or a pharmaceutically-acceptable salt thereof.

12. Compound of Claim 11 selected from a family of compounds of Formula II




wherein R^1 is selected from methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, 4-methylbutyl, n-pentyl, isopentyl, neopentyl, phenyl, benzyl, phenethyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cyclopropylmethyl, cyclopropylethyl, cyclobutylmethyl, cyclobutylethyl, cyclopentylmethyl, cyclopentylethyl, cyclohexylmethyl, cyclohexylethyl, cyclohexanoyl, methylbutyl, ethylbutyl, dimethylbutyl, thienylmethyl, thienylethyl, thienylpropyl, cyclopentenylmethyl, cyclopentenylethyl, cyclopentenylpropyl, cyclohexenyl, cyclohexenylmethyl, cyclohexenylethyl, adamantyl, adamantylmethyl, adamantylethyl, phenyl, chlorophenyl, dichlorophenyl, fluorophenyl, difluorophenyl, methoxyphenyl, ethoxyphenyl, methylphenyl, ethylphenyl, propylphenyl, isopropylphenyl, dimethylphenyl, diethylphenyl, benzyl, 1-oxo-2-cyclohexylethyl, benzoyl, 1-oxo-2-phenethyl, 1-oxopropyl, 1-oxobutyl, 1-oxopentyl and 2-hydroxybutyl; wherein R^0 is selected from hydrido, methyl, fluoro, chloro, monofluoromethyl, difluoromethyl, trifluoromethyl, formyl, carboxyl and dimethoxymethyl; wherein R^2 is selected from methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, n-pentyl, isopentyl, neopentyl, phenyl, benzyl, phenethyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cyclopropylmethyl, cyclopropylethyl, cyclobutylmethyl, cyclobutylethyl, cyclopentylmethyl, cyclopentylethyl, cyclohexylmethyl, cyclohexylethyl, methylbutyl, ethylbutyl, dimethylbutyl, thienylmethyl, thienylethyl, thienylpropyl, cyclopentenylmethyl, cyclopentenylethyl, cyclopentenylpropyl, cyclohexenyl, cyclohexenylmethyl, cyclohexenylethyl, adamantyl, adamantylmethyl, adamantylethyl, phenyl, chlorophenyl, dichlorophenyl, fluorophenyl, difluorophenyl, methoxyphenyl, ethoxyphenyl, methylphenyl, ethylphenyl, propylphenyl, isopropylphenyl, dimethylphenyl, diethylphenyl and benzyl; wherein R^5 is an acidic group

selected from CO₂H and



or a tautomer thereof or a pharmaceutically-acceptable salt thereof.

13. Compound of Claim 12 wherein R¹ is selected from methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, 4-methylbutyl, n-pentyl, isopentyl, neopentyl, phenyl, benzyl, phenethyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cyclopropylmethyl, cyclopropylethyl, cyclobutylmethyl, cyclobutylethyl, cyclopentylmethyl, cyclopentylethyl, cyclohexylmethyl, cyclohexylethyl, cyclohexanoyl, methylbutyl, ethylbutyl, dimethylbutyl, thienylmethyl, thienylethyl, thienylpropyl, cyclopentenylmethyl, cyclopentenylethyl, cyclopentenylpropyl, cyclohexenyl, cyclohexenylmethyl, cyclohexenylethyl, adamantyl, adamantylmethyl, adamantylethyl, phenyl, chlorophenyl, dichlorophenyl, fluorophenyl, difluorophenyl, methoxyphenyl, ethoxyphenyl, methylphenyl, ethylphenyl, propylphenyl, isopropylphenyl, dimethylphenyl, diethylphenyl, benzyl, 1-oxo-2-cyclohexylethyl, benzoyl, 1-oxo-2-phenethyl, 1-oxopropyl, 1-oxobutyl, 1-oxopentyl and 2-hydroxybutyl; wherein R⁰ is hydrido; wherein R² is selected from methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, n-pentyl, isopentyl, neopentyl, phenyl, benzyl, phenethyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cyclopropylmethyl, cyclopropylethyl, cyclobutylmethyl, cyclobutylethyl, cyclopentylmethyl, cyclopentylethyl, cyclohexylmethyl, cyclohexylethyl, methylbutyl, ethylbutyl, dimethylbutyl, thienylmethyl, thienylethyl, thienylpropyl, cyclopentenylmethyl, cyclopentenylethyl, cyclopentenylpropyl, cyclohexenyl, cyclohexenylmethyl, cyclohexenylethyl, adamantyl, adamantylmethyl, adamantylethyl, phenyl, chlorophenyl, dichlorophenyl, fluorophenyl, difluorophenyl, methoxyphenyl, ethoxyphenyl, methylphenyl, ethylphenyl, propylphenyl, isopropylphenyl, dimethylphenyl, diethylphenyl and benzyl; wherein R⁵ is an acidic group selected from CO₂H and




or a tautomer thereof or a pharmaceutically-acceptable salt thereof.

14. Compound of Claim 13 selected from compounds and their pharmaceutically-acceptable salts of the group of compounds consisting of
- 1-propyl-4-butyl-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
 - 1,4-diethyl-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
 - 1-pentyl-4-butyl-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
 - 1-isopentyl-4-butyl-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
 - 1-cyclohexylmethyl-4-butyl-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
 - 1-phenylmethyl-4-butyl-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
 - 1-(2-cyclohexylethyl)-4-butyl-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
 - 1-(2-phenylethyl)-4-butyl-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
 - 1-propyl-4-butyl-5-methyl-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;


- 1,4-dibutyl-5-methyl-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
 1-pentyl-4-butyl-5-methyl-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
 1-isopentyl-4-butyl-5-methyl-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
 1-cyclohexylmethyl-4-butyl-5-methyl-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
 1-phenylmethyl-4-butyl-5-methyl-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
 1-(2-cyclohexylethyl)-4-butyl-5-methyl-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
 1-(2-phenylethyl)-4-butyl-5-methyl-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
 1-propyl-4-butyl-5-chloro-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
 1,4-dibutyl-5-chloro-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
 1-pentyl-4-butyl-5-chloro-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
 1-isopentyl-4-butyl-5-chloro-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
 1-cyclohexylmethyl-4-butyl-5-chloro-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
 1-phenylmethyl-4-butyl-5-chloro-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
 1-(2-cyclohexylethyl)-4-butyl-5-chloro-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one; and
 1-(2-phenylethyl)-4-butyl-5-chloro-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one.

15. Compound of Claim 11 selected from a family of compounds of Formula III




- wherein R¹ is selected from methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, 4-methylbutyl, n-pentyl, isopentyl, neopentyl, phenyl, benzyl, phenethyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cyclopropylmethyl, cyclopropylethyl, cyclobutylmethyl, cyclobutylethyl, cyclopentylmethyl, cyclopentylethyl, cyclohexylmethyl, cyclohexylethyl, cyclohexanoyl, methylbutyl, ethylbutyl, dimethylbutyl, thienylmethyl, thienylethyl, thienylpropyl, cyclopentenylmethyl, cyclopentenylethyl, cyclopentenylpropyl, cyclohexenyl, cyclohexenylmethyl, cyclohexenylethyl, adamantyl, adamantylmethyl, adamantylethyl, phenyl, chlorophenyl, dichlorophenyl, fluorophenyl, difluorophenyl, methoxyphenyl, ethoxyphenyl, methylphenyl, ethylphenyl, propylphenyl, isopropylphenyl, dimethylphenyl, diethylphenyl, benzyl, 1-oxo-2-cyclohexylethyl, benzoyl, 1-oxo-2-phenethyl, 1-oxopropyl, 1-oxobutyl, 1-oxopentyl and 2-hydroxybutyl; wherein R⁰ is selected from hydrido, methyl, fluoro, chloro, monofluoromethyl, difluoromethyl, trifluoromethyl, formyl, carboxyl and dimethoxymethyl; wherein R² is selected from methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, n-pentyl, isopentyl, neopentyl, phenyl, benzyl, phenethyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cyclopropylmethyl, cyclopropylethyl, cyclobutylmethyl, cyclobutylethyl, cyclopentylmethyl, cyclopentylethyl, cyclohexylmethyl, cyclohexylethyl, methylbutyl, ethylbutyl, dimethylbutyl, thienylmethyl, thienylethyl, thienylpropyl, cyclopentenylmethyl, cyclopentenylethyl, cyclopentenylpropyl, cyclohexenyl, cyclohexenylmethyl, cyclohexenylethyl, adamantyl, adamantylmethyl, adamantylethyl, phenyl, chlorophenyl, dichlorophenyl, fluorophenyl, difluorophenyl, methoxyphenyl, ethoxyphenyl, methylphenyl, ethylphenyl, propylphenyl, isopropylphenyl, dimethylphenyl, diethylphenyl and benzyl; wherein R⁵ is an acidic group

selected from CO₂H and



or a tautomer thereof or a pharmaceutically-acceptable salt thereof.

16. Compound of Claim 15 wherein R¹ is selected from methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, 4-methylbutyl, n-pentyl, isopentyl, neopentyl, phenyl, benzyl, phenethyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cyclopropylmethyl, cyclopropylethyl, cyclobutylmethyl, cyclobutylethyl, cyclopentylmethyl, cyclopentylethyl, cyclohexylmethyl, cyclohexylethyl, cyclohexanoyl, methylbutyl, ethylbutyl, dimethylbutyl, thienylmethyl, thienylethyl, thienylpropyl, cyclopentenylmethyl, cyclopentenylethyl, cyclopentenylpropyl, cyclohexenyl, cyclohexenylmethyl, cyclohexenylethyl, adamantyl, adamantylmethyl, adamantylethyl, phenyl, chlorophenyl, dichlorophenyl, fluorophenyl, difluorophenyl, methoxyphenyl, ethoxyphenyl, methylphenyl, ethylphenyl, propylphenyl, isopropylphenyl, dimethylphenyl, diethylphenyl, benzyl, 1-oxo-2-cyclohexylethyl, benzoyl, 1-oxo-2-phenethyl, 1-oxopropyl, 1-oxobutyl, 1-oxopentyl and 2-hydroxybutyl; wherein R⁰ is hydrido; wherein R² is selected from methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, n-pentyl, isopentyl, neopentyl, phenyl, benzyl, phenethyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cyclopropylmethyl, cyclopropylethyl, cyclobutylmethyl, cyclobutylethyl, cyclopentylmethyl, cyclopentylethyl, cyclohexylmethyl, cyclohexylethyl, methylbutyl, ethylbutyl, dimethylbutyl, thienylmethyl, thienylethyl, thienylpropyl, cyclopentenylmethyl, cyclopentenylethyl, cyclopentenylpropyl, cyclohexenyl, cyclohexenylmethyl, cyclohexenylethyl, adamantyl, adamantylmethyl, adamantylethyl, phenyl, chlorophenyl, dichlorophenyl, fluorophenyl, difluorophenyl, methoxyphenyl, ethoxyphenyl, methylphenyl, ethylphenyl, propylphenyl, isopropylphenyl, dimethylphenyl, diethylphenyl and benzyl; wherein R⁵ is an acidic group selected from CO₂H and
- 15
- 20
- 25
- 30



or a tautomer thereof or a pharmaceutically-acceptable salt thereof.

17. Compound of Claim 16 selected from compounds and their pharmaceutically-acceptable salts of the group of compounds consisting of

1,4-dibutyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;

1,4-dipropyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;

1-propyl-4-isobutyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;

1-butyl-4-isobutyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;

1-butyl-4-isopentyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;

1-butyl-4-pentyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;

1-butyl-4-cyclohexyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;

1-butyl-4-cyclopentyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;

1-butyl-4-(2-cyclopropylethyl)-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;

1-methyl-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;

1-ethyl-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;

1-propyl-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;

1-isobutyl-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;

1-isopentyl-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;

1-pentyl-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;

1-isopropyl-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;

1-tertbutyl-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;

1-neopentyl-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;

1-(3,3-dimethylbutyl)-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;

1-(2-ethylbutyl)-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;

1-cyclohexyl-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;

1-(2-cyclopropylethyl)-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;

1-cyclopentyl-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;

1-cyclopentylmethyl-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;

1-[2-(2-thienyl)ethyl]-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;

1-(2-phenylethyl)-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;

1-(2-cyclopentylethyl)-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;

1-[2-(cyclopenten-1-yl)ethyl]-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;

1-(2-adamantyl)-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;

1-(1-adamantyl)-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;

1-phenyl-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;

1-(2-chlorophenyl)-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;

1-(2-methylphenyl)-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;

1-(2-methoxyphenyl)-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;


1-(2-isopropylphenyl)-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;

1-(2,6-difluorophenyl)-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;


1-(2,6-dichlorophenyl)-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one; and

1-(2,6-dimethylphenyl)-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one.

18. Compound of Claim 11 selected from a family of compounds of Formula IV




wherein R¹ is selected from methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, 4-methylbutyl, n-pentyl, isopentyl, neopentyl, phenyl, benzyl, phenethyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cyclopropylmethyl, cyclopropylethyl, cyclobutylmethyl, cyclobutylethyl, cyclopentylmethyl, cyclopentylethyl, cyclohexylmethyl, cyclohexylethyl, cyclohexanoyl, methylbutyl, ethylbutyl, dimethylbutyl, thienylmethyl, thienylethyl, thienylpropyl, cyclopentenylmethyl, cyclopentenylethyl, cyclopentenylpropyl, cyclohexenyl, cyclohexenylmethyl, cyclohexenylethyl, adamantyl, adamantylmethyl, adamantylethyl, phenyl, chlorophenyl, dichlorophenyl, fluorophenyl, difluorophenyl, methoxyphenyl, ethoxyphenyl, methylphenyl, ethylphenyl, propylphenyl, isopropylphenyl, dimethylphenyl, diethylphenyl and benzyl; wherein R⁰ is selected from hydrido, methyl, fluoro, chloro, monofluoromethyl, difluoromethyl, trifluoromethyl, formyl, carboxyl and dimethoxymethyl; wherein R² is selected from methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, n-pentyl, isopentyl, neopentyl, phenyl, benzyl, phenethyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cyclopropylmethyl, cyclopropylethyl, cyclobutylmethyl, cyclobutylethyl, cyclopentylmethyl, cyclopentylethyl, cyclohexylmethyl, cyclohexylethyl, methylbutyl, ethylbutyl, dimethylbutyl, thienylmethyl, thienylethyl, thienylpropyl, cyclopentenylmethyl, cyclopentenylethyl, cyclopentenylpropyl, cyclohexenyl, cyclohexenylmethyl, cyclohexenylethyl, adamantyl, adamantylmethyl, adamantylethyl, phenyl, chlorophenyl, dichlorophenyl, fluorophenyl, difluorophenyl, methoxyphenyl, ethoxyphenyl, methylphenyl, ethylphenyl, propylphenyl, isopropylphenyl, dimethylphenyl, diethylphenyl and benzyl; wherein R⁵ is an acidic group selected from CO₂H and



or a tautomer thereof or a pharmaceutically-acceptable salt thereof.

19. Compound of Claim 18 wherein R¹ is selected from methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, 4-methylbutyl, n-pentyl, isopentyl, neopentyl, phenyl, benzyl, phenethyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cyclopropylmethyl, cyclopropylethyl, cyclobutylmethyl, cyclobutylethyl, cyclopentylmethyl, cyclopentylethyl, cyclohexylmethyl, cyclohexylethyl, cyclohexanoyl, methylbutyl, ethylbutyl, dimethylbutyl, thienylmethyl, thienylethyl, thienylpropyl, cyclopentenylmethyl, cyclopentenylethyl, cyclopentenylpropyl, cyclohexenyl, cyclohexenylmethyl, cyclohexenylethyl, adamantyl, adamantylmethyl, adamantylethyl, phenyl, chlorophenyl, dichlorophenyl, fluorophenyl, difluorophenyl, methoxyphenyl, ethoxyphenyl, methylphenyl, ethylphenyl, propylphenyl, isopropylphenyl, dimethylphenyl, diethylphenyl, benzyl, 1-oxo-2-cyclohexylethyl, benzoyl, 1-oxo-2-phenethyl, 1-oxopropyl, 1-oxobutyl, 1-oxopentyl and 2-hydroxybutyl; wherein R⁰ is hydrido; wherein R² is selected from methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, n-pentyl, isopentyl, neopentyl, phenyl, benzyl, phenethyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cyclopropylmethyl, cyclopropylethyl, cyclobutylmethyl, cyclobutylethyl, cyclopentylmethyl, cyclopentylethyl, cyclohexylmethyl, cyclohexylethyl, methylbutyl, ethylbutyl, dimethylbutyl, thienylmethyl, thienylethyl, thienylpropyl, cyclopentenylmethyl, cyclopentenylethyl, cyclopentenylpropyl, cyclohexenyl, cyclohexenylmethyl, cyclohexenylethyl, adamantyl, adamantylmethyl, adamantylethyl, phenyl, chlorophenyl, dichlorophenyl, fluorophenyl, difluorophenyl, methoxyphenyl, ethoxyphenyl, methylphenyl, ethylphenyl, propylphenyl, isopropylphenyl, dimethylphenyl, diethylphenyl and benzyl; wherein R⁵ is an acidic group selected from CO₂H and



or a tautomer thereof or a pharmaceutically-acceptable salt thereof.

20. Compound of Claim 19 selected from compounds and their pharmaceutically-acceptable salts of the group of compounds consisting of
- 1-propyl-4-butyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 - 1,4-dibutyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 - 1-pentyl-4-butyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 - 1-isopentyl-4-butyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 - 1-cyclohexylmethyl-4-butyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 - 1-phenylmethyl-4-butyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 - 1-(2-cyclohexylethyl)-4-butyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 - 1-(2-phenylethyl)-4-butyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 - 1-propyl-4-butyl-5-methyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 - 1,4-dibutyl-5-methyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 - 1-pentyl-4-butyl-5-methyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

2-one;

1-isopentyl-4-butyl-5-methyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

1-cyclohexylmethyl-4-butyl-5-methyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

1-phenylmethyl-4-butyl-5-methyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

1-(2-cyclohexylethyl)-4-butyl-5-methyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

1-(2-phenylethyl)-4-butyl-5-methyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

1-propyl-4-butyl-5-chloro-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

1,4-dibutyl-5-chloro-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

1-pentyl-4-butyl-5-chloro-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

1-isopentyl-4-butyl-5-chloro-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;


1-cyclohexylmethyl-4-butyl-5-chloro-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

1-phenylmethyl-4-butyl-5-chloro-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;


1-(2-cyclohexylethyl)-4-butyl-5-chloro-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one; and

1-(2-phenylethyl)-4-butyl-5-chloro-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one.

21. Compound of Claim 11 selected from a family of compounds of Formula V




wherein R¹ is selected from methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, 4-methylbutyl, n-pentyl, isopentyl, neopentyl, phenyl, benzyl, phenethyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cyclopropylmethyl, cyclopropylethyl, cyclobutylmethyl, cyclobutylethyl, cyclopentylmethyl, cyclopentylethyl, cyclohexylmethyl, cyclohexylethyl, cyclohexanoyl, methylbutyl, ethylbutyl, dimethylbutyl, thienylmethyl, thienylethyl, thienylpropyl, cyclopentenylmethyl, cyclopentenylethyl, cyclopentenylpropyl, cyclohexenyl, cyclohexenylmethyl, cyclohexenylethyl, adamantyl, adamantylmethyl, adamantylethyl, phenyl, chlorophenyl, dichlorophenyl, fluorophenyl, difluorophenyl, methoxyphenyl, ethoxyphenyl, methylphenyl, ethylphenyl, propylphenyl, isopropylphenyl, dimethylphenyl, diethylphenyl, benzyl, 1-oxo-2-cyclohexylethyl, benzoyl, 1-oxo-2-phenethyl, 1-oxopropyl, 1-oxobutyl, 1-oxopentyl and 2-hydroxybutyl; wherein R⁰ is selected from hydrido, methyl, fluoro, chloro, monofluoromethyl, difluoromethyl, trifluoromethyl, formyl, carboxyl and dimethoxymethyl; wherein R² is selected from methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, n-pentyl, isopentyl, neopentyl, phenyl, benzyl, phenethyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cyclopropylmethyl, cyclopropylethyl, cyclobutylmethyl, cyclobutylethyl, cyclopentylmethyl, cyclopentylethyl, cyclohexylmethyl, cyclohexylethyl, methylbutyl, ethylbutyl, dimethylbutyl, thienylmethyl, thienylethyl, thienylpropyl, cyclopentenylmethyl, cyclopentenylethyl, cyclopentenylpropyl, cyclohexenyl, cyclohexenylmethyl, cyclohexenylethyl, adamantyl, adamantylmethyl, adamantylethyl, phenyl, chlorophenyl, dichlorophenyl, fluorophenyl, difluorophenyl, methoxyphenyl, ethoxyphenyl, methylphenyl, ethylphenyl, propylphenyl, isopropylphenyl, dimethylphenyl, diethylphenyl and benzyl; wherein R⁵ is an acidic group selected from CO₂H and



or a tautomer thereof or a pharmaceutically-acceptable salt thereof.

22. Compound of Claim 21 wherein R¹ is selected from methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, 4-methylbutyl, n-pentyl, isopentyl, neopentyl, phenyl, benzyl, phenethyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cyclopropylmethyl, cyclopropylethyl, cyclobutylmethyl, cyclobutylethyl, cyclopentylmethyl, cyclopentylethyl, cyclohexylmethyl, cyclohexylethyl, cyclohexanoyl, methylbutyl, ethylbutyl, dimethylbutyl, thienylethyl, thienylpropyl, cyclopentenylmethyl, cyclopentenylethyl, cyclopentenylpropyl, cyclohexenyl, cyclohexenylmethyl, cyclohexenylethyl, adamantyl, adamantylmethyl, adamantylethyl, phenyl, chlorophenyl, dichlorophenyl, fluorophenyl, difluorophenyl, methoxyphenyl, ethoxyphenyl, methylphenyl, ethylphenyl, propylphenyl, isopropylphenyl, dimethylphenyl, diethylphenyl, benzyl, 1-oxo-2-cyclohexylethyl, benzoyl, 1-oxo-2-phenethyl, 1-oxopropyl, 1-oxobutyl, 1-oxopentyl and 2-hydroxybutyl; wherein R⁰ is hydrido; wherein R² is selected from methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, n-pentyl, isopentyl, neopentyl, phenyl, benzyl, phenethyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cyclopropylmethyl, cyclopropylethyl, cyclobutylmethyl, cyclobutylethyl, cyclopentylmethyl, cyclopentylethyl, cyclohexylmethyl, cyclohexylethyl, methylbutyl, ethylbutyl, dimethylbutyl, thienylmethyl, thienylethyl, thienylpropyl, cyclopentenylmethyl, cyclopentenylethyl, cyclopentenylpropyl, cyclohexenyl, cyclohexenylmethyl, cyclohexenylethyl, adamantyl, adamantylmethyl, adamantylethyl, phenyl, chlorophenyl, dichlorophenyl, fluorophenyl, difluorophenyl, methoxyphenyl, ethoxyphenyl, methylphenyl, ethylphenyl, propylphenyl, isopropylphenyl, dimethylphenyl, diethylphenyl and benzyl; wherein R⁵ is an acidic group selected from CO₂H and



or a tautomer thereof or a pharmaceutically-acceptable salt thereof.

23. Compound of Claim 22 selected from compounds and their pharmaceutically-acceptable salts of the group of compounds consisting of
- 1-propyl-4-butyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
- 1,4-dibutyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
- 1-pentyl-4-butyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
- 1-isopentyl-4-butyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
- 1-cyclohexylmethyl-4-butyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
- 1-phenylmethyl-4-butyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
- 1-(2-cyclohexylethyl)-4-butyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
- 1-(2-phenylethyl)-4-butyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
- 1-propyl-4-butyl-5-methyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
- 1,4-dibutyl-5-methyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
- 1-pentyl-4-butyl-5-methyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

2-one;

1-isopentyl-4-butyl-5-methyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

1-cyclohexylmethyl-4-butyl-5-methyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

1-phenylmethyl-4-butyl-5-methyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

1-(2-cyclohexylethyl)-4-butyl-5-methyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

1-(2-phenylethyl)-4-butyl-5-methyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

1-propyl-4-butyl-5-chloro-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

1,4-dibutyl-5-chloro-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

1-pentyl-4-butyl-5-chloro-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

1-isopentyl-4-butyl-5-chloro-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;


1-cyclohexylmethyl-4-butyl-5-chloro-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

1-phenylmethyl-4-butyl-5-chloro-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;


1-(2-cyclohexylethyl)-4-butyl-5-chloro-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one; and

1-(2-phenylethyl)-4-butyl-5-chloro-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one.

24. Compound of Claim 11 selected from a family of compounds of Formula VI




wherein R¹ is selected from methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, 4-methylbutyl, n-pentyl, isopentyl, neopentyl, phenyl, benzyl, phenethyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cyclopropylmethyl, cyclopropylethyl, cyclobutylmethyl, cyclobutylethyl, cyclopentylmethyl, cyclopentylethyl, cyclohexylmethyl, cyclohexylethyl, cyclohexanoyl, methylbutyl, ethylbutyl, dimethylbutyl, thienylmethyl, thienylethyl, thienylpropyl, cyclopentenylmethyl, cyclopentenylethyl, cyclopentenylpropyl, cyclohexenyl, cyclohexenylmethyl, cyclohexenylethyl, adamantyl, adamantylmethyl, adamantylethyl, phenyl, chlorophenyl, dichlorophenyl, fluorophenyl, difluorophenyl, methoxyphenyl, ethoxyphenyl, methylphenyl, ethylphenyl, propylphenyl, isopropylphenyl, dimethylphenyl, diethylphenyl, benzyl, 1-oxo-2-cyclohexylethyl, benzoyl, 1-oxo-2-phenethyl, 1-oxopropyl, 1-oxobutyl, 1-oxopentyl and 2-hydroxybutyl; wherein R⁰ is selected from hydrido, methyl, fluoro, chloro, monofluoromethyl, difluoromethyl, trifluoromethyl, formyl, carboxyl and dimethoxymethyl; wherein R² is selected from methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, n-pentyl, isopentyl, neopentyl, phenyl, benzyl, phenethyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cyclopropylmethyl, cyclopropylethyl, cyclobutylmethyl, cyclobutylethyl, cyclopentylmethyl, cyclopentylethyl, cyclohexylmethyl, cyclohexylethyl, methylbutyl, ethylbutyl, dimethylbutyl, thienylmethyl, thienylethyl, thienylpropyl, cyclopentenylmethyl, cyclopentenylethyl, cyclopentenylpropyl, cyclohexenyl, cyclohexenylmethyl, cyclohexenylethyl, adamantyl, adamantylmethyl, adamantylethyl, phenyl, chlorophenyl, dichlorophenyl, fluorophenyl, difluorophenyl, methoxyphenyl, ethoxyphenyl, methylphenyl, ethylphenyl, propylphenyl, isopropylphenyl, dimethylphenyl, diethylphenyl and benzyl; wherein R⁵ is an acidic group selected from CO₂H and



or a tautomer thereof or a pharmaceutically-acceptable salt thereof.

25. Compound of Claim 24 wherein R¹ is selected from methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, 4-methylbutyl, n-pentyl, isopentyl, neopentyl, phenyl, benzyl, phenethyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cyclopropylmethyl, cyclopropylethyl, cyclobutylmethyl, cyclobutylethyl, cyclopentylmethyl, cyclopentylethyl, cyclohexylmethyl, cyclohexylethyl, cyclohexanoyl, methylbutyl, ethylbutyl, dimethylbutyl, thienylmethyl, thienylethyl, thienylpropyl, cyclopentenylmethyl, cyclopentenylethyl, cyclopentenylpropyl, cyclohexenyl, cyclohexenylmethyl, cyclohexenylethyl, adamantyl, adamantylmethyl, adamantylethyl, phenyl, chlorophenyl, dichlorophenyl, fluorophenyl, difluorophenyl, methoxyphenyl, ethoxyphenyl, methylphenyl, ethylphenyl, propylphenyl, isopropylphenyl, dimethylphenyl, diethylphenyl, benzyl, 1-oxo-2-cyclohexylethyl, benzoyl, 1-oxo-2-phenethyl, 1-oxopropyl, 1-oxobutyl, 1-oxopentyl and 2-hydroxybutyl; wherein R⁰ is hydrido; wherein R² is selected from methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, n-pentyl, isopentyl, neopentyl, phenyl, benzyl, phenethyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cyclopropylmethyl, cyclopropylethyl, cyclobutylmethyl, cyclobutylethyl, cyclopentylmethyl, cyclopentylethyl, cyclohexylmethyl, cyclohexylethyl, methylbutyl, ethylbutyl, dimethylbutyl, thienylmethyl, thienylethyl, thienylpropyl, cyclopentenylmethyl, cyclopentenylethyl, cyclopentenylpropyl, cyclohexenyl, cyclohexenylmethyl, cyclohexenylethyl, adamantyl, adamantylmethyl, adamantylethyl, phenyl, chlorophenyl, dichlorophenyl, fluorophenyl, difluorophenyl, methoxyphenyl, ethoxyphenyl, methylphenyl, ethylphenyl, propylphenyl, isopropylphenyl, dimethylphenyl, diethylphenyl and benzyl; wherein R⁵ is an acidic group selected from CO₂H and



or a tautomer thereof or a pharmaceutically-acceptable salt thereof.

26. Compound of Claim 25 selected from compounds and their pharmaceutically-acceptable salts of the group of compounds consisting of
- 1-propyl-4-butyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 - 1,4-dibutyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 - 1-pentyl-4-butyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 - 1-isopentyl-4-butyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 - 1-cyclohexylmethyl-4-butyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 - 1-phenylmethyl-4-butyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 - 1-(2-cyclohexylethyl)-4-butyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 - 1-(2-phenylethyl)-4-butyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 - 1-propyl-4-butyl-5-methyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 - 1,4-dibutyl-5-methyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 - 1-pentyl-4-butyl-5-methyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

2-one;

1-isopentyl-4-butyl-5-methyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

1-cyclohexylmethyl-4-butyl-5-methyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

1-phenylmethyl-4-butyl-5-methyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

1-(2-cyclohexylethyl)-4-butyl-5-methyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

1-(2-phenylethyl)-4-butyl-5-methyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

1-propyl-4-butyl-5-chloro-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

1,4-dibutyl-5-chloro-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

1-pentyl-4-butyl-5-chloro-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

1-isopentyl-4-butyl-5-chloro-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;


1-cyclohexylmethyl-4-butyl-5-chloro-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

1-phenylmethyl-4-butyl-5-chloro-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

1-(2-cyclohexylethyl)-4-butyl-5-chloro-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one; and


1-(2-phenylethyl)-4-butyl-5-chloro-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one.

27. Compound of Claim 11 selected from a family of compounds of Formula VII




(VII)

wherein R¹ is selected from methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, 4-methylbutyl, n-pentyl, isopentyl, neopentyl, phenyl, benzyl, phenethyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cyclopropylmethyl, cyclopropylethyl, cyclobutylmethyl, cyclobutylethyl, cyclopentylmethyl, cyclopentylethyl, cyclohexylmethyl, cyclohexylethyl, cyclohexanoyl, methylbutyl, ethylbutyl, dimethylbutyl, thienylmethyl, thienylethyl, thienylpropyl, cyclopentenylmethyl, cyclopentenylethyl, cyclopentenylpropyl, cyclohexenyl, cyclohexenylmethyl, cyclohexenylethyl, adamantyl, adamantylmethyl, adamantylethyl, phenyl, chlorophenyl, dichlorophenyl, fluorophenyl, difluorophenyl, methoxyphenyl, ethoxyphenyl, methylphenyl, ethylphenyl, propylphenyl, isopropylphenyl, dimethylphenyl, diethylphenyl, benzyl, 1-oxo-2-cyclohexylethyl, benzoyl, 1-oxo-2-phenethyl, 1-oxopropyl, 1-oxobutyl, 1-oxopentyl and 2-hydroxybutyl; wherein R⁰ is selected from hydrido, methyl, fluoro, chloro, monofluoromethyl, difluoromethyl, trifluoromethyl, formyl, carboxyl and dimethoxymethyl; wherein R² is selected from methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, n-pentyl, isopentyl, neopentyl, phenyl, benzyl, phenethyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cyclopropylmethyl, cyclopropylethyl, cyclobutylmethyl, cyclobutylethyl, cyclopentylmethyl, cyclopentylethyl, cyclohexylmethyl, cyclohexylethyl, methylbutyl, ethylbutyl, dimethylbutyl, thienylmethyl, thienylethyl, thienylpropyl, cyclopentenylmethyl, cyclopentenylethyl, cyclopentenylpropyl, cyclohexenyl, cyclohexenylmethyl, cyclohexenylethyl, adamantyl, adamantylmethyl, adamantylethyl, phenyl, chlorophenyl, dichlorophenyl, fluorophenyl, difluorophenyl, methoxyphenyl, ethoxyphenyl, methylphenyl, ethylphenyl, propylphenyl, isopropylphenyl, dimethylphenyl, diethylphenyl and benzyl; wherein R⁵ is an acidic group selected from CO₂H and



or a tautomer thereof or a pharmaceutically-acceptable salt thereof.

28. Compound of Claim 27 wherein R¹ is selected from methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, 4-methylbutyl, n-pentyl, isopentyl, neopentyl, phenyl, benzyl, phenethyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cyclopropylmethyl, cyclopropylethyl, cyclobutylmethyl, cyclobutylethyl, cyclopentylmethyl, cyclopentylethyl, cyclohexylmethyl, cyclohexylethyl, cyclohexanoyl, methylbutyl, ethylbutyl, dimethylbutyl, thienylmethyl, thienylethyl, thienylpropyl, cyclopentenylmethyl, cyclopentenylethyl, cyclopentenylpropyl, cyclohexenyl, cyclohexenylmethyl, cyclohexenylethyl, adamantyl, adamantylmethyl, adamantylethyl, phenyl, chlorophenyl, dichlorophenyl, fluorophenyl, difluorophenyl, methoxyphenyl, ethoxyphenyl, methylphenyl, ethylphenyl, propylphenyl, isopropylphenyl, dimethylphenyl, diethylphenyl, benzyl, 1-oxo-2-cyclohexylethyl, benzoyl, 1-oxo-2-phenethyl, 1-oxopropyl, 1-oxobutyl, 1-oxopentyl and 2-hydroxybutyl; wherein R⁰ is hydrido; wherein R² is selected from methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, n-pentyl, isopentyl, neopentyl, phenyl, benzyl, phenethyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cyclopropylmethyl, cyclopropylethyl, cyclobutylmethyl, cyclobutylethyl, cyclopentylmethyl, cyclopentylethyl, cyclohexylmethyl, cyclohexylethyl, methylbutyl, ethylbutyl, dimethylbutyl, thienylmethyl, thienylethyl, thienylpropyl, cyclopentenylmethyl, cyclopentenylethyl, cyclopentenylpropyl, cyclohexenyl, cyclohexenylmethyl, cyclohexenylethyl, adamantyl, adamantylmethyl, adamantylethyl, phenyl, chlorophenyl, dichlorophenyl, fluorophenyl, difluorophenyl, methoxyphenyl, ethoxyphenyl, methylphenyl, ethylphenyl, propylphenyl, isopropylphenyl, dimethylphenyl, diethylphenyl and benzyl; wherein R⁵ is an acidic group selected from CO₂H and



or a tautomer thereof or a pharmaceutically-acceptable salt thereof.

29. Compound of Claim 28 selected from compounds and their pharmaceutically-acceptable salts of the group of compounds consisting of
- 1-propyl-4-butyl-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 - 1,4-dibutyl-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 - 1-pentyl-4-butyl-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 - 1-isopentyl-4-butyl-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 - 1-cyclohexylmethyl-4-butyl-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 - 1-phenylmethyl-4-butyl-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 - 1-(2-cyclohexylethyl)-4-butyl-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 - 1-(2-phenylethyl)-4-butyl-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 - 1-propyl-4-butyl-5-methyl-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 - 1,4-dibutyl-5-methyl-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 - 1-pentyl-4-butyl-5-methyl-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-

2-one;

1-isopentyl-4-butyl-5-methyl-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

1-cyclohexylmethyl-4-butyl-5-methyl-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

1-phenylmethyl-4-butyl-5-methyl-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

1-(2-cyclohexylethyl)-4-butyl-5-methyl-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

1-(2-phenylethyl)-4-butyl-5-methyl-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

1-propyl-4-butyl-5-chloro-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

1,4-dibutyl-5-chloro-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

1-pentyl-4-butyl-5-chloro-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

1-isopentyl-4-butyl-5-chloro-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

1-cyclohexylmethyl-4-butyl-5-chloro-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

1-phenylmethyl-4-butyl-5-chloro-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;


1-(2-cyclohexylethyl)-4-butyl-5-chloro-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one; and

1-(2-phenylethyl)-4-butyl-5-chloro-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one.

30. Compound of Claim 17 which is 1,4-dibutyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one or a pharmaceutically-acceptable salt thereof.


31. Compound of Claim 17 which is 1-(2-ethylbutyl)-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one or a pharmaceutically-acceptable salt thereof.

32. A pharmaceutical composition comprising a therapeutically-effective amount of an angiotensin II antagonist compound and a pharmaceutically-acceptable carrier or diluent, said antagonist compound selected from a family of compounds of Formula I




(I)

wherein A is selected from



wherein m is a number selected from one to four, inclusive;

wherein R¹ is selected from alkyl, hydroxyalkyl, haloalkyl, cycloalkyl, cycloalkylalkyl, formyl, thienylalkyl, phenylalkyl, polycycloalkyl, polycycloalkylalkyl, phenyl, halophenyl, alkylphenyl, alkoxyphenyl, cycloalkenyl, cycloalkenylalkyl, aroyl, alkoxyalkyl, alkylcarbonyl, alkylcarbonylalkyl, cycloalkylcarbonyl, cycloalkylalkylcarbonyl, aralkylcarbonyl, alkoxycarbonyl, alkenyl, cycloalkenyl, aralkoxycarbonyl, alkynyl, alkylthiocarbonyl, alkylthiothiocarbonyl, arylthiocarbonyl, arylthiothiocarbonyl, aralkylthiocarbonyl, alkylthiocarbonyl, alkylsulfinyl, alkylsulfonyl, aralkylsulfinyl, aralkylsulfonyl, arylsulfinyl, arylsulfonyl, heteroaryl having one or more ring atoms selected from oxygen, sulfur and nitrogen atoms, and amido radicals of the formula



wherein X is oxygen atom or sulfur atom;

wherein each of R¹² and R¹³ is independently selected from hydrido, alkyl, cycloalkyl, cyano, amino, monoalkylamino, dialkylamino, hydroxyalkyl, cycloalkylalkyl, alkoxyalkyl, aralkyl and aryl, and wherein R¹² and R¹³ taken together may form a heterocyclic group having five to seven ring members including the nitrogen atom of said amido radical and which heterocyclic group may further contain one or more hetero atoms as ring members selected from oxygen, nitrogen and sulfur atoms and which heterocyclic group may be saturated or partially unsaturated; wherein each of R¹² and R¹³ taken together may form an aromatic heterocyclic group having five ring members including the nitrogen atom of said amido radical and which aromatic heterocyclic group may further contain one or more additional nitrogen atoms;


wherein each of R⁰ and R² through R¹¹ is independently selected from hydrido, alkyl, hydroxyalkyl, formyl, halo, haloalkyl, cycloalkyl, cycloalkylalkyl, cycloalkylhaloalkyl, cycloalkylcarbonyl, alkoxy, thienylalkyl, phenylalkyl, polycycloalkyl, polycycloalkylalkyl, phenyl, halophenyl, alkylphenyl, alkoxyphenyl, cycloalkenyl, cycloalkenylalkyl, aralkylhaloalkyl, aroyl, aryloxy, aryloxyalkyl, aralkoxy, alkoxyalkyl, alkylcarbonyl, alkylcarbonylalkyl, alkoxycarbonyl, alkenyl, cycloalkenyl, alkynyl, cyano, nitro, carboxyl, carboxyalkyl, alkylcarbonyloxy, alkylcarbonyloxyalkyl, alkoxycarbonylalkyl, aralkoxycarbonylalkyl, aralkylcarbonyloxyalkyl, mercaptocarbonyl, mercaptothiocarbonyl, mercaptoalkyl, alkoxycarbonyloxy, alkylthio, cycloalkylthio, cycloalkylalkylthio, alkylthiocarbonyl, alkylcarbonylthio, alkylthiocarbonyloxy, alkylthiocarbonylthio, alkylthiothiocarbonyl, alkylthiothiocarbonylthio, arylthio, arylthiocarbonyl, arylcarbonylthio, arylthiocarbonyloxy, arylthiocarbonylthio, arylthiothiocarbonyl, arylthiothiocarbonylthio, aralkylthio, aralkylthiocarbonyl, aralkylcarbonylthio, aralkylthiocarbonyloxy, aralkylthiocarbonylthio, alkylthiocarbonyl, aralkylthiocarbonylthio, mercapto, alkylsulfinyl, alkylsulfonyl, aralkylsulfinyl, aralkylsulfonyl, arylsulfinyl, arylsulfonyl, phthalimido, phthalimidoalkyl, heteroaryl, heteroarylalkyl, cycloheteroalkyl,

cycloheteroalkylalkyl and cycloheteroalkylcarbonylalkyl wherein each of said heteroaryl- and cyclohetero-containing groups has one or more ring atoms selected from oxygen, sulfur and nitrogen atoms, and wherein each of R⁰ and R² through R¹¹ may be further independently selected from amino and amido radicals of the formula

5

10

15



wherein X is oxygen atom or sulfur atom;

20

wherein each n is a number independently selected from zero to six, inclusive;

25

30

wherein each of R¹⁴ through R²⁶ is independently selected from hydrido, alkyl, cycloalkyl, cyano, amino, monoalkylamino, dialkylamino, hydroxyalkyl, cycloalkylalkyl, alkoxyalkyl, aralkyl and aryl, and wherein R¹⁴ and R¹⁵ taken together, R¹⁶ and R¹⁷ taken together, R¹⁸ and R¹⁹ taken together, R²¹ and R²² taken together and R²³ and R²⁴ taken together may each form a heterocyclic group having five to seven ring members including the nitrogen atom of said amino or amido radical and which heterocyclic group may further contain one or more hetero atoms as ring members selected from oxygen, nitrogen and sulfur atoms and which heterocyclic group may be saturated or partially unsaturated; wherein R¹⁴ and R¹⁵ taken together, R¹⁶ and R¹⁷ taken together, R²¹ and R²² taken together and R²³ and R²⁴ taken together may each form an aromatic heterocyclic group having five ring members including the nitrogen atom of said amino or amido radical and which aromatic heterocyclic group may further contain one or more additional nitrogen atoms;

35

and wherein each of R⁰ and R³ through R¹¹ may be further independently selected from hydroxy and acidic moieties of the formula



40

wherein n is a number selected from zero through three, inclusive, and wherein A is an acidic group selected to contain at least one acidic hydrogen atom, and the amide, ester and salt derivatives of said acidic moieties;


45

wherein Y is a spacer group independently selected from one or more of alkyl, cycloalkyl, cycloalkylalkyl, alkenyl, alkynyl, aryl, aralkyl and heteroaryl having one or more ring atoms selected from oxygen, sulfur and nitrogen atoms;


50

and wherein any of the foregoing R⁰ through R²⁶, Y and A groups having a substitutable position may be substituted by one or more groups independently selected from hydroxy, alkyl, alkenyl, alkynyl, aralkyl, hydroxyalkyl, haloalkyl, halo, oxo, alkoxy, aryloxy, aralkoxy, aralkylthio, alkoxyalkyl, cycloalkyl, cycloalkylalkyl, aryl, aroyl, cycloalkenyl, cyano, cyanoamino, nitro, alkylcarbonyloxy, alkoxyalkyl, alkylcarbonyl, alkoxyalkyl, aralkoxyalkyl, carboxyl, mercapto, mercaptoalkyl, alkylthio, arylthio, alkylthiocarbonyl, alkylsulfonyl, alkylsulfonate, haloalkylsulfonate, aralkylsulfonate, aralkylsulfonate, arylsulfonate, arylsulfonate, heteroaryl having one or more ring atoms selected from oxygen, sulfur and nitrogen atoms, and amino and amido radicals of the formula


55



wherein X is oxygen atom or sulfur atom; wherein each of R²⁷ through R³¹ is independently selected from hydrido, alkyl, cycloalkyl, cycloalkylalkyl, aralkyl, aryl, DR³² and




wherein D is selected from oxygen atom and sulfur atom and R³² is selected from hydrido, alkyl, cycloalkyl, cycloalkylalkyl, aralkyl and aryl; wherein each of R²⁷, R²⁸, R²⁹, R³⁰, R³¹, R³³ and R³⁴ is independently selected from hydrido, alkyl, cycloalkyl, cyano, hydroxyalkyl, haloalkyl, cycloalkylalkyl, alkoxyalkyl, alkylcarbonyl, alkoxy carbonyl, carboxyl, alkylsulfinyl, alkylsulfonyl, arylsulfinyl, arylsulfonyl, haloalkylsulfinyl, haloalkylsulfonyl, aralkyl and aryl, and wherein each of R²⁷, R²⁸, R²⁹, R³⁰, R³¹, R³³ and R³⁴ is further independently selected from amino and amido radicals of the formula



wherein X is oxygen atom or sulfur atom;

wherein each of R³⁵, R³⁶, R³⁷, R³⁸, R³⁹ and R⁴⁰ is independently selected from hydrido, alkyl, cycloalkyl, cyano, hydroxyalkyl, cycloalkylalkyl, alkoxyalkyl, haloalkylsulfinyl, haloalkylsulfonyl, aralkyl and aryl, and wherein each of R²⁸ and R²⁹ taken together and each of R³⁰ and R³¹ taken together may form a heterocyclic group having five to seven ring members including the nitrogen atom of said amino or amido radical, which heterocyclic group may further contain one or more hetero atoms as ring members selected from oxygen, nitrogen and sulfur atoms and which heterocyclic group may be saturated or partially unsaturated; wherein each of R²⁸ and R²⁹ taken together and each of R³³ and R³⁴ taken together may form an aromatic heterocyclic group having five ring members including the nitrogen atom of said amino or amido radical and which aromatic heterocyclic group may further contain one or more additional nitrogen atoms; or a tautomer thereof or a pharmaceutically-acceptable salt thereof.


33. The composition of Claim 32 wherein m is one; wherein R¹ is selected from alkyl, hydroxyalkyl, haloalkyl, cycloalkyl, cycloalkylalkyl, thienylalkyl, phenylalkyl, polycycloalkyl, polycycloalkylalkyl, phenyl, halophenyl, alkylphenyl, alkoxyphenyl, cycloalkenyl, cycloalkenylalkyl, aroyl, alkoxyalkyl, alkylcarbonyl, cycloalkylcarbonyl, cycloalkylalkylcarbonyl, aralkylcarbonyl, alkoxy carbonyl, alkenyl, cycloalkenyl, alkynyl, cycloalkynyl, alkylthiocarbonyl, arylthiocarbonyl, arylthiothiocarbonyl, aralkylthiocarbonyl, alkylsulfonyl, aralkylsulfonyl, arylsulfonyl, heteroaryl having one or more ring atoms selected from oxygen, sulfur and nitrogen atoms, and amido radicals of the formula



wherein X is oxygen atom or sulfur atom;

wherein each of R¹² and R¹³ is independently selected from hydrido, alkyl, cycloalkyl, cyano, amino, monoalkylamino, dialkylamino, hydroxyalkyl, cycloalkylalkyl, alkoxyalkyl, aralkyl and aryl;

5 wherein each of R⁰ and R² is independently selected from hydrido, alkyl, hydroxyalkyl, formyl, halo, haloalkyl, cycloalkyl, cycloalkylalkyl, cycloalkylhaloalkyl, cycloalkylcarbonyl, alkoxy, thienylalkyl, phenylalkyl, polycycloalkyl, polycycloalkylalkyl, phenyl, halophenyl, alkylphenyl, alkoxyphenyl, cycloalkenyl, cycloalkenylalkyl, aralkylhaloalkyl, aroyl, aryloxy, aryloxyalkyl, aralkoxy, alkoxyalkyl, alkylcarbonyl, alkoxyalkyl, alkenyl, cycloalkenyl, alkynyl, cycloalkynyl, cyano, nitro, carboxyl, carboxyalkyl, alkylcarbonyloxy, alkylcarbonyloxyalkyl, alkoxyalkyl, aralkoxyalkyl, aralkylcarbonyloxy, mercaptocarbonyl, mercaptothiocarbonyl, mercaptoalkyl, alkoxyalkyl, alkylthio, cycloalkylthio, cycloalkylalkylthio, alkylthiocarbonyl, alkylcarbonylthio, alkylthiocarbonyloxy, alkylthiocarbonylthio, alkylthiothiocarbonyl, alkylthiothiocarbonylthio, arylthio, arylthiocarbonyl, arylcarbonylthio, arylthiocarbonyloxy, arylthiocarbonylthio, arylthiothiocarbonyl, arylthiothiocarbonylthio, aralkylthio, aralkylthiocarbonyl, aralkylcarbonylthio, aralkylthiocarbonyloxy, aralkylthiocarbonylthio, aralkylthiothiocarbonyl, aralkylthiothiocarbonylthio, mercapto, alkylsulfonyl, alkylsulfonyl, aralkylsulfonyl, aralkylsulfonyl, arylsulfonyl, phthalimido, phthalimidoalkyl, heteroaryl, heteroarylalkyl, cycloheteroalkyl, cycloheteroalkylalkyl and cycloheteroalkylcarbonylalkyl wherein each of said heteroaryl- and cycloheteroalkyl-containing groups has one or more hetero ring atoms selected from oxygen, sulfur and nitrogen atoms, and
 10 15 20 wherein each of R⁰ and R² through R¹¹ may be further independently selected from amino and amido radicals of the formula




35 wherein X is selected from oxygen atom or sulfur atom;

wherein each n is a number independently selected from zero to six, inclusive;

40 wherein each of R¹⁴ through R²⁶ is independently selected from hydrido, alkyl, cycloalkyl, cyano, amino, monoalkylamino, dialkylamino, hydroxyalkyl, cycloalkylalkyl, alkoxyalkyl, aralkyl and aryl;

45 wherein each of R³ through R¹¹ is independently selected from hydrido, hydroxy, alkyl, hydroxyalkyl, halo, haloalkyl, cycloalkyl, cycloalkylalkyl, alkoxy, aralkyl, aryl, aroyl, aryloxy, aralkoxy, alkoxyalkyl, alkylcarbonyl, alkoxyalkyl, alkenyl, cycloalkenyl, alkynyl, cycloalkynyl, cyano, nitro, carboxyl, alkylcarbonyloxy, mercaptocarbonyl, mercaptothiocarbonyl, alkoxyalkyl, alkylthio, alkylthiocarbonyl, alkylcarbonylthio, alkylthiocarbonyloxy, alkylthiocarbonylthio, alkylthiothiocarbonyl, arylthio, arylthiocarbonyl, arylcarbonylthio, arylthiocarbonyloxy, arylthiothiocarbonyl, aralkylthio, aralkylthiocarbonyl, aralkylcarbonylthio, aralkylthiocarbonyloxy, aralkylthiocarbonylthio, aralkylthiothiocarbonyl, aralkylthiothiocarbonylthio, mercapto, alkylsulfonyl, aralkylsulfonyl and arylsulfonyl, and amino and amido radicals of the formula



wherein X is oxygen atom or sulfur atom;


wherein each of R¹⁴, R¹⁵, R¹⁶, R¹⁷, R¹⁸ and R¹⁹ is independently selected from hydrido, alkyl, cycloalkyl, cyano, hydroxyalkyl, cycloalkylalkyl, alkoxyalkyl, aralkyl and aryl;

5 and wherein each of R⁰ and R³ through R¹¹ may be further independently selected from acidic moieties of the formula



10 wherein n is a number selected from zero through three, inclusive; wherein A is an acidic group selected from acids containing one or more atoms selected from oxygen, sulfur, phosphorus and nitrogen atoms, and wherein said acidic group is selected to contain at least one acidic hydrogen atom, and the amide, ester and salt derivatives of said acidic moieties; wherein Y is a spacer group
15 independently selected from one or more of alkyl, cycloalkyl, cycloalkylalkyl, alkenyl, alkynyl, aryl, aralkyl and heteroaryl having one or more ring atoms selected from oxygen, sulfur and nitrogen atoms;

and wherein any of the foregoing R⁰ through R²⁶, Y and A groups having a substitutable position may be substituted by one or more groups independently selected from hydroxy, alkyl, alkenyl, aralkyl, hydroxyalkyl, halo, haloalkyl, oxo, alkoxy, aryloxy, aralkoxy, alkoxyalkyl, alkylcarbonyl, alkoxycarbonyl,
20 carboxyl, cyano, nitro, alkylsulfonyl, haloalkylsulfonyl, aryl, aralkyl, mercaptocarbonyl, alkylthio and alkylthiocarbonyl, and amino and amido radicals of the formula



30 wherein X is oxygen atom or sulfur atom; wherein each of R²⁷ through R³¹ is independently selected from hydrido, alkyl, cycloalkyl, cycloalkylalkyl, aralkyl, aryl, and DR³² and



40 wherein D is selected from oxygen atom and sulfur atom, and R³² is selected from hydrido, alkyl, cycloalkyl, cycloalkylalkyl, aralkyl and aryl; wherein each of R²⁷, R²⁸, R²⁹, R³⁰, R³¹, R³³ and R³⁴ is independently selected from hydrido, alkyl, cycloalkyl, cyano, hydroxyalkyl, haloalkyl, cycloalkylalkyl, alkoxyalkyl, alkanoyl, alkoxycarbonyl, carboxyl, haloalkylsulfinyl, haloalkylsulfonyl, aralkyl and aryl;


or a tautomer thereof or a pharmaceutically-acceptable salt thereof.

45 **34.** The composition of Claim 33 wherein m is one; wherein R¹ is selected from alkyl, hydroxyalkyl, haloalkyl, cycloalkyl, cycloalkylalkyl, thienylalkyl, phenylalkyl, polycycloalkyl, polycycloalkylalkyl, phenyl, halophenyl, alkylphenyl, alkoxyphenyl, cycloalkenyl, cycloalkenylalkyl, aroyl, alkoxyalkyl, alkylcarbonyl, cycloalkylcarbonyl, cycloalkylalkylcarbonyl, aralkylcarbonyl, alkoxycarbonyl, alkenyl, cycloalkenyl, alkynyl, cycloalkynyl, alkylsulfonyl, aralkylsulfonyl and arylsulfonyl, and amido radicals of the
50 formula



wherein each of R^{12} and R^{13} is independently selected from hydrido, alkyl, cycloalkyl, cyano, amino, monoalkylamino, dialkylamino, hydroxyalkyl, cycloalkylalkyl, alkoxyalkyl, aralkyl and aryl;

wherein each of R^0 and R^2 is independently selected from hydrido, alkyl, hydroxyalkyl, formyl, halo, haloalkyl, cycloalkyl, cycloalkylalkyl, cycloalkylhaloalkyl, cycloalkylcarbonyl, alkoxy, thienylalkyl, phenylalkyl, polycycloalkyl, polycycloalkylalkyl, phenyl, halophenyl, alkylphenyl, alkoxyphenyl, cycloalkenyl, cycloalkenylalkyl, aralkylhaloalkyl, aroyl, aryloxy, aryloxyalkyl, aralkoxy, alkoxyalkyl, alkylcarbonyl, alkoxyalkyl, alkenyl, cycloalkenyl, alkynyl, cycloalkynyl, cyano, nitro, carboxyl, carboxyalkyl, alkylcarbonyloxy, mercaptocarbonyl, alkoxyalkoxy, alkylcarbonyloxyalkyl, alkoxyalkoxyalkyl, aralkoxyalkoxyalkyl, aralkylcarbonyloxyalkyl, alkylthio, cycloalkylthio, cycloalkylalkylthio, arylthio, aralkylthio, aralkylthiocarbonylthio, mercapto, alkylsulfinyl, alkylsulfonyl, aralkylsulfinyl, aralkylsulfonyl, arylsulfinyl, arylsulfonyl, phthalimido, phthalimidoalkyl, heteroaryl, heteroarylalkyl, cycloheteroalkyl, cycloheteroalkylalkyl and cycloheteroalkylcarbonylalkyl wherein each of said heteroaryl- and cycloheteroalkyl-containing groups has one or more hetero ring atoms selected from oxygen, sulfur and nitrogen atoms, and wherein each of R^0 and R^2 may be further independently selected from amino and amido radicals of the formula




wherein X is selected from oxygen atom or sulfur atom;

wherein each n is a number independently selected from zero to six, inclusive;

wherein each of R^{14} through R^{26} is independently selected from hydrido, alkyl, cycloalkyl, cyano, amino, monoalkylamino, dialkylamino, hydroxyalkyl, cycloalkylalkyl, alkoxyalkyl, aralkyl and aryl;

wherein each of R^3 through R^{11} is independently selected from hydrido, hydroxy, alkyl, hydroxyalkyl, halo, haloalkyl, cycloalkyl, alkoxy, aralkyl, aryl, aroyl, aryloxy, aralkoxy, alkoxyalkyl, alkylcarbonyl, alkoxyalkyl, alkenyl, cycloalkenyl, alkynyl, cyano, nitro, carboxyl, alkylcarbonyloxy, mercaptocarbonyl, alkoxyalkoxy, alkylthio, arylthio, aralkylthio, mercapto, alkylsulfonyl, aralkylsulfonyl and arylsulfonyl, and amino and amido radicals of the formula



wherein each of R^{14} , R^{15} , R^{16} , R^{17} , R^{18} and R^{19} is independently selected from hydrido, alkyl, cycloalkyl, cyano, amino, monoalkylamino, dialkylamino, hydroxyalkyl, cycloalkylalkyl, alkoxyalkyl, aralkyl and aryl;


and wherein each of R^0 and R^3 through R^{11} may be further independently selected from acidic moieties of the formula

-Y_nA


wherein n is a number selected from zero through three, inclusive;

wherein A is selected from carboxylic acid and bioisosteres of carboxylic acid selected from


-OH, -SH,



-S-WH,



wherein each W is independently selected from oxygen atom, sulfur atom and NR³⁹; wherein each of R³⁵, R³⁶, R³⁷, R³⁸ and R³⁹ is independently selected from hydrido, alkyl, haloalkyl, haloalkylsulfonyl, haloalkylcarbonyl, cycloalkyl, cycloalkylalkyl, aryl and aralkyl; wherein each of R³⁵, R³⁶, R³⁷ and R³⁹ may be further independently selected from amino radical of the formula




wherein each of R⁴⁰ and R⁴¹ is independently selected from hydrido, alkyl, cycloalkyl, hydroxyalkyl, haloalkyl, cycloalkylalkyl, alkoxyalkyl, aralkyl and aryl, and wherein R⁴⁰ and R⁴¹ taken together may form a heterocyclic group having five to seven ring members including the nitrogen atom of said amino radical, which heterocyclic group may further contain one or more hetero atoms as ring members selected from oxygen, nitrogen and sulfur atoms and which heterocyclic group may be saturated or partially unsaturated; wherein R⁴⁰ and R⁴¹ taken together may form an aromatic heterocyclic group having five ring members including the nitrogen atom of said amino radical and which aromatic heterocyclic group may further contain one or more hetero atoms as ring atoms selected from oxygen, nitrogen and sulfur atoms; wherein each of R³⁶ and R³⁷ may be further independently selected from hydroxy, alkoxy, alkylthio, aryloxy, arylthio, aralkylthio and aralkoxy; and the amide, ester and salt derivatives of said acidic groups;

wherein said bioisostere of carboxylic acid may be further selected from heterocyclic acidic groups consisting of heterocyclic rings of four to about nine ring members, which heterocyclic ring contains at least one hetero atom selected from oxygen, sulfur and nitrogen atoms, which heterocyclic ring may be saturated, fully unsaturated or partially unsaturated, and which heterocyclic ring may be attached at a single position selected from R³ through R¹¹ or may be attached at any two adjacent positions selected from R³ through R¹¹ so as to form a fused-ring system with one of the phenyl rings of Formula I; and the amide, ester and salt derivatives of said heterocyclic acidic groups;


wherein Y is a spacer group independently selected from one or more of alkyl, cycloalkyl, cycloalkylalkyl, alkenyl, aryl and aralkyl;

and wherein any of the foregoing R⁰ through R²⁶ and R³⁵ through R⁴¹, Y and A groups having a substitutable position may be substituted by one or more groups independently selected from hydroxy,

alkyl, alkenyl, aralkyl, hydroxyalkyl, halo, oxo, haloalkyl, alkoxy, aryloxy, aralkoxy, alkoxyalkyl, alkylcarbonyl, alkoxycarbonyl, carboxyl, cyano, nitro, alkylsulfonyl, haloalkylsulfonyl, aryl, aralkyl, mercaptocarbonyl, alkylthio and alkylthiocarbonyl, and amino and amido radicals of the formula



wherein X is selected from oxygen atom and sulfur atom; wherein R²⁷ is selected from hydrido, alkyl, cycloalkyl, cycloalkylalkyl, aralkyl, aryl and DR³² and




wherein D is selected from oxygen atom and sulfur atom; wherein R³² is selected from hydrido, alkyl, cycloalkyl, cycloalkylalkyl, aralkyl and aryl;

wherein each of R²⁷, R²⁸, R²⁹, R³⁰, R³¹, R³³ and R³⁴ is independently selected from hydrido, alkyl, cycloalkyl, cyano, hydroxyalkyl, haloalkyl, cycloalkylalkyl, alkoxyalkyl, alkanoyl, alkoxycarbonyl, carboxyl, haloalkylsulfinyl, haloalkylsulfonyl, aralkyl and aryl;

or a tautomer thereof or a pharmaceutically-acceptable salt thereof.


35. The composition of Claim 34 wherein m is one; wherein R¹ is selected from alkyl, hydroxyalkyl, haloalkyl, cycloalkyl, cycloalkylalkyl, thienylalkyl, phenylalkyl, polycycloalkyl, polycycloalkylalkyl, phenyl, halophenyl, alkylphenyl, alkoxyphenyl, cycloalkenyl, cycloalkenylalkyl, aroyl, alkoxyalkyl, alkylcarbonyl, cycloalkylcarbonyl, cycloalkylalkylcarbonyl, aralkylcarbonyl, alkenyl, cycloalkenyl, alkynyl, mercaptocarbonyl, alkylsulfonyl, aralkylsulfonyl, arylsulfonyl and amido radicals of the formula



wherein each of R¹² and R¹³ is independently selected from hydrido, alkyl, cycloalkyl, cyano, amino, monoalkylamino, dialkylamino, hydroxyalkyl, cycloalkylalkyl, alkoxyalkyl, aralkyl and aryl;

wherein each of R⁰ and R² is independently selected from hydrido, alkyl, hydroxyalkyl, formyl, halo, haloalkyl, cycloalkyl, cycloalkylalkyl, cycloalkylhaloalkyl, cycloalkylcarbonyl, alkoxy, thienylalkyl, phenylalkyl, polycycloalkyl, polycycloalkylalkyl, phenyl, halophenyl, alkylphenyl, alkoxyphenyl, cycloalkenyl, cycloalkenylalkyl, aralkylhaloalkyl, aroyl, aryloxy, aryloxyalkyl, aralkoxy, alkoxyalkyl, alkylcarbonyl, alkoxycarbonyl, alkenyl, cycloalkenyl, alkynyl, cycloalkynyl, cyano, nitro, carboxyl, carboxyalkyl, alkylcarbonyloxy, alkylcarbonyloxyalkyl, alkoxycarbonyloxy, aralkoxycarbonyloxy, aralkylcarbonyloxyalkyl, mercaptocarbonyl, mercaptoalkyl, alkoxycarbonyloxy, alkylthio, cycloalkylthio, cycloalkylalkylthio, arylthio, aralkylthio, mercapto, alkylsulfinyl, alkylsulfonyl, aralkylsulfinyl, aralkylsulfonyl, arylsulfinyl, arylsulfonyl, phthalimido, phthalimidoalkyl, heteroaryl, heteroarylalkyl, cycloheteroalkyl, cycloheteroalkylalkyl and cycloheteroalkylcarbonylalkyl wherein each of said heteroaryl- and cycloheteroalkyl-containing groups has one or more hetero ring atoms selected from oxygen, sulfur and nitrogen atoms, and wherein each of R² through R¹¹ may be further independently selected from amino and amido

radicals of the formula



wherein X is selected from oxygen atom and sulfur atom;

wherein each n is a number independently selected from zero to six, inclusive;

wherein each of R¹⁴ through R²⁶ is independently selected from hydrido, alkyl, cycloalkyl, cyano, amino, monoalkylamino, dialkylamino, hydroxyalkyl, cycloalkylalkyl, alkoxyalkyl, aralkyl and aryl;

wherein each of R³ through R¹¹ is independently selected from hydrido, hydroxy, alkyl, hydroxyalkyl, halo, haloalkyl, cycloalkyl, alkoxy, aralkyl, aryl, aroyl, aryloxy, aralkoxy, alkoxyalkyl, alkylcarbonyl, alkoxyalkyl, alkenyl, cycloalkenyl, alkynyl, cyano, nitro, carboxyl, alkylthio, aralkylthio and mercapto;

and wherein each of R⁰ and R³ through R¹¹ may be further independently selected from acidic moieties of the formula


-Y_nA

wherein n is a number selected from zero through three, inclusive; wherein A is selected from carboxylic acid and bioisosteres of carboxylic acid selected from

-OH, -SH,



wherein each W is independently selected from oxygen atom, sulfur atom and NR³⁹; wherein each of R³⁵, R³⁸ and R³⁹ is independently selected from hydrido, alkyl, haloalkyl, haloalkylsulfonyl, haloalkylcarbonyl, cycloalkyl, cycloalkylalkyl, aryl and aralkyl; wherein each of R³⁵ and R³⁹ may be further independently selected from amino radical of the formula



5

10

15

20

wherein each of R^{40} and R^{41} is independently selected from hydrido, alkyl, cycloalkyl, hydroxyalkyl, haloalkyl, cycloalkylalkyl, alkoxyalkyl, aralkyl and aryl, and wherein R^{40} and R^{41} taken together may form a heterocyclic group having five to seven ring members including the nitrogen atom of said amino radical, which heterocyclic group may further contain one or more hetero atoms as ring members selected from oxygen, nitrogen and sulfur atoms, and which heterocyclic group may be saturated or partially unsaturated; wherein R^{40} and R^{41} taken together may form an aromatic heterocyclic group having five ring members including the nitrogen atom of said amino radical and which aromatic heterocyclic group may further contain one or more hetero atoms as ring atoms selected from oxygen, nitrogen and sulfur atoms; and the amide, ester and salt derivatives of said acidic groups; wherein said bioisostere of carboxylic acid may be further selected from heterocyclic acidic groups consisting of heterocyclic rings of four to about nine ring members, which ring contains at least one hetero atom, selected from oxygen, sulfur and nitrogen atoms, which heterocyclic ring may be saturated, fully unsaturated or partially unsaturated, and which heterocyclic ring may be attached at a single position selected from R^3 through R^{11} or may be attached at any two adjacent positions selected from R^3 through R^{11} so as to form a fused-ring system with one of the phenyl rings of Formula I; and the amide, ester and salt derivatives of said heterocyclic acidic groups;

25

wherein Y is a spacer group independently selected from one or more of alkyl, cycloalkyl, cycloalkylalkyl, alkenyl, aryl and aralkyl;

30


wherein each of R^0 through R^{26} , R^{35} and R^{38} through R^{41} , Y and A independently may be substituted at any substitutable position with one or more groups selected from alkyl, hydroxy, halo, oxo, haloalkyl, alkoxy, aryloxy and aralkoxy;

or a tautomer thereof or a pharmaceutically-acceptable salt thereof.

35

36. The composition of Claim 35 wherein m is one; wherein R^1 is selected from alkyl, hydroxyalkyl, haloalkyl, cycloalkyl, cycloalkylalkyl, thienylalkyl, phenylalkyl, polycycloalkyl, polycycloalkylalkyl, phenyl, halophenyl, alkylphenyl, alkoxyphenyl, cycloalkenyl, cycloalkenylalkyl, aroyl, alkoxyalkyl, alkylcarbonyl, cycloalkylcarbonyl, cycloalkylalkylcarbonyl, aralkylcarbonyl, alkenyl, alkynyl, alkylsulfonyl, aralkylsulfonyl, arylsulfonyl and amido radicals of the formula

40



45


wherein each of R^{12} and R^{13} is independently selected from hydrido, alkyl, cycloalkyl, cyano, amino, hydroxyalkyl, alkoxyalkyl, phenalkyl and phenyl;

50

55

wherein each of R^0 and R^2 is independently selected from hydrido, alkyl, hydroxyalkyl, formyl, halo, haloalkyl, cycloalkyl, cycloalkylalkyl, cycloalkylhaloalkyl, cycloalkylcarbonyl, alkoxy, thienylalkyl, phenylalkyl, polycycloalkyl, polycycloalkylalkyl, phenyl, halophenyl, alkylphenyl, alkoxyphenyl, cycloalkenyl, cycloalkenylalkyl, aralkylhaloalkyl, benzoyl, phenoxy, phenoxyalkyl, phenalkyloxy, phenylthio, phenalkylthio, aralkoxy, alkoxyalkyl, alkylcarbonyl, alkoxyalkyl, alkenyl, cycloalkenyl, alkynyl, cyano, nitro, carboxyl, carboxyalkyl, alkylcarbonyloxy, alkylcarbonyloxyalkyl, alkoxyalkyl, aralkoxy, aralkoxyalkyl, aralkylcarbonyloxyalkyl, mercaptocarbonyl, mercaptoalkyl, alkoxyalkyl, alkoxyalkyl, cycloalkylthio, cycloalkylalkylthio, phthalimido, phthalimidoalkyl, heteroaryl, heteroarylalkyl, cycloheteroalkyl, cycloheteroalkylalkyl and cycloheteroalkylcarbonylalkyl wherein each of said heteroaryl- and cycloheteroalkyl-containing groups has one or more hetero ring atoms selected from

oxygen, sulfur and nitrogen atoms, and wherein each of R^0 and R^2 through R^{11} may be further independently selected from amino and amido radicals of the formula



wherein X is selected from oxygen atom and sulfur atom;

wherein each n is a number independently selected from zero to six, inclusive;




wherein each of R^{14} through R^{26} is independently selected from hydrido, alkyl, cycloalkyl, cyano, amino, hydroxyalkyl, alkoxyalkyl, phenalkyl and phenyl;

wherein each of R^3 through R^{11} is independently selected from hydrido, hydroxy, alkyl, hydroxyalkyl, halo, haloalkyl, cycloalkyl, alkoxy, phenalkyl, phenyl, benzoyl, phenoxy, phenalkyloxy, alkoxyalkyl, alkylcarbonyl, alkoxy carbonyl, alkenyl, cyano, nitro, carboxyl, alkylthio and mercapto;


and wherein each of R^0 and R^3 through R^{11} may be further independently selected from acidic moieties of the formula



wherein n is a number selected from zero through two, inclusive; wherein A is selected from carboxylic acid and bioisosteres of carboxylic acid selected from



wherein each W is independently selected from oxygen atom, sulfur atom and NR^{39} ; wherein each of R^{35} , R^{38} and R^{39} is independently selected from hydrido, alkyl, haloalkyl, haloalkylsulfonyl, haloalkylcarbonyl, cycloalkyl, phenyl and benzyl; wherein each of R^{35} and R^{39} may be further independently selected from amino radical of the formula



wherein each of R^{40} and R^{41} is independently selected from hydrido, alkyl, cycloalkyl, hydroxyalkyl, haloalkyl, alkoxyalkyl, benzyl and phenyl; and the amide, ester and salt derivatives of said acidic groups;

wherein said bioisostere of carboxylic acid may be further selected from heterocyclic acidic groups consisting of heterocyclic rings of four to about nine ring members, which ring contains at least one hetero atom, selected from oxygen, sulfur and nitrogen atoms, which heterocyclic ring may be saturated, fully unsaturated or partially unsaturated, and which heterocyclic ring may be attached at a single position selected from R^3 through R^{11} or may be attached at any two adjacent positions selected from R^3 through R^{11} so as to form a fused-ring system with one of the phenyl rings of Formula I; and the amide, ester and salt derivatives of said heterocyclic acidic groups;


wherein Y is a spacer group independently selected from one or more of alkyl, cycloalkyl, cycloalkylalkyl, alkenyl, phenyl, phenalkyl and aralkyl;

wherein each of R^0 through R^{26} , R^{35} and R^{38} through R^{41} , Y and A and independently may be substituted at any substitutable position with one or more groups selected from alkyl, cycloalkyl, cycloalkylalkyl, hydroxy, halo, oxo, haloalkyl, alkoxycarbonyl, cyano, nitro, alkylsulfonyl, haloalkylsulfonyl, aryl, aralkyl, alkoxy, aryloxy and aralkoxy;

or a tautomer thereof or a pharmaceutically-acceptable salt thereof.

37. The composition of Claim 36 wherein m is one; wherein R^1 is selected from alkyl, hydroxyalkyl, haloalkyl, cycloalkyl, cycloalkylalkyl, thienylalkyl, phenylalkyl, polycycloalkyl, polycycloalkylalkyl, phenyl, halophenyl, alkylphenyl, alkoxyphenyl, cycloalkenyl, cycloalkenylalkyl, benzoyl, alkoxyalkyl, alkylcarbonyl, alkoxycarbonyl, cycloalkylcarbonyl, cycloalkylalkylcarbonyl, aralkylcarbonyl, alkenyl and alkynyl;

where each of R^0 and R^2 is independently selected from hydrido, alkyl, aminoalkyl, hydroxyalkyl, formyl, halo, haloalkyl, cycloalkyl, cycloalkylalkyl, cycloalkylhaloalkyl, cycloalkylcarbonyl, alkoxy, thienylalkyl, phenylalkyl, polycycloalkyl, polycycloalkylalkyl, phenyl, halophenyl, alkylphenyl, alkoxyphenyl, cycloalkenyl, cycloalkenylalkyl, aralkylhaloalkyl, benzoyl, phenoxy, phenoxyalkyl, phenalkyloxy, phenylthio, phenalkylthio, aralkoxy, alkoxyalkyl, acetyl, alkoxycarbonyl, alkenyl, cycloalkenyl, alkynyl, cyano, nitro, carboxyl, carboxyalkyl, alkylcarbonyl, alkylcarbonyloxy, mercaptoalkyl, mercaptocarbonyl, alkoxycarbonyloxy, alkylcarbonyloxyalkyl, alkoxycarbonylalkyl, aralkoxycarbonylalkyl, aralkylcarbonyloxyalkyl, phthalimido, phthalimidoalkyl, imidazoalkyl, tetrazole, tetrazolealkyl, alkylthio, cycloalkylthio, cycloalkylalkylthio, and amino and amido radicals of the formula




wherein X is selected from oxygen atom and sulfur atom;

wherein each n is a number independently selected from zero to six, inclusive;


wherein each of R^{14} through R^{26} is independently selected from hydrido, alkyl, cycloalkyl, cyano, amino, hydroxyalkyl, alkoxyalkyl, phenalkyl and phenyl;

wherein each of R^3 through R^{11} is independently selected from hydrido, hydroxy, alkyl, hydroxyalkyl, halo, haloalkyl, alkoxy, phenyl, benzoyl, phenoxy, alkoxyalkyl, acetyl, alkoxyalkyl, alkenyl, cyano, nitro, carboxyl, alkylthio and mercapto;

and wherein each of R^0 and R^3 through R^{11} may be further independently selected from acidic moieties consisting of CO_2H , CO_2CH_3 , SH , CH_2SH , C_2H_4SH , PO_3H_2 , $NHSO_2CF_3$, $NHSO_2C_6F_5$, SO_3H , $CONHNH_2$, $CONHNHSO_2CF_3$, $CONHOCH_3$, $CONHOC_2H_5$, $CONHCF_3$, OH , CH_2OH , C_2H_4OH , OPO_3H_2 , OSO_3H ,



wherein each of R^{42} , R^{43} and R^{44} is independently selected from H, Cl, CN, NO_2 , CF_3 , C_2F_5 , C_3F_7 , CHF_2 , CH_2F , CO_2CH_3 , $CO_2C_2H_5$, SO_2CH_3 , SO_2CF_3 and $SO_2C_6F_5$; wherein Z is selected from O, S, NR^{45} and CH_2 ; wherein R^{45} is selected from hydrido, CH_3 and $CH_2C_6H_5$; and wherein said acidic moiety may be a heterocyclic acidic group attached at any two adjacent positions of R^3 through R^{11} so as to form a fused ring system with one of the phenyl rings of the biphenyl moiety of Formula I, said biphenyl fused ring system selected from



30 and the esters, amides and salts of said acidic moieties;
or a tautomer thereof or a pharmaceutically-acceptable salt thereof.

38. The composition of Claim 37 wherein m is one; wherein R¹ is selected from methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, n-pentyl, isopentyl, neopentyl, phenyl, benzyl, phenethyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cyclopropylmethyl, cyclopropylethyl, cyclobutylmethyl, cyclobutylethyl, cyclopentylmethyl, cyclopentylethyl, cyclohexylmethyl, cyclohexylethyl, cyclohexanoyl, methylbutyl, ethylbutyl, dimethylbutyl, thienylmethyl, thienylethyl, thienylpropyl, cyclopentenylmethyl, cyclopentenylethyl, cyclopentenylpropyl, cyclohexenyl, cyclohexenylmethyl, cyclohexenylethyl, adamantyl, adamantylmethyl, adamantylethyl, phenyl, chlorophenyl, dichlorophenyl, fluorophenyl, difluorophenyl, methoxyphenyl, ethoxyphenyl, methylphenyl, ethylphenyl, propylphenyl, isopropylphenyl, dimethylphenyl, diethylphenyl, benzyl, 1-oxo-2-cyclohexylethyl, benzoyl, 1-oxo-2-phenethyl, 1-oxoethyl, 1-oxopropyl, 1-oxobutyl, 1-oxopentyl, 2-butenyl, 3-butenyl, 2-butenyl, 3-butenyl, 2-butenyl, 3-butenyl and 2-hydroxybutyl; wherein R⁰ is selected from hydrido, C₄H₉(n), CH₃CH₂CH=CH, C₃H₇(n), SC₃H₇,

45




50





C₂H₅, C₅H₁₁(n), C₆H₁₃(n), SC₄H₉,

55




CH₃CH=CH, CH₃CH₂CH₂CH=CH-, amino, aminomethyl, aminoethyl, aminopropyl, CH₂OH, CH₂OCOCH₃, CH₂Cl, CH₂OCH₃, CH₂OCH(CH₃)₂, CHO,

CH₂CO₂H, CH(CH₃)CO₂H, NO₂, Cl,




20

-CO₂CH₃, -CONH₂, -CONHCH₃, CON(CH₃)₂, -CH₂-NHCO₂C₂H₅,




-CH₂NHCO₂CH₃, -CH₂NHCO₂C₃H₇, -CH₂NHCO₂CH₂(CH₃)₂, -CH₂NHCO₂C₄H₉, CH₂NHCO₂-adamantyl,
 -CH₂NHCO₂-(1-naphthyl), -CH₂NHCONHCH₃, -CH₂NHCONHC₂H₅, -CH₂NHCONHC₃H₇,
 -CH₂NHCONHC₄H₉, -CH₂NHCONHCH(CH₃)₂, -CH₂NHCONH(1-naphthyl), -CH₂NHCONH(1-adamantyl),




35

-CH₂CH₂CH₂CO₂H, -CH₂CH₂F, -CH₂OCONHCH₃, -CH₂OCSNHCH₃, -CH₂NHCSOC₃H₇, -CH₂CH₂CH₂F,
 -CH₂ONO₂,



45


-CH₂SH,



50


Cl, NO₂, CF₃, CH₂OH, Br, F, I, methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl,
 n-pentyl, isopentyl, neopentyl, phenyl, cyclohexyl, cyclohexylmethyl, carboxyl, formyl, 1-oxoethyl, 1-
 oxopropyl, 1-oxobutyl, 1-oxopentyl, dimethoxymethyl, 1,1-dimethoxypropyl, 1,1-dimethoxypentyl,
 hydroxyalkyl, halo, 1-oxo-2-phenylethyl, 1-oxo-2-cyclohexylethyl, 1,1-difluoro-2-phenylethyl, mon-
 ofluoromethyl, 1,1-difluoro-2-cyclohexylethyl, 2-cyclohexylethyl, 1,1-difluoro-3-cyclohexylpropyl, 1,1-
 55 dimethoxybutyl, 1,1-difluoroethyl, 1,1-difluoropropyl, 1,1-difluorobutyl, 1,1-difluoropentyl, benzyl, 2-
 phenylethyl, 1,1-difluoro-3-phenylpropyl, cyclohexylmethyl, cyclohexylethyl, cyclohexanoyl, 1-butenyl,
 2-butenyl, 3-butenyl, 1-butyryl, 2-butyryl, 3-butyryl, difluoromethyl, CO₂H, SH, PO₃H₂, SO₃H,
 CONHNH₂, CONHNHSO₂CF₃, OH,

5



10 wherein each of R⁴² and R⁴³ is independently selected from chloro, cyano, nitro, trifluoromethyl, methoxycarbonyl and trifluoromethylsulfonyl; wherein R² is selected from methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, n-pentyl, isopentyl, neopentyl, phenyl, benzyl, phenethyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cyclopropylmethyl, cyclopropylethyl, cyclobutylmethyl, cyclobutylethyl, cyclopentylmethyl, cyclopentylethyl, cyclohexylmethyl, cyclohexylethyl, methylbutyl, ethylbutyl, dimethylbutyl, thienylmethyl, thienylethyl, thienylpropyl, cyclopentenylmethyl, cyclopentenylethyl, cyclopentenylpropyl, cyclohexenyl, cyclohexenylmethyl, cyclohexenylethyl, adamantyl, adamantylmethyl, adamantylethyl, phenyl, chlorophenyl, dichlorophenyl, fluorophenyl, difluorophenyl, methoxyphenyl, ethoxyphenyl, methylphenyl, ethylphenyl, propylphenyl, isopropylphenyl, dimethylphenyl, diethylphenyl, benzyl, propylthio, butylthio and hydroxyalkyl; wherein each of R³ through R¹¹ is hydrido with the proviso that at least one of R⁵, R⁶, R⁸ and R⁹ is an acidic group selected from CO₂H, SH, PO₃H₂, SO₃H, CONHNH₂, CONHNHSO₂CF₃, OH,

25



30


wherein each of R⁴² and R⁴³ is independently selected from chloro, cyano, nitro, trifluoromethyl, methoxycarbonyl and trifluoromethylsulfonyl;

35

or a tautomer thereof or a pharmaceutically-acceptable salt thereof.


39. The composition of Claim 38 wherein m is one; wherein R¹ is selected from methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, 4-methylbutyl, n-pentyl, isopentyl, neopentyl, phenyl, benzyl, phenethyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cyclopropylmethyl, cyclopropylethyl, cyclobutylmethyl, cyclobutylethyl, cyclopentylmethyl, cyclopentylethyl, cyclohexylmethyl, cyclohexylethyl, cyclohexanoyl, methylbutyl, ethylbutyl, dimethylbutyl, thienylmethyl, thienylethyl, thienylpropyl, cyclopentenylmethyl, cyclopentenylethyl, cyclopentenylpropyl, cyclohexenyl, cyclohexenylmethyl, cyclohexenylethyl, adamantyl, adamantylmethyl, adamantylethyl, phenyl, chlorophenyl, dichlorophenyl, fluorophenyl, difluorophenyl, methoxyphenyl, ethoxyphenyl, methylphenyl, ethylphenyl, propylphenyl, isopropylphenyl, dimethylphenyl, diethylphenyl, benzyl, 1-oxo-2-cyclohexylethyl, benzoyl, 1-oxo-2-phenethyl, 1-oxoethyl, 1-oxopropyl, 1-oxobutyl, 1-oxopentyl, 2-butenyl, 3-butenyl, 2-butyryl, 3-butyryl and 2-hydroxybutyl; wherein R⁰ is selected from hydrido, methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, 4-methylbutyl, tert-butyl, n-pentyl, neopentyl, 1-oxoethyl, 1-oxopropyl, 1-oxobutyl, 1-oxopentyl, 1-oxo-2-phenylethyl, 1-oxo-2-cyclohexylethyl, 1,1-difluoro-2-phenylethyl, 1,1-difluoro-2-cyclohexylethyl, 2-cyclohexylethyl, 1,1-difluoro-3-cyclohexylpropyl, fluoro, chloro, monofluoromethyl, difluoromethyl, trifluoromethyl, formyl, carboxyl, dimethoxymethyl, 1,1-dimethoxybutyl, 1,1-difluoroethyl, 1,1-difluoropropyl, 1,1-difluorobutyl, 1,1-difluoropentyl, benzyl, 2-phenylethyl, 1,1-difluoro-3-phenylpropyl, cyclohexylmethyl, cyclohexylethyl, cyclohexanoyl, 1-butenyl, 2-butenyl, 3-butenyl, 1-butyryl, 2-butyryl, 3-butyryl, propylthio, butylthio, CO₂H, SH, PO₃H₂, SO₃H, CONHNH₂, CONHNHSO₂CF₃, OH,

5



10 wherein each of R⁴² and R⁴³ is independently selected from chloro, cyano, nitro, trifluoromethyl, methoxycarbonyl and trifluoromethylsulfonyl; wherein R² is selected from methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, n-pentyl, isopentyl, neopentyl, phenyl, benzyl, phenethyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cyclopropylmethyl, cyclopropylethyl, cyclobutylmethyl, cyclobutylethyl, cyclopentylmethyl, cyclopentylethyl, cyclohexylmethyl, cyclohexylethyl, methylbutyl, ethylbutyl, dimethylbutyl, thienylmethyl, thienylethyl, thienylpropyl, cyclopentenylmethyl, cyclopentenylethyl, cyclopentenylpropyl, cyclohexenyl, cyclohexenylmethyl, cyclohexenylethyl, adamantyl, adamantylmethyl, adamantylethyl, phenyl, chlorophenyl, dichlorophenyl, fluorophenyl, difluorophenyl, methoxyphenyl, ethoxyphenyl, methylphenyl, ethylphenyl, propylphenyl, isopropylphenyl, dimethylphenyl, diethylphenyl, benzyl, propylthio, butylthio and hydroxyalkyl; wherein each of R³ through R¹¹ is hydrido with the proviso that at least one of R⁵, R⁶, R⁸ and R⁹ is an acidic group selected from CO₂H, SH, PO₃H₂, SO₃H, CONHNH₂, CONHNHSO₂CF₃, OH,

25



30


wherein each of R⁴² and R⁴³ is independently selected from chloro, cyano, nitro, trifluoromethyl, methoxycarbonyl and trifluoromethylsulfonyl;

35

or a tautomer thereof or a pharmaceutically-acceptable salt thereof.

40. The composition of Claim 39 wherein m is one; wherein R¹ is selected from methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tertbutyl, 4-methylbutyl, n-pentyl, isopentyl, neopentyl, phenyl, benzyl, phenethyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cyclopropylmethyl, cyclopropylethyl, cyclobutylmethyl, cyclobutylethyl, cyclopentylmethyl, cyclopentylethyl, cyclohexylmethyl, cyclohexylethyl, cyclohexanoyl, methylbutyl, ethylbutyl, dimethylbutyl, thienylmethyl, thienylethyl, thienylpropyl, cyclopentenylmethyl, cyclopentenylethyl, cyclopentenylpropyl, cyclohexenyl, cyclohexenylmethyl, cyclohexenylethyl, adamantyl, adamantylmethyl, adamantylethyl, phenyl, chlorophenyl, dichlorophenyl, fluorophenyl, difluorophenyl, methoxyphenyl, ethoxyphenyl, methylphenyl, ethylphenyl, propylphenyl, isopropylphenyl, dimethylphenyl, diethylphenyl, benzyl, 1-oxo-2-cyclohexylethyl, benzoyl, 1-oxo-2-phenethyl, 1-oxopropyl, 1-oxobutyl, 1-oxopentyl and 2-hydroxybutyl; wherein R⁰ is selected from hydrido, methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, 4-methylbutyl, n-pentyl, 1-oxo-2-phenylethyl, 1-oxo-2-cyclohexylethyl, fluoro, chloro, monofluoromethyl, difluoromethyl, trifluoromethyl, formyl, carboxyl, dimethoxymethyl, 1,1-difluoro-2-phenylethyl, 1,1-difluoro-2-cyclohexylethyl, 2-cyclohexylethyl, 1,1-difluoro-3-cyclohexylpropyl, dimethoxymethyl, 1,1-dimethoxybutyl, 1,1-difluoroethyl, 1,1-difluoropropyl, 1,1-difluorobutyl, 1,1-difluoropentyl, benzyl, 2-phenylethyl, 1,1-difluoro-3-phenylpropyl, cyclohexylmethyl, cyclohexylethyl, cyclohexanoyl, 1-butenyl, 2-butenyl, 3-butenyl, 1-butylnyl, 2-butylnyl, 3-butylnyl, propylthio and butylthio; wherein R² is selected from methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, n-pentyl, isopentyl, neopentyl, phenyl, benzyl, phenethyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cyclopropylmethyl, cyclopropylethyl, cyclobutylmethyl, cyclobutylethyl, cyclopentylmethyl, cyclopentylethyl, cyclohexylmethyl, cyclohexylethyl, methylbutyl, ethylbutyl, dimethylbutyl, thienylmethyl, thienylethyl, thienylpropyl, cyclopentenylmethyl, cyclopentenylethyl, cyclopentenylpropyl, cyclohexenyl, cyclohex-


enylmethyl, cyclohexenylethyl, adamantyl, adamantylmethyl, adamantylethyl, phenyl, chlorophenyl, dichlorophenyl, fluorophenyl, difluorophenyl, methoxyphenyl, ethoxyphenyl, methylphenyl, ethylphenyl, propylphenyl, isopropylphenyl, dimethylphenyl, diethylphenyl, benzyl, propylthio, butylthio and hydroxyalkyl; wherein each of R^3 , R^4 , R^6 , R^7 , R^8 , R^{10} and R^{11} is hydrido; wherein one of R^5 and R^9 is hydrido and the other of R^5 and R^9 is an acidic group selected from COOH , SH , PO_3H_2 , SO_3H , CONHNH_2 , $\text{CONHNHSO}_2\text{CF}_3$, OH ,



wherein each of R^{42} and R^{43} is independently selected from chloro, cyano, nitro, trifluoromethyl, methoxycarbonyl and trifluoromethylsulfonyl;

or a tautomer thereof or a pharmaceutically-acceptable salt thereof.

41. The composition of Claim 40 wherein m is one; wherein R^1 is selected from methyl, ethyl, n -propyl, isopropyl, n -butyl, sec -butyl, isobutyl, $tert$ -butyl, 4-methylbutyl, n -pentyl, isopentyl, neopentyl, phenyl, benzyl, phenethyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cyclopropylmethyl, cyclopropylethyl, cyclobutylmethyl, cyclobutylethyl, cyclopentylmethyl, cyclopentylethyl, cyclohexylmethyl, cyclohexylethyl, cyclohexanoyl, methylbutyl, ethylbutyl, dimethylbutyl, thienylmethyl, thienylethyl, thienylpropyl, cyclopentenylmethyl, cyclopentenylethyl, cyclopentenylpropyl, cyclohexenyl, cyclohexenylmethyl, cyclohexenylethyl, adamantyl, adamantylmethyl, adamantylethyl, phenyl, chlorophenyl, dichlorophenyl, fluorophenyl, difluorophenyl, methoxyphenyl, ethoxyphenyl, methylphenyl, ethylphenyl, propylphenyl, isopropylphenyl, dimethylphenyl, diethylphenyl, benzyl, 1-oxo-2-cyclohexylethyl, benzoyl, 1-oxo-2-phenethyl, 1-oxopropyl, 1-oxobutyl, 1-oxopentyl and 2-hydroxybutyl; wherein R^0 is selected from hydrido, methyl, fluoro, chloro, monofluoromethyl, difluoromethyl, trifluoromethyl, formyl, carboxyl and dimethoxymethyl; wherein R^2 is selected from methyl, ethyl, n -propyl, isopropyl, n -butyl, sec -butyl, isobutyl, $tert$ -butyl, n -pentyl, isopentyl, neopentyl, phenyl, benzyl, phenethyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cyclopropylmethyl, cyclopropylethyl, cyclobutylmethyl, cyclobutylethyl, cyclopentylmethyl, cyclopentylethyl, cyclohexylmethyl, cyclohexylethyl, methylbutyl, ethylbutyl, dimethylbutyl, thienylmethyl, thienylethyl, thienylpropyl, cyclopentenylmethyl, cyclopentenylethyl, cyclopentenylpropyl, cyclohexenyl, cyclohexenylmethyl, cyclohexenylethyl, adamantyl, adamantylmethyl, adamantylethyl, phenyl, chlorophenyl, dichlorophenyl, fluorophenyl, difluorophenyl, methoxyphenyl, ethoxyphenyl, methylphenyl, ethylphenyl, propylphenyl, isopropylphenyl, dimethylphenyl, diethylphenyl, benzyl, propylthio, butylthio and hydroxyalkyl; wherein each of R^3 , R^4 , R^6 , R^7 , R^8 , R^{10} and R^{11} is hydrido; with the proviso that at least one of R^5 and R^9 must be selected from COOH , SH , PO_3H_2 , SO_3H , CONHNH_2 , $\text{CONHNHSO}_2\text{CF}_3$, OH ,




wherein each of R^{42} and R^{43} is independently selected from chloro, cyano, nitro, trifluoromethyl, methoxycarbonyl and trifluoromethylsulfonyl;

or a tautomer thereof or a pharmaceutically-acceptable salt thereof.


42. The composition of Claim 41 wherein m is one; wherein R^1 is selected from methyl, ethyl, n -propyl, isopropyl, n -butyl, sec -butyl, isobutyl, $tert$ -butyl, 4-methylbutyl, n -pentyl, isopentyl, neopentyl, phenyl, benzyl, phenethyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cyclopropylmethyl, cyclopropylethyl, cyclobutylmethyl, cyclobutylethyl, cyclopentylmethyl, cyclopentylethyl, cyclohexyl-

methyl, cyclohexylethyl cyclohexanoyl, methylbutyl, ethylbutyl, dimethylbutyl, thienylmethyl, thienylethyl, thienylpropyl, cyclopentenylmethyl, cyclopentenylethyl, cyclopentenylpropyl, cyclohexenyl, cyclohexenylmethyl, cyclohexenylethyl, adamantyl, adamantylmethyl, adamantylethyl, phenyl, chlorophenyl, dichlorophenyl, fluorophenyl, difluorophenyl, methoxyphenyl, ethoxyphenyl, methylphenyl, ethylphenyl, propylphenyl, isopropylphenyl, dimethylphenyl, diethylphenyl, benzyl, 1-oxo-2-cyclohexylethyl, benzoyl, 1-oxo-2-phenethyl, 1-oxopropyl, 1-oxobutyl, 1-oxopentyl and 2-hydroxybutyl; wherein R⁰ is selected from hydrido, methyl, fluoro, chloro, monofluoromethyl, difluoromethyl, trifluoromethyl, formyl, carboxyl and dimethoxymethyl; wherein R² is selected from methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, n-pentyl, isopentyl, neopentyl, phenyl, benzyl, phenethyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cyclopropylmethyl, cyclopropylethyl, cyclobutylmethyl, cyclobutylethyl, cyclopentylmethyl, cyclopentylethyl, cyclohexylmethyl, cyclohexylethyl, methylbutyl, ethylbutyl, dimethylbutyl, thienylmethyl, thienylethyl, thienylpropyl, cyclopentenylmethyl, cyclopentenylethyl, cyclopentenylpropyl, cyclohexenyl, cyclohexenylmethyl, cyclohexenylethyl, adamantyl, adamantylmethyl, adamantylethyl, phenyl, chlorophenyl, dichlorophenyl, fluorophenyl, difluorophenyl, methoxyphenyl, ethoxyphenyl, methylphenyl, ethylphenyl, propylphenyl, isopropylphenyl, dimethylphenyl, diethylphenyl, benzyl, propylthio, butylthio and hydroxyalkyl; wherein each of R³, R⁴, R⁶, R⁷, R⁸, R¹⁰ and R¹¹ is hydrido; wherein one of R⁵ and R⁹ is hydrido and the other of R⁵ and R⁹ is an acidic group selected from CO₂H and




or a tautomer thereof or a pharmaceutically-acceptable salt thereof.

43. The composition of Claim 42 wherein said antagonist compound is selected from a family of compounds of Formula II




wherein R¹ is selected from methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, 4-methylbutyl, n-pentyl, isopentyl, neopentyl, phenyl, benzyl, phenethyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cyclopropylmethyl, cyclopropylethyl, cyclobutylmethyl, cyclobutylethyl, cyclopentylmethyl, cyclopentylethyl, cyclohexylmethyl, cyclohexylethyl, cyclohexanoyl, methylbutyl, ethylbutyl, dimethylbutyl, thienylmethyl, thienylethyl, thienylpropyl, cyclopentenylmethyl, cyclopentenylethyl, cyclopentenylpropyl, cyclohexenyl, cyclohexenylmethyl, cyclohexenylethyl, adamantyl, adamantylmethyl, adamantylethyl, phenyl, chlorophenyl, dichlorophenyl, fluorophenyl, difluorophenyl, methoxyphenyl, ethoxyphenyl, methylphenyl, ethylphenyl, propylphenyl, isopropylphenyl, dimethylphenyl, diethylphenyl, benzyl, 1-oxo-2-cyclohexylethyl, benzoyl, 1-oxo-2-phenethyl, 1-oxopropyl, 1-oxobutyl, 1-oxopentyl and 2-hydroxybutyl; wherein R⁰ is selected from hydrido, methyl, fluoro, chloro, monofluoromethyl, difluoromethyl, trifluoromethyl, formyl, carboxyl and dimethoxymethyl; wherein R² is selected from methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, n-pentyl, isopentyl, neopentyl, phenyl, benzyl, phenethyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cyclopropylmethyl, cyclopropylethyl, cyclobutylmethyl, cyclobutylethyl, cyclopentylmethyl, cyclopentylethyl, cyclohexylmethyl, cyclohexylethyl, methylbutyl, ethylbutyl, dimethylbutyl, thienylmethyl, thienylethyl, thienylpropyl, cyclopentenylmethyl, cyclopentenylethyl, cyclopentenylpropyl, cyclohexenyl, cyclohexenylmethyl, cyclohexenylethyl, adamantyl, adamantylmethyl, adamantylethyl, phenyl, chlorophenyl,

dichlorophenyl, fluorophenyl, difluorophenyl, methoxyphenyl, ethoxyphenyl, methylphenyl, ethylphenyl, propylphenyl, isopropylphenyl, dimethylphenyl, diethylphenyl and benzyl; wherein R⁵ is an acidic group selected from CO₂H and



or a tautomer thereof or a pharmaceutically-acceptable salt thereof.

44. The composition of Claim 43 wherein R¹ is selected from methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, 4-methylbutyl, n-pentyl, isopentyl, neopentyl, phenyl, benzyl, phenethyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cyclopropylmethyl, cyclopropylethyl, cyclobutylmethyl, cyclobutylethyl, cyclopentylmethyl, cyclopentylethyl, cyclohexylmethyl, cyclohexylethyl, cyclohexanoyl, methylbutyl, ethylbutyl, dimethylbutyl, thienylmethyl, thienylethyl, thienylpropyl, cyclopentenylmethyl, cyclopentenylethyl, cyclopentenylpropyl, cyclohexenyl, cyclohexenylmethyl, cyclohexenylethyl, adamantyl, adamantylmethyl, adamantylethyl, phenyl, chlorophenyl, dichlorophenyl, fluorophenyl, difluorophenyl, methoxyphenyl, ethoxyphenyl, methylphenyl, ethylphenyl, propylphenyl, isopropylphenyl, dimethylphenyl, diethylphenyl, benzyl, 1-oxo-2-cyclohexylethyl, benzoyl, 1-oxo-2-phenethyl, 1-oxopropyl, 1-oxobutyl, 1-oxopentyl and 2-hydroxybutyl; wherein R⁰ is hydrido; wherein R² is selected from methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, n-pentyl, isopentyl, neopentyl, phenyl, benzyl, phenethyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cyclopropylmethyl, cyclopropylethyl, cyclobutylmethyl, cyclobutylethyl, cyclopentylmethyl, cyclopentylethyl, cyclohexylmethyl, cyclohexylethyl, methylbutyl, ethylbutyl, dimethylbutyl, thienylmethyl, thienylethyl, thienylpropyl, cyclopentenylmethyl, cyclopentenylethyl, cyclopentenylpropyl, cyclohexenyl, cyclohexenylmethyl, cyclohexenylethyl, adamantyl, adamantylmethyl, adamantylethyl, phenyl, chlorophenyl, dichlorophenyl, fluorophenyl, difluorophenyl, methoxyphenyl, ethoxyphenyl, methylphenyl, ethylphenyl, propylphenyl, isopropylphenyl, dimethylphenyl, diethylphenyl and benzyl; wherein R⁵ is an acidic group selected from CO₂H and




or a tautomer thereof or a pharmaceutically-acceptable salt thereof.

45. The composition of Claim 44 wherein said antagonist compound is selected from compounds and their pharmaceutically-acceptable salts of the group of compounds consisting of
- 1-propyl-4-butyl-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
 - 1,4-dibutyl-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
 - 1-pentyl-4-butyl-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
 - 1-isopentyl-4-butyl-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
 - 1-cyclohexylmethyl-4-butyl-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
 - 1-phenylmethyl-4-butyl-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
 - 1-(2-cyclohexylethyl)-4-butyl-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
 - 1-(2-phenylethyl)-4-butyl-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;


- 1-propyl-4-butyl-5-methyl-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
 1,4-dibutyl-5-methyl-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
 1-pentyl-4-butyl-5-methyl-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
 1-isopentyl-4-butyl-5-methyl-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
 1-cyclohexylmethyl-4-butyl-5-methyl-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
 1-phenylmethyl-4-butyl-5-methyl-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
 1-(2-cyclohexylethyl)-4-butyl-5-methyl-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
 1-(2-phenylethyl)-4-butyl-5-methyl-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
 1-propyl-4-butyl-5-chloro-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
 1,4-dibutyl-5-chloro-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
 1-pentyl-4-butyl-5-chloro-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
 1-isopentyl-4-butyl-5-chloro-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
 1-cyclohexylmethyl-4-butyl-5-chloro-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
 1-phenylmethyl-4-butyl-5-chloro-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
 1-(2-cyclohexylethyl)-4-butyl-5-chloro-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one; and
 1-(2-phenylethyl)-4-butyl-5-chloro-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one.

46. The composition of Claim 42 wherein said antagonist compound is selected from a family of compounds of Formula III




wherein R¹ is selected from methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, 4-methylbutyl, n-pentyl, isopentyl, neopentyl, phenyl, benzyl, phenethyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cyclopropylmethyl, cyclopropylethyl, cyclobutylmethyl, cyclobutylethyl, cyclopentylmethyl, cyclopentylethyl, cyclohexylmethyl, cyclohexylethyl, cyclohexanoyl, methylbutyl, ethylbutyl, dimethylbutyl, thienylmethyl, thienylethyl, thienylpropyl, cyclopentenylmethyl, cyclopentenylethyl, cyclopentenylpropyl, cyclohexenyl, cyclohexenylmethyl, cyclohexenylethyl, adamantyl, adamantylmethyl, adamantylethyl, phenyl, chlorophenyl, dichlorophenyl, fluorophenyl, difluorophenyl, methoxyphenyl, ethoxyphenyl, methylphenyl, ethylphenyl, propylphenyl, isopropylphenyl, dimethylphenyl, diethylphenyl, benzyl, 1-oxo-2-cyclohexylethyl, benzoyl, 1-oxo-2-phenethyl, 1-oxopropyl, 1-oxobutyl, 1-oxopentyl and 2-hydroxybutyl; wherein R⁰ is selected from hydrido, methyl, fluoro, chloro, monofluoromethyl, difluoromethyl, trifluoromethyl, formyl, carboxyl and dimethoxymethyl; wherein R² is selected from methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, n-pentyl, isopentyl, neopentyl, phenyl, benzyl, phenethyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cyclopropylmethyl, cyclopropylethyl, cyclobutylmethyl, cyclobutylethyl, cyclopentylmethyl, cyclopentylethyl, cyclohexylmethyl, cyclohexylethyl, methylbutyl, ethylbutyl, dimethylbutyl, thienylmethyl, thienylethyl, thienylpropyl, cyclopentenylmethyl, cyclopentenylethyl, cyclopentenylpropyl, cyclohexenyl, cyclohex-

enylmethyl, cyclohexenylethyl, adamantyl, adamantylmethyl, adamantylethyl, phenyl, chlorophenyl, dichlorophenyl, fluorophenyl, difluorophenyl, methoxyphenyl, ethoxyphenyl, methylphenyl, ethylphenyl, propylphenyl, isopropylphenyl, dimethylphenyl, diethylphenyl and benzyl; wherein R⁵ is an acidic group selected from CO₂H and



or a tautomer thereof or a pharmaceutically-acceptable salt thereof.

47. The composition of Claim 46 wherein R¹ is selected from methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, 4-methylbutyl, n-pentyl, isopentyl, neopentyl, phenyl, benzyl, phenethyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cyclopropylmethyl, cyclopropylethyl, cyclobutylmethyl, cyclobutylethyl, cyclopentylmethyl, cyclopentylethyl, cyclohexylmethyl, cyclohexylethyl, cyclohexanoyl, methylbutyl, ethylbutyl, dimethylbutyl, thienylmethyl, thienylethyl, thienylpropyl, cyclopentenylmethyl, cyclopentenylethyl, cyclopentenylpropyl, cyclohexenyl, cyclohexenylmethyl, cyclohexenylethyl, adamantyl, adamantylmethyl, adamantylethyl, phenyl, chlorophenyl, dichlorophenyl, fluorophenyl, difluorophenyl, methoxyphenyl, ethoxyphenyl, methylphenyl, ethylphenyl, propylphenyl, isopropylphenyl, dimethylphenyl, diethylphenyl, benzyl, 1-oxo-2-cyclohexylethyl, benzoyl, 1-oxo-2-phenethyl, 1-oxopropyl, 1-oxobutyl, 1-oxopentyl and 2-hydroxybutyl; wherein R⁰ is hydrido; wherein R² is selected from methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, n-pentyl, isopentyl, neopentyl, phenyl, benzyl, phenethyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cyclopropylmethyl, cyclopropylethyl, cyclobutylmethyl, cyclobutylethyl, cyclopentylmethyl, cyclopentylethyl, cyclohexylmethyl, cyclohexylethyl, methylbutyl, ethylbutyl, dimethylbutyl, thienylmethyl, thienylethyl, thienylpropyl, cyclopentenylmethyl, cyclopentenylethyl, cyclopentenylpropyl, cyclohexenyl, cyclohexenylmethyl, cyclohexenylethyl, adamantyl, adamantylmethyl, adamantylethyl, phenyl, chlorophenyl, dichlorophenyl, fluorophenyl, difluorophenyl, methoxyphenyl, ethoxyphenyl, methylphenyl, ethylphenyl, propylphenyl, isopropylphenyl, dimethylphenyl, diethylphenyl and benzyl; wherein R⁵ is an acidic group selected from CO₂H and



or a tautomer thereof or a pharmaceutically-acceptable salt thereof.

48. The composition of Claim 47 wherein said antagonist compound is selected from compounds and their pharmaceutically-acceptable salts of the group of compounds consisting of
- 1,4-dibutyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
- 1,4-dipropyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
- 1-propyl-4-isobutyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
- 1-butyl-4-isobutyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
- 1-butyl-4-isopentyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
- 1-butyl-4-pentyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;

- 1-butyl-4-cyclohexyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
- 5 1-butyl-4-cyclopentyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
- 1-butyl-4-(2-cyclopropylethyl)-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
- 10 1-methyl-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
- 1-ethyl-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
- 15 1-propyl-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
- 1-isobutyl-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
- 1-isopentyl-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
- 20 1-pentyl-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
- 1-isopropyl-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
- 25 1-tertbutyl-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
- 1-neopentyl-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
- 30 1-(3,3-dimethylbutyl)-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
- 1-(2-ethylbutyl)-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
- 35 1-cyclohexyl-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
- 1-(2-cyclopropylethyl)-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
- 40 1-cyclopentyl-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
- 1-cyclopentylmethyl-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
- 45 1-[2-(2-thienyl)ethyl]-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
- 1-(2-phenylethyl)-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
- 50 1-(2-cyclopentylethyl)-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
- 1-[2-(cyclopenten-1-yl)ethyl]-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
- 55 1-(2-adamantyl)-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;

5

10

15

20

25

30




40

45


50

55



or a tautomer thereof or a pharmaceutically-acceptable salt thereof.

50. The composition of Claim 49 wherein R¹ is selected from methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, 4-methylbutyl, n-pentyl, isopentyl, neopentyl, phenyl, benzyl, phenethyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cyclopropylmethyl, cyclopropylethyl, cyclobutylmethyl, cyclobutylethyl, cyclopentylmethyl, cyclopentylethyl, cyclohexylmethyl, cyclohexylethyl, cyclohexanoyl, methylbutyl, ethylbutyl, dimethylbutyl, thienylmethyl, thienylethyl, thienylpropyl, cyclopentenylmethyl, cyclopentenylethyl, cyclopentenylpropyl, cyclohexenyl, cyclohexenylmethyl, cyclohexenylethyl, adamantyl, adamantylmethyl, adamantylethyl, phenyl, chlorophenyl, dichlorophenyl, fluorophenyl, difluorophenyl, methoxyphenyl, ethoxyphenyl, methylphenyl, ethylphenyl, propylphenyl, isopropylphenyl, dimethylphenyl, diethylphenyl, benzyl, 1-oxo-2-cyclohexylethyl, benzoyl, 1-oxo-2-phenethyl, 1-oxopropyl, 1-oxobutyl, 1-oxopentyl and 2-hydroxybutyl; wherein R⁰ is hydrido; wherein R² is selected from methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, n-pentyl, isopentyl, neopentyl, phenyl, benzyl, phenethyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cyclopropylmethyl, cyclopropylethyl, cyclobutylmethyl, cyclobutylethyl, cyclopentylmethyl, cyclopentylethyl, cyclohexylmethyl, cyclohexylethyl, methylbutyl, ethylbutyl, dimethylbutyl, thienylmethyl, thienylethyl, thienylpropyl, cyclopentenylmethyl, cyclopentenylethyl, cyclopentenylpropyl, cyclohexenyl, cyclohexenylmethyl, cyclohexenylethyl, adamantyl, adamantylmethyl, adamantylethyl, phenyl, chlorophenyl, dichlorophenyl, fluorophenyl, difluorophenyl, methoxyphenyl, ethoxyphenyl, methylphenyl, ethylphenyl, propylphenyl, isopropylphenyl, dimethylphenyl, diethylphenyl and benzyl; wherein R⁵ is an acidic group selected from CO₂H and




or a tautomer thereof or a pharmaceutically-acceptable salt thereof.


51. The composition of Claim 50 wherein said antagonist compound is selected from compounds and their pharmaceutically-acceptable salts of the group of compounds consisting of
- 1-propyl-4-butyl-1,3-dihydro-3-[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]-2H-imidazol-2-one;
 - 1,4-dibutyl-1,3-dihydro-3-[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]-2H-imidazol-2-one;
 - 1-pentyl-4-butyl-1,3-dihydro-3-[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]-2H-imidazol-2-one;
 - 1-isopentyl-4-butyl-1,3-dihydro-3-[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]-2H-imidazol-2-one;
 - 1-cyclohexylmethyl-4-butyl-1,3-dihydro-3-[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]-2H-imidazol-2-one;
 - 1-phenylmethyl-4-butyl-1,3-dihydro-3-[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]-2H-imidazol-2-one;
 - 1-(2-cyclohexylethyl)-4-butyl-1,3-dihydro-3-[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]-2H-imidazol-2-one;
 - 1-(2-phenylethyl)-4-butyl-1,3-dihydro-3-[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]-2H-imidazol-2-one;
 - 1-propyl-4-butyl-5-methyl-1,3-dihydro-3-[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]-2H-imidazol-2-one;
 - 1,4-dibutyl-5-methyl-1,3-dihydro-3-[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]-2H-imidazol-2-one;
 - 1-pentyl-4-butyl-5-methyl-1,3-dihydro-3-[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]-2H-imidazol-2-one;
 - 1-isopentyl-4-butyl-5-methyl-1,3-dihydro-3-[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]-2H-imidazol-2-one;
 - 1-cyclohexylmethyl-4-butyl-5-methyl-1,3-dihydro-3-[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]-2H-imidazol-2-one;
 - 1-phenylmethyl-4-butyl-5-methyl-1,3-dihydro-3-[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]-2H-imidazol-2-one;

1-(2-cyclohexylethyl)-4-butyl-5-methyl-1,3-dihydro-3-[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]-2H-imidazol-2-one;
 1-(2-phenylethyl)-4-butyl-5-methyl-1,3-dihydro-3-[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]-2H-imidazol-2-one;
 1-propyl-4-butyl-5-chloro-1,3-dihydro-3-[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]-2H-imidazol-2-one;
 1,4-dibutyl-5-chloro-1,3-dihydro-3-[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]-2H-imidazol-2-one;
 1-pentyl-4-butyl-5-chloro-1,3-dihydro-3-[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]-2H-imidazol-2-one;
 1-isopentyl-4-butyl-5-chloro-1,3-dihydro-3-[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]-2H-imidazol-2-one;
 1-cyclohexylmethyl-4-butyl-5-chloro-1,3-dihydro-3-[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]-2H-imidazol-2-one;
 1-phenylmethyl-4-butyl-5-chloro-1,3-dihydro-3-[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]-2H-imidazol-2-one;
 1-(2-cyclohexylethyl)-4-butyl-5-chloro-1,3-dihydro-3-[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]-2H-imidazol-2-one; and
 1-(2-phenylethyl)-4-butyl-5-chloro-1,3-dihydro-3-[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]-2H-imidazol-2-one.

52. The composition of Claim 42 wherein said antagonist compound is selected from a family of compounds of Formula V




wherein R¹ is selected from methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, 4-methylbutyl, n-pentyl, isopentyl, neopentyl, phenyl, benzyl, phenethyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cyclopropylmethyl, cyclopropylethyl, cyclobutylmethyl, cyclobutylethyl, cyclopentylmethyl, cyclopentylethyl, cyclohexylmethyl, cyclohexylethyl, cyclohexanoyl, methylbutyl, ethylbutyl, dimethylbutyl, thienylmethyl, thienylethyl, thienylpropyl, cyclopentenylmethyl, cyclopentenylethyl, cyclopentenylpropyl, cyclohexenyl, cyclohexenylmethyl, cyclohexenylethyl, adamantyl, adamantylmethyl, adamantylethyl, phenyl, chlorophenyl, dichlorophenyl, fluorophenyl, difluorophenyl, methoxyphenyl, ethoxyphenyl, methylphenyl, ethylphenyl, propylphenyl, isopropylphenyl, dimethylphenyl, diethylphenyl, benzyl, 1-oxo-2-cyclohexylethyl, benzoyl, 1-oxo-2-phenethyl, 1-oxopropyl, 1-oxobutyl, 1-oxopentyl and 2-hydroxybutyl; wherein R⁰ is selected from hydrido, methyl, fluoro, chloro, monofluoromethyl, difluoromethyl, trifluoromethyl, formyl, carboxyl and dimethoxymethyl; wherein R² is selected from methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, n-pentyl, isopentyl, neopentyl, phenyl, benzyl, phenethyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cyclopropylmethyl, cyclopropylethyl, cyclobutylmethyl, cyclobutylethyl, cyclopentylmethyl, cyclopentylethyl, cyclohexylmethyl, cyclohexylethyl, methylbutyl, ethylbutyl, dimethylbutyl, thienylmethyl, thienylethyl, thienylpropyl, cyclopentenylmethyl, cyclopentenylethyl, cyclopentenylpropyl, cyclohexenyl, cyclohexenylmethyl, cyclohexenylethyl, adamantyl, adamantylmethyl, adamantylethyl, phenyl, chlorophenyl, dichlorophenyl, fluorophenyl, difluorophenyl, methoxyphenyl, ethoxyphenyl, methylphenyl, ethylphenyl, propylphenyl, isopropylphenyl, dimethylphenyl, diethylphenyl and benzyl; wherein R⁵ is an acidic group selected from CO₂H and



or a tautomer thereof or a pharmaceutically-acceptable salt thereof.

53. The composition of Claim 52 wherein R¹ is selected from methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, 4-methylbutyl, n-pentyl, isopentyl, neopentyl, phenyl, benzyl, phenethyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cyclopropylmethyl, cyclopropylethyl, cyclobutylmethyl, cyclobutylethyl, cyclopentylmethyl, cyclopentylethyl, cyclohexylmethyl, cyclohexylethyl, cyclohexanoyl, methylbutyl, ethylbutyl, dimethylbutyl, thienylmethyl, thienylethyl, thienylpropyl, cyclopentenylmethyl, cyclopentenylethyl, cyclopentenylpropyl, cyclohexenyl, cyclohexenylmethyl, cyclohexenylethyl, adamantyl, adamantylmethyl, adamantylethyl, phenyl, chlorophenyl, dichlorophenyl, fluorophenyl, difluorophenyl, methoxyphenyl, ethoxyphenyl, methylphenyl, ethylphenyl, propylphenyl, isopropylphenyl, dimethylphenyl, diethylphenyl, benzyl, 1-oxo-2-cyclohexylethyl, benzoyl, 1-oxo-2-phenethyl, 1-oxopropyl, 1-oxobutyl, 1-oxopentyl and 2-hydroxybutyl; wherein R⁰ is hydrido; wherein R² is selected from methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, n-pentyl, isopentyl, neopentyl, phenyl, benzyl, phenethyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cyclopropylmethyl, cyclopropylethyl, cyclobutylmethyl, cyclobutylethyl, cyclopentylmethyl, cyclopentylethyl, cyclohexylmethyl, cyclohexylethyl, methylbutyl, ethylbutyl, dimethylbutyl, thienylmethyl, thienylethyl, thienylpropyl, cyclopentenylmethyl, cyclopentenylethyl, cyclopentenylpropyl, cyclohexenyl, cyclohexenylmethyl, cyclohexenylethyl, adamantyl, adamantylmethyl, adamantylethyl, phenyl, chlorophenyl, dichlorophenyl, fluorophenyl, difluorophenyl, methoxyphenyl, ethoxyphenyl, methylphenyl, ethylphenyl, propylphenyl, isopropylphenyl, dimethylphenyl, diethylphenyl and benzyl; wherein R⁵ is an acidic group selected from CO₂H and



or a tautomer thereof or a pharmaceutically-acceptable salt thereof.

54. The composition of Claim 53 wherein said antagonist compound is selected from compounds and their pharmaceutically-acceptable salts of the group of compounds consisting of

1-propyl-4-butyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

1,4-dibutyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

1-pentyl-4-butyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

1-isopentyl-4-butyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

1-cyclohexylmethyl-4-butyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

1-phenylmethyl-4-butyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

1-(2-cyclohexylethyl)-4-butyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

1-(2-phenylethyl)-4-butyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

1-propyl-4-butyl-5-methyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

1,4-dibutyl-5-methyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

1-pentyl-4-butyl-5-methyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

1-isopentyl-4-butyl-5-methyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

1-cyclohexylmethyl-4-butyl-5-methyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

1-phenylmethyl-4-butyl-5-methyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

1-(2-cyclohexylethyl)-4-butyl-5-methyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

1-(2-phenylethyl)-4-butyl-5-methyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-

imidazol-2-one;

1-propyl-4-butyl-5-chloro-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

1,4-dibutyl-5-chloro-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

1-pentyl-4-butyl-5-chloro-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

1-isopentyl-4-butyl-5-chloro-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;


1-cyclohexylmethyl-4-butyl-5-chloro-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

1-phenylmethyl-4-butyl-5-chloro-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;


1-(2-cyclohexylethyl)-4-butyl-5-chloro-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one; and

1-(2-phenylethyl)-4-butyl-5-chloro-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one.

55. The composition of Claim 42 wherein said antagonist compound is selected from a family of compounds of Formula VI




wherein R¹ is selected from methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, 4-methylbutyl, n-pentyl, isopentyl, neopentyl, phenyl, benzyl, phenethyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cyclopropylmethyl, cyclopropylethyl, cyclobutylmethyl, cyclobutylethyl, cyclopentylmethyl, cyclopentylethyl, cyclohexylmethyl, cyclohexylethyl, cyclohexanoyl, methylbutyl, ethylbutyl, dimethylbutyl, thienylmethyl, thienylethyl, thienylpropyl, cyclopentenylmethyl, cyclopentenylethyl, cyclopentenylpropyl, cyclohexenyl, cyclohexenylmethyl, cyclohexenylethyl, adamantyl, adamantylmethyl, adamantylethyl, phenyl, chlorophenyl, dichlorophenyl, fluorophenyl, difluorophenyl, methoxyphenyl, ethoxyphenyl, methylphenyl, ethylphenyl, propylphenyl, isopropylphenyl, dimethylphenyl, diethylphenyl, benzyl, 1-oxo-2-cyclohexylethyl, benzoyl, 1-oxo-2-phenethyl, 1-oxopropyl, 1-oxobutyl, 1-oxopentyl and 2-hydroxybutyl; wherein R⁰ is selected from hydrido, methyl, fluoro, chloro, monofluoromethyl, difluoromethyl, trifluoromethyl, formyl, carboxyl and dimethoxymethyl; wherein R² is selected from methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, n-pentyl, isopentyl, neopentyl, phenyl, benzyl, phenethyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cyclopropylmethyl, cyclopropylethyl, cyclobutylmethyl, cyclobutylethyl, cyclopentylmethyl, cyclopentylethyl, cyclohexylmethyl, cyclohexylethyl, methylbutyl, ethylbutyl, dimethylbutyl, thienylmethyl, thienylethyl, thienylpropyl, cyclopentenylmethyl, cyclopentenylethyl, cyclopentenylpropyl, cyclohexenyl, cyclohexenylmethyl, cyclohexenylethyl, adamantyl, adamantylmethyl, adamantylethyl, phenyl, chlorophenyl, dichlorophenyl, fluorophenyl, difluorophenyl, methoxyphenyl, ethoxyphenyl, methylphenyl, ethylphenyl, propylphenyl, isopropylphenyl, dimethylphenyl, diethylphenyl and benzyl; wherein R⁵ is an acidic group selected from CO₂H and



or a tautomer thereof or a pharmaceutically-acceptable salt thereof.

56. The composition of Claim 55 wherein R¹ is selected from methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, 4-methylbutyl, n-pentyl, isopentyl, neopentyl, phenyl, benzyl, phenethyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cyclopropylmethyl, cyclopropylethyl, cyclobutylmethyl, cyclobutylethyl, cyclopentylmethyl, cyclopentylethyl, cyclohexylmethyl, cyclohexylethyl, cyclohexanoyl, methylbutyl, ethylbutyl, dimethylbutyl, thienylmethyl, thienylethyl, thienylpropyl, cyclopentenylmethyl, cyclopentenylethyl, cyclopentenylpropyl, cyclohexenyl, cyclohexenylmethyl, cyclohexenylethyl, adamantyl, adamantylmethyl, adamantylethyl, phenyl, chlorophenyl, dichlorophenyl, fluorophenyl, difluorophenyl, methoxyphenyl, ethoxyphenyl, methylphenyl, ethylphenyl, propylphenyl, isopropylphenyl, dimethylphenyl, diethylphenyl, benzyl, 1-oxo-2-cyclohexylethyl, benzoyl, 1-oxo-2-phenethyl, 1-oxopropyl, 1-oxobutyl, 1-oxopentyl and 2-hydroxybutyl; wherein R⁰ is hydrido; wherein R² is selected from methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, n-pentyl, isopentyl, neopentyl, phenyl, benzyl, phenethyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cyclopropylmethyl, cyclopropylethyl, cyclobutylmethyl, cyclobutylethyl, cyclopentylmethyl, cyclopentylethyl, cyclohexylmethyl, cyclohexylethyl, methylbutyl, ethylbutyl, dimethylbutyl, thienylmethyl, thienylethyl, thienylpropyl, cyclopentenylmethyl, cyclopentenylethyl, cyclopentenylpropyl, cyclohexenyl, cyclohexenylmethyl, cyclohexenylethyl, adamantyl, adamantylmethyl, adamantylethyl, phenyl, chlorophenyl, dichlorophenyl, fluorophenyl, difluorophenyl, methoxyphenyl, ethoxyphenyl, methylphenyl, ethylphenyl, propylphenyl, isopropylphenyl, dimethylphenyl, diethylphenyl and benzyl; wherein R⁵ is an acidic group selected from CO₂H and



or a tautomer thereof or a pharmaceutically-acceptable salt thereof.

57. The composition of Claim 56 wherein said antagonist compound is selected from compounds and their pharmaceutically-acceptable salts of the group of compounds consisting of

- 1-propyl-4-butyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
- 1,4-dibutyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
- 1-pentyl-4-butyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
- 1-isopentyl-4-butyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
- 1-cyclohexylmethyl-4-butyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
- 1-phenylmethyl-4-butyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
- 1-(2-cyclohexylethyl)-4-butyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
- 1-(2-phenylethyl)-4-butyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
- 1-propyl-4-butyl-5-methyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
- 1,4-dibutyl-5-methyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
- 1-pentyl-4-butyl-5-methyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
- 1-isopentyl-4-butyl-5-methyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
- 1-cyclohexylmethyl-4-butyl-5-methyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
- 1-phenylmethyl-4-butyl-5-methyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
- 1-(2-cyclohexylethyl)-4-butyl-5-methyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
- 1-(2-phenylethyl)-4-butyl-5-methyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

imidazol-2-one;

1-propyl-4-butyl-5-chloro-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

1,4-dibutyl-5-chloro-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

1-pentyl-4-butyl-5-chloro-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

1-isopentyl-4-butyl-5-chloro-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;


1-cyclohexylmethyl-4-butyl-5-chloro-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

1-phenylmethyl-4-butyl-5-chloro-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;


1-(2-cyclohexylethyl)-4-butyl-5-chloro-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one; and

1-(2-phenylethyl)-4-butyl-5-chloro-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one.

58. The composition of Claim 42 wherein said antagonist compound is selected from a family of compounds of Formula VII




wherein R¹ is selected from methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, 4-methylbutyl, n-pentyl, isopentyl, neopentyl, phenyl, benzyl, phenethyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cyclopropylmethyl, cyclopropylethyl, cyclobutylmethyl, cyclobutylethyl, cyclopentylmethyl, cyclopentylethyl, cyclohexylmethyl, cyclohexylethyl, cyclohexanoyl, methylbutyl, ethylbutyl, dimethylbutyl, thienylmethyl, thienylethyl, thienylpropyl, cyclopentenylmethyl, cyclopentenylethyl, cyclopentenylpropyl, cyclohexenyl, cyclohexenylmethyl, cyclohexenylethyl, adamantyl, adamantylmethyl, adamantylethyl, phenyl, chlorophenyl, dichlorophenyl, fluorophenyl, difluorophenyl, methoxyphenyl, ethoxyphenyl, methylphenyl, ethylphenyl, propylphenyl, isopropylphenyl, dimethylphenyl, diethylphenyl, benzyl, 1-oxo-2-cyclohexylethyl, benzoyl, 1-oxo-2-phenethyl, 1-oxopropyl, 1-oxobutyl, 1-oxopentyl and 2-hydroxybutyl; wherein R⁰ is selected from hydrido, methyl, fluoro, chloro, monofluoromethyl, difluoromethyl, trifluoromethyl, formyl, carboxyl and dimethoxymethyl; wherein R² is selected from methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, n-pentyl, isopentyl, neopentyl, phenyl, benzyl, phenethyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cyclopropylmethyl, cyclopropylethyl, cyclobutylmethyl, cyclobutylethyl, cyclopentylmethyl, cyclopentylethyl, cyclohexylmethyl, cyclohexylethyl, methylbutyl, ethylbutyl, dimethylbutyl, thienylmethyl, thienylethyl, thienylpropyl, cyclopentenylmethyl, cyclopentenylethyl, cyclopentenylpropyl, cyclohexenyl, cyclohexenylmethyl, cyclohexenylethyl, adamantyl, adamantylmethyl, adamantylethyl, phenyl, chlorophenyl, dichlorophenyl, fluorophenyl, difluorophenyl, methoxyphenyl, ethoxyphenyl, methylphenyl, ethylphenyl, propylphenyl, isopropylphenyl, dimethylphenyl, diethylphenyl and benzyl; wherein R⁵ is an acidic group selected from CO₂H and



or a tautomer thereof or a pharmaceutically-acceptable salt thereof.

59. The composition of Claim 58 wherein R¹ is selected from methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, 4-methylbutyl, n-pentyl, isopentyl, neopentyl, phenyl, benzyl, phenethyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cyclopropylmethyl, cyclopropylethyl, cyclobutylmethyl, cyclobutylethyl, cyclopentylmethyl, cyclopentylethyl, cyclohexylmethyl, cyclohexylethyl, cyclohexanoyl, methylbutyl, ethylbutyl, dimethylbutyl, thienylmethyl, thienylethyl, thienylpropyl, cyclopentenylmethyl, cyclopentenylethyl, cyclopentenylpropyl, cyclohexenyl, cyclohexenylmethyl, cyclohexenylethyl, adamantyl, adamantylmethyl, adamantylethyl, phenyl, chlorophenyl, dichlorophenyl, fluorophenyl, difluorophenyl, methoxyphenyl, ethoxyphenyl, methylphenyl, ethylphenyl, propylphenyl, isopropylphenyl, dimethylphenyl, diethylphenyl, benzyl, 1-oxo-2-cyclohexylethyl, benzoyl, 1-oxo-2-phenethyl, 1-oxopropyl, 1-oxobutyl, 1-oxopentyl and 2-hydroxybutyl; wherein R⁰ is hydrido; wherein R² is selected from methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, n-pentyl, isopentyl, neopentyl, phenyl, benzyl, phenethyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cyclopropylmethyl, cyclopropylethyl, cyclobutylmethyl, cyclobutylethyl, cyclopentylmethyl, cyclopentylethyl, cyclohexylmethyl, cyclohexylethyl, methylbutyl, ethylbutyl, dimethylbutyl, thienylmethyl, thienylethyl, thienylpropyl, cyclopentenylmethyl, cyclopentenylethyl, cyclopentenylpropyl, cyclohexenyl, cyclohexenylmethyl, cyclohexenylethyl, adamantyl, adamantylmethyl, adamantylethyl, phenyl, chlorophenyl, dichlorophenyl, fluorophenyl, difluorophenyl, methoxyphenyl, ethoxyphenyl, methylphenyl, ethylphenyl, propylphenyl, isopropylphenyl, dimethylphenyl, diethylphenyl and benzyl; wherein R⁵ is an acidic group selected from CO₂H and



or a tautomer thereof or a pharmaceutically-acceptable salt thereof.

60. The composition of Claim 59 wherein said antagonist compound is selected from compounds and their pharmaceutically-acceptable salts of the group of compounds consisting of
- 1-propyl-4-butyl-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 - 1,4-dibutyl-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 - 1-pentyl-4-butyl-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 - 1-isopentyl-4-butyl-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 - 1-cyclohexylmethyl-4-butyl-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 - 1-phenylmethyl-4-butyl-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 - 1-(2-cyclohexylethyl)-4-butyl-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 - 1-(2-phenylethyl)-4-butyl-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 - 1-propyl-4-butyl-5-methyl-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 - 1,4-dibutyl-5-methyl-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 - 1-pentyl-4-butyl-5-methyl-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 - 1-isopentyl-4-butyl-5-methyl-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 - 1-cyclohexylmethyl-4-butyl-5-methyl-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 - 1-phenylmethyl-4-butyl-5-methyl-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 - 1-(2-cyclohexylethyl)-4-butyl-5-methyl-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 - 1-(2-phenylethyl)-4-butyl-5-methyl-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-

imidazol-2-one;

1-propyl-4-butyl-5-chloro-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

1,4-dibutyl-5-chloro-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

5 1-pentyl-4-butyl-5-chloro-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

1-isopentyl-4-butyl-5-chloro-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

10 1-cyclohexylmethyl-4-butyl-5-chloro-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

1-phenylmethyl-4-butyl-5-chloro-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;


1-(2-cyclohexylethyl)-4-butyl-5-chloro-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one; and

15 1-(2-phenylethyl)-4-butyl-5-chloro-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one.


61. The composition of Claim 48 wherein said antagonist compound is 1,4-dibutyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one or a pharmaceutically-acceptable salt thereof.

62. The composition of Claim 48 wherein said antagonist compound is 1-(2-ethylbutyl)-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one or a pharmaceutically-acceptable salt thereof.

63. A therapeutic method for treating a circulatory disorder, said method comprising administering to a subject susceptible to or afflicted with such disorder a therapeutically-effective amount of an active compound of Formula I:




wherein A is selected from



wherein m is a number selected from one to four, inclusive;


wherein R¹ is selected from alkyl, hydroxyalkyl, haloalkyl, cycloalkyl, cycloalkylalkyl, formyl, thienylalkyl, phenylalkyl, polycycloalkyl, polycycloalkylalkyl, phenyl, halophenyl, alkylphenyl, alkoxyphenyl, cycloalkenyl, cycloalkenylalkyl, aroyl, alkoxyalkyl, alkylcarbonyl, alkylcarbonylalkyl, cycloalkylcarbonyl, cycloalkylalkylcarbonyl, aralkylcarbonyl, alkoxyalkyl, alkenyl, cycloalkenyl, aralkoxyalkyl, alkynyl, alkylthiocarbonyl, alkylthiothiocarbonyl, arylthiocarbonyl, arylthiothiocarbonyl, aralkylthiocarbonyl, alkylthiocarbonyl, alkylsulfinyl, alkylsulfonyl, aralkylsulfinyl, aralkylsulfonyl, arylsulfinyl, arylsulfonyl, heteroaryl having one or more ring atoms selected from oxygen, sulfur and nitrogen atoms, and amido radicals of the formula



wherein X is oxygen atom or sulfur atom;

wherein each of R¹² and R¹³ is independently selected from hydrido, alkyl, cycloalkyl, cyano, amino, monoalkylamino, dialkylamino, hydroxyalkyl, cycloalkylalkyl, alkoxyalkyl, aralkyl and aryl, and wherein R¹² and R¹³ taken together may form a heterocyclic group having five to seven ring members including the nitrogen atom of said amido radical and which heterocyclic group may further contain one or more hetero atoms as ring members selected from oxygen, nitrogen and sulfur atoms and which heterocyclic group may be saturated or partially unsaturated; wherein each of R¹² and R¹³ taken together may form an aromatic heterocyclic group having five ring members including the nitrogen atom of said amido radical and which aromatic heterocyclic group may further contain one or more additional nitrogen atoms;

wherein each of R⁰ and R² through R¹¹ is independently selected from hydrido, alkyl, hydroxyalkyl, formyl, halo, haloalkyl, cycloalkyl, cycloalkylalkyl, cycloalkylhaloalkyl, cycloalkylcarbonyl, alkoxy, thienylalkyl, phenylalkyl, polycycloalkyl, polycycloalkylalkyl, phenyl, halophenyl, alkylphenyl, alkoxyphenyl, cycloalkenyl, cycloalkenylalkyl, aralkylhaloalkyl, aroyl, aryloxy, aryloxyalkyl, aralkoxy, alkoxyalkyl, alkylcarbonyl, alkylcarbonylalkyl, alkoxyalkyl, alkenyl, cycloalkenyl, alkynyl, cyano, nitro, carboxyl, carboxyalkyl, alkylcarbonyloxy, alkylcarbonyloxyalkyl, alkoxyalkyl, aralkoxyalkyl, aralkylcarbonyloxyalkyl, mercaptocarbonyl, mercaptothiocarbonyl, mercaptoalkyl, alkoxyalkyl, alkylthio, cycloalkylthio, cycloalkylalkylthio, alkylthiocarbonyl, alkylcarbonylthio, alkylthiocarbonyloxy, alkylthiocarbonylthio, alkylthiothiocarbonyl, alkylthiothiocarbonylthio, arylthio, arylthiocarbonyl, arylcarbonylthio, arylthiocarbonyloxy, arylthiocarbonylthio, arylthiothiocarbonyl, arylthiothiocarbonylthio, aralkylthio, aralkylthiocarbonyl, aralkylcarbonylthio, aralkylthiocarbonyloxy, aralkylthiocarbonylthio, alkylthiocarbonyl, aralkylthiocarbonylthio, mercapto, alkylsulfinyl, alkylsulfonyl, aralkylsulfinyl, aralkylsulfonyl, arylsulfinyl, arylsulfonyl, phthalimido, phthalimidoalkyl, heteroaryl, heteroarylalkyl, cycloheteroalkyl, cycloheteroalkylalkyl and cycloheteroalkylcarbonylalkyl wherein each of said heteroaryl- and cyclohetero-containing groups has one or more ring atoms selected from oxygen, sulfur and nitrogen atoms, and wherein each of R⁰ and R² through R¹¹ may be further independently selected from amino and amido radicals of the formula



wherein X is oxygen atom or sulfur atom;

wherein each n is a number independently selected from zero to six, inclusive;

5 wherein each of R¹⁴ through R²⁶ is independently selected from hydrido, alkyl, cycloalkyl, cyano, amino, monoalkylamino, dialkylamino, hydroxyalkyl, cycloalkylalkyl, alkoxyalkyl, aralkyl and aryl, and wherein R¹⁴ and R¹⁵ taken together, R¹⁶ and R¹⁷ taken together, R¹⁸ and R¹⁹ taken together, R²¹ and R²² taken together and R²³ and R²⁴ taken together may each form a heterocyclic group having five to
10 seven ring members including the nitrogen atom of said amino or amido radical and which heterocyclic group may further contain one or more hetero atoms as ring members selected from oxygen, nitrogen and sulfur atoms and which heterocyclic group may be saturated or partially unsaturated; wherein R¹⁴ and R¹⁵ taken together, R¹⁶ and R¹⁷ taken together, R²¹ and R²² taken together and R²³ and R²⁴ taken together may each form an aromatic heterocyclic group having five ring members including the nitrogen atom of said amino or amido radical and which aromatic heterocyclic group may further
15 contain one or more additional nitrogen atoms;


and wherein each of R⁰ and R³ through R¹¹ may be further independently selected from hydroxy and acidic moieties of the formula

20 -Y_nA

wherein n is a number selected from zero through three, inclusive, and wherein A is an acidic group selected to contain at least one acidic hydrogen atom, and the amide, ester and salt derivatives of said acidic moieties;

25 wherein Y is a spacer group independently selected from one or more of alkyl, cycloalkyl, cycloalkylalkyl, alkenyl, alkynyl, aryl, aralkyl and heteroaryl having one or more ring atoms selected from oxygen, sulfur and nitrogen atoms;

30 and wherein any of the foregoing R⁰ through R²⁶, Y and A groups having a substitutable position may be substituted by one or more groups independently selected from hydroxy, alkyl, alkenyl, alkynyl, aralkyl, hydroxyalkyl, haloalkyl, halo, oxo, alkoxy, aryloxy, aralkoxy, aralkylthio, alkoxyalkyl, cycloalkyl, cycloalkylalkyl, aryl, aroyl, cycloalkenyl, cyano, cyanoamino, nitro, alkylcarbonyloxy, alkoxyalkoxy, alkylcarbonyl, alkoxyalkoxy, alkoxyalkoxy, carboxyl, mercapto, mercaptocarbonyl, alkylthio, arylthio, alkylthiocarbonyl, alkylsulfanyl, alkylsulfonyl, haloalkylsulfanyl, aralkylsulfanyl, aralkylsulfonyl, arylsulfanyl, arylsulfonyl, heteroaryl having one or more ring atoms selected from oxygen, sulfur and
35 nitrogen atoms, and amino and amido radicals of the formula




45 wherein X is oxygen atom or sulfur atom; wherein each of R²⁷ through R³¹ is independently selected from hydrido, alkyl, cycloalkyl, cycloalkylalkyl, aralkyl, aryl, DR³² and



55 wherein D is selected from oxygen atom and sulfur atom and R³² is selected from hydrido, alkyl, cycloalkyl, cycloalkylalkyl, aralkyl and aryl; wherein each of R²⁷, R²⁸, R²⁹, R³⁰, R³¹, R³³ and R³⁴ is independently selected from hydrido, alkyl, cycloalkyl, cyano, hydroxyalkyl, haloalkyl, cycloalkylalkyl, alkoxyalkyl, alkylcarbonyl, alkoxyalkoxy, carboxyl, alkylsulfanyl, alkylsulfonyl, arylsulfanyl, arylsulfonyl, haloalkylsulfanyl, haloalkylsulfonyl, aralkyl and aryl, and wherein each of R²⁷, R²⁸, R²⁹, R³⁰, R³¹, R³³ and


R³⁴ is further independently selected from amino and amido radicals of the formula



wherein X is oxygen atom or sulfur atom;

wherein each of R³⁵, R³⁶, R³⁷, R³⁸, R³⁹ and R⁴⁰ is independently selected from hydrido, alkyl, cycloalkyl, cyano, hydroxyalkyl, cycloalkylalkyl, alkoxyalkyl, haloalkylsulfinyl, haloalkylsulfonyl, aralkyl and aryl, and wherein each of R²⁸ and R²⁹ taken together and each of R³⁰ and R³¹ taken together may form a heterocyclic group having five to seven ring members including the nitrogen atom of said amino or amido radical, which heterocyclic group may further contain one or more hetero atoms as ring members selected from oxygen, nitrogen and sulfur atoms and which heterocyclic group may be saturated or partially unsaturated; wherein each of R²⁸ and R²⁹ taken together and each of R³³ and R³⁴ taken together may form an aromatic heterocyclic group having five ring members including the nitrogen atom of said amino or amido radical and which aromatic heterocyclic group may further contain one or more additional nitrogen atoms; or a tautomer thereof or a pharmaceutically-acceptable salt thereof.

64. The method of Claim 63 wherein m is one; wherein R¹ is selected from alkyl, hydroxyalkyl, haloalkyl, cycloalkyl, cycloalkylalkyl, thienylalkyl, phenylalkyl, polycycloalkyl, polycycloalkylalkyl, phenyl, halophenyl, alkylphenyl, alkoxyphenyl, cycloalkenyl, cycloalkenylalkyl, aroyl, alkoxyalkyl, alkylcarbonyl, cycloalkylcarbonyl, cycloalkylalkylcarbonyl, aralkylcarbonyl, alkoxycarbonyl, alkenyl, cycloalkenyl, alkynyl, cycloalkynyl, alkylthiocarbonyl, arylthiocarbonyl, arylthiothiocarbonyl, aralkylthiocarbonyl, alkylsulfonyl, aralkylsulfonyl, arylsulfonyl, heteroaryl having one or more ring atoms selected from oxygen, sulfur and nitrogen atoms, and amido radicals of the formula




wherein X is oxygen atom or sulfur atom;

wherein each of R¹² and R¹³ is independently selected from hydrido, alkyl, cycloalkyl, cyano, amino, monoalkylamino, dialkylamino, hydroxyalkyl, cycloalkylalkyl, alkoxyalkyl, aralkyl and aryl;

wherein each of R⁰ and R² is independently selected from hydrido, alkyl, hydroxyalkyl, formyl, halo, haloalkyl, cycloalkyl, cycloalkylalkyl, cycloalkylhaloalkyl, cycloalkylcarbonyl, alkoxy, thienylalkyl, phenylalkyl, polycycloalkyl, polycycloalkylalkyl, phenyl, halophenyl, alkylphenyl, alkoxyphenyl, cycloalkenyl, cycloalkenylalkyl, aralkylhaloalkyl, aroyl, aryloxy, aryloxyalkyl, aralkoxy, alkoxyalkyl, alkylcarbonyl, alkoxycarbonyl, alkenyl, cycloalkenyl, alkynyl, cycloalkynyl, cyano, nitro, carboxyl, carboxyalkyl, alkylcarbonyloxy, alkylcarbonyloxyalkyl, alkoxycarbonylalkyl, aralkoxycarbonylalkyl, aralkylcarbonyloxyalkyl, mercaptocarbonyl, mercaptothiocarbonyl, mercaptoalkyl, alkoxycarbonyloxy, alkylthio, cycloalkylthio, cycloalkylalkylthio, alkylthiocarbonyl, alkylcarbonylthio, alkylthiocarbonyloxy, alkylthiocarbonylthio, alkylthiothiocarbonyl, alkylthiothiocarbonylthio, arylthio, arylthiocarbonyl, arylcarbonylthio, arylthiocarbonyloxy, arylthiocarbonylthio, arylthiothiocarbonyl, arylthiothiocarbonylthio, aralkylthio, aralkylthiocarbonyl, aralkylcarbonylthio, aralkylthiothiocarbonyl, aralkylthiothiocarbonylthio, aralkylthiothiocarbonyloxy, aralkylthiothiocarbonylthio, aralkylthiothiocarbonylthio, mercapto, alkylsulfinyl, alkylsulfonyl, aralkylsulfinyl, aralkylsulfonyl, arylsulfinyl, arylsulfonyl, phthalimido, phthalimidoalkyl, heteroaryl, heteroarylalkyl, cycloheteroalkyl, cycloheteroalkylalkyl and cycloheteroalkylcarbonylalkyl wherein each of said heteroaryl- and cycloheteroalkyl-containing groups has one or more hetero ring atoms selected from oxygen, sulfur and nitrogen atoms, and wherein each of R⁰ and R² through R¹¹ may be further independently selected from amino and amido

radicals of the formula




wherein X is selected from oxygen atom or sulfur atom;

wherein each n is a number independently selected from zero to six, inclusive;

wherein each of R¹⁴ through R²⁶ is independently selected from hydrido, alkyl, cycloalkyl, cyano, amino, monoalkylamino, dialkylamino, hydroxyalkyl, cycloalkylalkyl, alkoxyalkyl, aralkyl and aryl;

wherein each of R³ through R¹¹ is independently selected from hydrido, hydroxy, alkyl, hydroxyalkyl, halo, haloalkyl, cycloalkyl, cycloalkylalkyl, alkoxy, aralkyl, aryl, aroyl, aryloxy, aralkoxy, alkoxyalkyl, alkylcarbonyl, alkoxy carbonyl, alkenyl, cycloalkenyl, alkynyl, cycloalkynyl, cyano, nitro, carboxyl, alkylcarbonyloxy, mercaptocarbonyl, mercaptothiocarbonyl, alkoxy carbonyloxy, alkylthio, alkylthiocarbonyl, alkylcarbonylthio, alkylthiocarbonyloxy, alkylthiocarbonylthio, alkylthiothiocarbonyl, arylthio, arylthiocarbonyl, arylcarbonylthio, arylthiocarbonyloxy, arylthiothiocarbonyl, aralkylthio, aralkylthiocarbonyl, aralkylcarbonylthio, aralkylthiocarbonyloxy, aralkylthiocarbonylthio, aralkylthiocarbonyl, aralkylthiocarbonylthio, mercapto, alkylsulfonyl, aralkylsulfonyl and arylsulfonyl, and amino and amido radicals of the formula



wherein X is oxygen atom or sulfur atom;

wherein each of R¹⁴, R¹⁵, R¹⁶, R¹⁷, R¹⁸ and R¹⁹ is independently selected from hydrido, alkyl, cycloalkyl, cyano, hydroxyalkyl, cycloalkylalkyl, alkoxyalkyl, aralkyl and aryl;


and wherein each of R⁰ and R³ through R¹¹ may be further independently selected from acidic moieties of the formula




wherein n is a number selected from zero through three, inclusive; wherein A is an acidic group selected from acids containing one or more atoms selected from oxygen, sulfur, phosphorus and nitrogen atoms, and wherein said acidic group is selected to contain at least one acidic hydrogen atom, and the amide, ester and salt derivatives of said acidic moieties; wherein Y is a spacer group independently selected from one or more of alkyl, cycloalkyl, cycloalkylalkyl, alkenyl, alkynyl, aryl, aralkyl and heteroaryl having one or more ring atoms selected from oxygen, sulfur and nitrogen atoms;

and wherein any of the foregoing R⁰ through R²⁶, Y and A groups having a substitutable position may be substituted by one or more groups independently selected from hydroxy, alkyl, alkenyl, aralkyl, hydroxyalkyl, halo, haloalkyl, oxo, alkoxy, aryloxy, aralkoxy, alkoxyalkyl, alkylcarbonyl, alkoxy carbonyl, carboxyl, cyano, nitro, alkylsulfonyl, haloalkylsulfonyl, aryl, aralkyl, mercaptocarbonyl, alkylthio and

alkylthiocarbonyl, and amino and amido radicals of the formula




wherein X is oxygen atom or sulfur atom; wherein each of R²⁷ through R³¹ is independently selected from hydrido, alkyl, cycloalkyl, cycloalkylalkyl, aralkyl, aryl, and DR³² and



wherein D is selected from oxygen atom and sulfur atom, and R³² is selected from hydrido, alkyl, cycloalkyl, cycloalkylalkyl, aralkyl and aryl; wherein each of R²⁷, R²⁸, R²⁹, R³⁰, R³¹, R³³ and R³⁴ is independently selected from hydrido, alkyl, cycloalkyl, cyano, hydroxyalkyl, haloalkyl, cycloalkylalkyl, alkoxyalkyl, alkanoyl, alkoxycarbonyl, carboxyl, haloalkylsulfinyl, haloalkylsulfonyl, aralkyl and aryl;


or a tautomer thereof or a pharmaceutically-acceptable salt thereof.

65. The method of Claim 64 wherein m is one; wherein R¹ is selected from alkyl, hydroxyalkyl, haloalkyl, cycloalkyl, cycloalkylalkyl, thienylalkyl, phenylalkyl, polycycloalkyl, polycycloalkylalkyl, phenyl, halophenyl, alkylphenyl, alkoxyphenyl, cycloalkenyl, cycloalkenylalkyl, aroyl, alkoxyalkyl, alkylcarbonyl, cycloalkylcarbonyl, cycloalkylalkylcarbonyl, aralkylcarbonyl, alkoxyalkyl, alkenyl, cycloalkenyl, alkynyl, cycloalkynyl, alkylsulfonyl, aralkylsulfonyl and arylsulfonyl, and amido radicals of the formula



wherein each of R¹² and R¹³ is independently selected from hydrido, alkyl, cycloalkyl, cyano, amino, monoalkylamino, dialkylamino, hydroxyalkyl, cycloalkylalkyl, alkoxyalkyl, aralkyl and aryl;

wherein each of R⁰ and R² is independently selected from hydrido, alkyl, hydroxyalkyl, formyl, halo, haloalkyl, cycloalkyl, cycloalkylalkyl, cycloalkylhaloalkyl, cycloalkylcarbonyl, alkoxy, thienylalkyl, phenylalkyl, polycycloalkyl, polycycloalkylalkyl, phenyl, halophenyl, alkylphenyl, alkoxyphenyl, cycloalkenyl, cycloalkenylalkyl, aralkylhaloalkyl, aroyl, aryloxy, aryloxyalkyl, aralkoxy, alkoxyalkyl, alkylcarbonyl, alkoxyalkyl, alkenyl, cycloalkenyl, alkynyl, cycloalkynyl, cyano, nitro, carboxyl, carboxyalkyl, alkylcarbonyloxy, mercaptocarbonyl, alkoxyalkyl, alkylcarbonyloxyalkyl, alkoxyalkyl, aralkoxyalkyl, aralkoxyalkyl, aralkylcarbonyloxyalkyl, alkylthio, cycloalkylthio, cycloalkylalkylthio, arylthio, aralkylthio, aralkylthiocarbonylthio, mercapto, alkylsulfinyl, alkylsulfonyl, aralkylsulfinyl, aralkylsulfonyl, arylsulfinyl, arylsulfonyl, phthalimido, phthalimidoalkyl, heteroaryl, heteroarylalkyl, cycloheteroalkyl, cycloheteroalkylalkyl and cycloheteroalkylcarbonylalkyl wherein each of said heteroaryl- and cycloheteroalkyl-containing groups has one or more hetero ring atoms selected from oxygen, sulfur and nitrogen atoms, and wherein each of R⁰ and R² may be further independently selected from amino and amido radicals of the formula




wherein X is selected from oxygen atom or sulfur atom;

wherein each n is a number independently selected from zero to six, inclusive;

wherein each of R¹⁴ through R¹⁵ is independently selected from hydrido, alkyl, cycloalkyl, cyano, amino, monoalkylamino, dialkylamino, hydroxyalkyl, cycloalkylalkyl, alkoxyalkyl, aralkyl and aryl;

wherein each of R³ through R¹¹ is independently selected from hydrido, hydroxy, alkyl, hydroxyalkyl, halo, haloalkyl, cycloalkyl, alkoxy, aralkyl, aryl, aryloxy, aralkoxy, alkoxyalkyl, alkylcarbonyl, alkoxycarbonyl, alkenyl, cycloalkenyl, alkynyl, cyano, nitro, carboxyl, alkylcarbonyloxy, mercaptocarboxyl, alkoxycarbonyloxy, alkylthio, arylthio, aralkylthio, mercapto, alkylsulfonyl, aralkylsulfonyl and arylsulfonyl, and amino and amido radicals of the formula



wherein each of R¹⁴, R¹⁵, R¹⁶, R¹⁷, R¹⁸ and R¹⁹ is independently selected from hydrido, alkyl, cycloalkyl, cyano, amino, monoalkylamino, dialkylamino, hydroxyalkyl, cycloalkylalkyl, alkoxyalkyl, aralkyl and aryl;


and wherein each of R⁰ and R³ through R¹¹ may be further independently selected from acidic moieties of the formula

$$-Y_{DA}$$


wherein n is a number selected from zero through three, inclusive;

wherein A is selected from carboxylic acid and bioisosteres of carboxylic acid selected from


-OH, -SH,



-S-WH,



wherein each W is independently selected from oxygen atom, sulfur atom and NR^{39} ; wherein each of R^{35} , R^{36} , R^{37} , R^{38} and R^{39} is independently selected from hydrido, alkyl, haloalkyl, haloalkylsulfonyl, haloalkylcarbonyl, cycloalkyl, cycloalkylalkyl, aryl and aralkyl; wherein each of R^{35} , R^{36} , R^{37} and R^{39} may be further independently selected from amino radical of the formula




wherein each of R^{40} and R^{41} is independently selected from hydrido, alkyl, cycloalkyl, hydroxyalkyl, haloalkyl, cycloalkylalkyl, alkoxyalkyl, aralkyl and aryl, and wherein R^{40} and R^{41} taken together may form a heterocyclic group having five to seven ring members including the nitrogen atom of said amino radical, which heterocyclic group may further contain one or more hetero atoms as ring members selected from oxygen, nitrogen and sulfur atoms and which heterocyclic group may be saturated or partially unsaturated; wherein R^{40} and R^{41} taken together may form an aromatic heterocyclic group having five ring members including the nitrogen atom of said amino radical and which aromatic heterocyclic group may further contain one or more hetero atoms as ring atoms selected from oxygen, nitrogen and sulfur atoms; wherein each of R^{36} and R^{37} may be further independently selected from hydroxy, alkoxy, alkylthio, aryloxy, arylthio, aralkylthio and aralkoxy; and the amide, ester and salt derivatives of said acidic groups;


wherein said bioisostere of carboxylic acid may be further selected from heterocyclic acidic groups consisting of heterocyclic rings of four to about nine ring members, which heterocyclic ring contains at least one hetero atom selected from oxygen, sulfur and nitrogen atoms, which heterocyclic ring may be saturated, fully unsaturated or partially unsaturated, and which heterocyclic ring may be attached at a single position selected from R^3 through R^{11} or may be attached at any two adjacent positions selected from R^3 through R^{11} so as to form a fused-ring system with one of the phenyl rings of Formula I; and the amide, ester and salt derivatives of said heterocyclic acidic groups;

wherein Y is a spacer group independently selected from one or more of alkyl, cycloalkyl, cycloalkylalkyl, alkenyl, aryl and aralkyl;

and wherein any of the foregoing R^0 through R^{26} and R^{35} through R^{41} , Y and A groups having a substitutable position may be substituted by one or more groups independently selected from hydroxy, alkyl, alkenyl, aralkyl, hydroxyalkyl, halo, oxo, haloalkyl, alkoxy, aryloxy, aralkoxy, alkoxyalkyl, alkylcarbonyl, alkoxy carbonyl, carboxyl, cyano, nitro, alkylsulfonyl, haloalkylsulfonyl, aryl, aralkyl, mercaptocarbonyl, alkylthio and alkylthiocarbonyl, and amino and amido radicals of the formula



wherein X is selected from oxygen atom and sulfur atom;
wherein R^{27} is selected from hydrido, alkyl, cycloalkyl, cycloalkylalkyl, aralkyl, aryl and DR^{32} and



5

wherein D is selected from oxygen atom and sulfur atom; wherein R³² is selected from hydrido, alkyl, cycloalkyl, cycloalkylalkyl, aralkyl and aryl;

10 wherein each of R²⁷, R²⁸, R²⁹, R³⁰, R³¹, R³³ and R³⁴ is independently selected from hydrido, alkyl, cycloalkyl, cyano, hydroxyalkyl, haloalkyl, cycloalkylalkyl, alkoxyalkyl, alkanoyl, alkoxycarbonyl, carboxyl, haloalkylsulfinyl, haloalkylsulfonyl, aralkyl and aryl;


or a tautomer thereof or a pharmaceutically-acceptable salt thereof.

15

66. The method of Claim 65 wherein m is one; wherein R¹ is selected from alkyl, hydroxyalkyl, haloalkyl, cycloalkyl, cycloalkylalkyl, thienylalkyl, phenylalkyl, polycycloalkyl, polycycloalkylalkyl, phenyl, halophenyl, alkylphenyl, alkoxyphenyl, cycloalkenyl, cycloalkenylalkyl, aroyl, alkoxyalkyl, alkylcarbonyl, cycloalkylcarbonyl, cycloalkylalkylcarbonyl, aralkylcarbonyl, alkenyl, cycloalkenyl, alkynyl, mercaptocar-

20

bonyl, alkylsulfonyl, aralkylsulfonyl, arylsulfonyl and amido radicals of the formula




25

wherein each of R¹² and R¹³ is independently selected from hydrido, alkyl, cycloalkyl, cyano, amino, monoalkylamino, dialkylamino, hydroxyalkyl, cycloalkylalkyl, alkoxyalkyl, aralkyl and aryl;

30

wherein each of R⁰ and R² is independently selected from hydrido, alkyl, hydroxyalkyl, formyl, halo, haloalkyl, cycloalkyl, cycloalkylalkyl, cycloalkylhaloalkyl, cycloalkylcarbonyl, alkoxy, thienylalkyl, phenylalkyl, polycycloalkyl, polycycloalkylalkyl, phenyl, halophenyl, alkylphenyl, alkoxyphenyl, cycloalkenyl, cycloalkenylalkyl, aralkylhaloalkyl, aroyl, aryloxy, aryloxyalkyl, aralkoxy, alkoxyalkyl, alkylcarbonyl, alkoxycarbonyl, alkenyl, cycloalkenyl, alkynyl, cycloalkynyl, cyano, nitro, carboxyl, carboxyalkyl, alkylcarbonyloxy, alkylcarbonyloxyalkyl, alkoxycarbonylalkyl, aralkoxy, alkoxyalkyl, alkylcarbonyl, mercaptocarbonyl, mercaptoalkyl, alkoxy, alkoxyalkyl, alkylthio, cycloalkylthio, cycloalkylalkylthio, arylthio, aralkylthio, mercapto, alkylsulfinyl, alkylsulfonyl, aralkylsulfinyl, aralkylsulfonyl, arylsulfinyl, arylsulfonyl, phthalimido, phthalimidoalkyl, heteroaryl, heteroarylalkyl, cycloheteroalkyl, cycloheteroalkylalkyl and cycloheteroalkylcarbonylalkyl wherein each of said heteroaryl- and cycloheteroalkyl-containing groups has one or more hetero ring atoms selected from oxygen, sulfur and nitrogen atoms, and wherein each of R² through R¹¹ may be further independently selected from amino and amido radicals of the formula

45



55

wherein X is selected from oxygen atom and sulfur atom;

wherein each n is a number independently selected from zero to six, inclusive;

wherein each of R^{14} through R^{26} is independently selected from hydrido, alkyl, cycloalkyl, cyano, amino, monoalkylamino, dialkylamino, hydroxyalkyl, cycloalkylalkyl, alkoxyalkyl, aralkyl and aryl;


wherein each of R^3 through R^{11} is independently selected from hydrido, hydroxy, alkyl, hydroxyalkyl, halo, haloalkyl, cycloalkyl, alkoxy, aralkyl, aryl, aroyl, aryloxy, aralkoxy, alkoxyalkyl, alkylcarbonyl, alkoxycarbonyl, alkenyl, cycloalkenyl, alkynyl, cyano, nitro, carboxyl, alkylthio, aralkylthio and mercapto;

and wherein each of R^0 and R^3 through R^{11} may be further independently selected from acidic moieties of the formula


$-Y_nA$

wherein n is a number selected from zero through three, inclusive; wherein A is selected from carboxylic acid and bioisosteres of carboxylic acid selected from


$-OH$, $-SH$,



$-S-\text{WH}$,



wherein each W is independently selected from oxygen atom, sulfur atom and NR^{39} ; wherein each of R^{35} , R^{38} and R^{39} is independently selected from hydrido, alkyl, haloalkyl, haloalkylsulfonyl, haloalkylcarbonyl, cycloalkyl, cycloalkylalkyl, aryl and aralkyl; wherein each of R^{35} and R^{39} may be further independently selected from amino radical of the formula



wherein each of R^{40} and R^{41} is independently selected from hydrido, alkyl, cycloalkyl, hydroxyalkyl, haloalkyl, cycloalkylalkyl, alkoxyalkyl, aralkyl and aryl, and wherein R^{40} and R^{41} taken together may form a heterocyclic group having five to seven ring members including the nitrogen atom of said amino radical, which heterocyclic group may further contain one or more hetero atoms as ring members selected from oxygen, nitrogen and sulfur atoms, and which heterocyclic group may be saturated or partially unsaturated; wherein R^{40} and R^{41} taken together may form an aromatic heterocyclic group having five ring members including the nitrogen atom of said amino radical and which aromatic heterocyclic group may further contain one or more hetero atoms as ring atoms selected from oxygen, nitrogen and sulfur atoms; and the amide, ester and salt derivatives of said acidic groups; wherein said bioisostere of carboxylic acid may be further selected from heterocyclic acidic groups consisting of heterocyclic rings of four to about nine ring members, which ring contains at least one hetero atom, selected from oxygen, sulfur and nitrogen atoms, which heterocyclic ring may be saturated, fully


unsaturated or partially unsaturated, and which heterocyclic ring may be attached at a single position selected from R³ through R¹¹ or may be attached at any two adjacent positions selected from R³ through R¹¹ so as to form a fused-ring system with one of the phenyl rings of Formula I; and the amide, ester and salt derivatives of said heterocyclic acidic groups;

wherein Y is a spacer group independently selected from one or more of alkyl, cycloalkyl, cycloalkylalkyl, alkenyl, aryl and aralkyl;

wherein each of R⁰ through R²⁶, R³⁵ and R³⁸ through R⁴¹, Y and A independently may be substituted at any substitutable position with one or more groups selected from alkyl, hydroxy, halo, oxo, haloalkyl, alkoxycarbonyl, cyano, nitro, alkylsulfonyl, haloalkylsulfonyl, aryl, aralkyl, alkoxy, aryloxy and aralkoxy;


or a tautomer thereof or a pharmaceutically-acceptable salt thereof.

67. The method of Claim 66 wherein m is one; wherein R¹ is selected from alkyl, hydroxyalkyl, haloalkyl, cycloalkyl, cycloalkylalkyl, thienylalkyl, phenylalkyl, polycycloalkyl, polycycloalkylalkyl, phenyl, halophenyl, alkylphenyl, alkoxyphenyl, cycloalkenyl, cycloalkenylalkyl, aroyl, alkoxyalkyl, alkylcarbonyl, cycloalkylcarbonyl, cycloalkylalkylcarbonyl, aralkylcarbonyl, alkenyl, alkynyl, alkylsulfonyl, aralkylsulfonyl, arylsulfonyl and amido radicals of the formula



wherein each of R¹² and R¹³ is independently selected from hydrido, alkyl, cycloalkyl, cyano, amino, hydroxyalkyl, alkoxyalkyl, phenalkyl and phenyl;

wherein each of R⁰ and R² is independently selected from hydrido, alkyl, hydroxyalkyl, formyl, halo, haloalkyl, cycloalkyl, cycloalkylalkyl, cycloalkylhaloalkyl, cycloalkylcarbonyl, alkoxy, thienylalkyl, phenylalkyl, polycycloalkyl, polycycloalkylalkyl, phenyl, halophenyl, alkylphenyl, alkoxyphenyl, cycloalkenyl, cycloalkenylalkyl, aralkylhaloalkyl, benzoyl, phenoxy, phenoxyalkyl, phenalkyloxy, phenylthio, phenalkylthio, aralkoxy, alkoxyalkyl, alkylcarbonyl, alkoxycarbonyl, alkenyl, cycloalkenyl, alkynyl, cyano, nitro, carboxyl, carboxyalkyl, alkylcarbonyloxy, alkylcarbonyloxyalkyl, alkoxycarbonylalkyl, aralkoxycarbonylalkyl, aralkylcarbonyloxyalkyl, mercaptocarbonyl, mercaptoalkyl, alkoxycarbonyloxy, alkylthio, cycloalkylthio, cycloalkylalkylthio, phthalimido, phthalimidoalkyl, heteroaryl, heteroarylalkyl, cycloheteroalkyl, cycloheteroalkylalkyl and cycloheteroalkylcarbonylalkyl wherein each of said heteroaryl- and cycloheteroalkyl-containing groups has one or more hetero ring atoms selected from oxygen, sulfur and nitrogen atoms, and wherein each of R⁰ and R² through R¹¹ may be further independently selected from amino and amido radicals of the formula



wherein X is selected from oxygen atom and sulfur atom;

wherein each n is a number independently selected from zero to six, inclusive;

wherein each of R¹⁴ through R²⁶ is independently selected from hydrido, alkyl, cycloalkyl, cyano, amino, hydroxyalkyl, alkoxyalkyl, phenalkyl and phenyl;


wherein each of R³ through R¹¹ is independently selected from hydrido, hydroxy, alkyl, hydroxyalkyl, halo, haloalkyl, cycloalkyl, alkoxy, phenalkyl, phenyl, benzoyl, phenoxy, phenalkyloxy, alkoxyalkyl, alkylcarbonyl, alkoxycarbonyl, alkenyl, cyano, nitro, carboxyl, alkylthio and mercapto;

and wherein each of R⁰ and R³ through R¹¹ may be further independently selected from acidic moieties of the formula


-Y_nA

wherein n is a number selected from zero through two, inclusive; wherein A is selected from carboxylic acid and bioisosteres of carboxylic acid selected from


-OH, -SH,



-S-WH,



wherein each W is independently selected from oxygen atom, sulfur atom and NR³⁹; wherein each of R³⁵, R³⁸ and R³⁹ is independently selected from hydrido, alkyl, haloalkyl, haloalkylsulfonyl, haloalkylcarbonyl, cycloalkyl, phenyl and benzyl; wherein each of R³⁵ and R³⁹ may be further independently selected from amino radical of the formula



wherein each of R⁴⁰ and R⁴¹ is independently selected from hydrido, alkyl, cycloalkyl, hydroxyalkyl, haloalkyl, alkoxyalkyl, benzyl and phenyl; and the amide, ester and salt derivatives of said acidic groups;

wherein said bioisostere of carboxylic acid may be further selected from heterocyclic acidic groups consisting of heterocyclic rings of four to about nine ring members, which ring contains at least one hetero atom, selected from oxygen, sulfur and nitrogen atoms, which heterocyclic ring may be saturated, fully unsaturated or partially unsaturated, and which heterocyclic ring may be attached at a single position selected from R³ through R¹¹ or may be attached at any two adjacent positions selected from R³ through R¹¹ so as to form a fused-ring system with one of the phenyl rings of Formula I; and the amide, ester and salt derivatives of said heterocyclic acidic groups;

wherein Y is a spacer group independently selected from one or more of alkyl, cycloalkyl, cycloalkylal-


kyl, alkenyl, phenyl, phenalkyl and aralkyl;

wherein each of R^0 through R^{26} , R^{35} and R^{38} through R^{41} , Y and A and independently may be substituted at any substitutable position with one or more groups selected from alkyl, cycloalkyl, cycloalkylalkyl, hydroxy, halo, oxo, haloalkyl, alkoxy, carbonyl, cyano, nitro, alkylsulfonyl, haloalkylsulfonyl, aryl, aralkyl, alkoxy, aryloxy and aralkoxy;

or a tautomer thereof or a pharmaceutically-acceptable salt thereof.

68. The method of Claim 67 wherein m is one; wherein R^1 is selected from alkyl, hydroxyalkyl, haloalkyl, cycloalkyl, cycloalkylalkyl, thienylalkyl, phenylalkyl, polycycloalkyl, polycycloalkylalkyl, phenyl, halophenyl, alkylphenyl, alkoxyphenyl, cycloalkenyl, cycloalkenylalkyl, benzoyl, alkoxyalkyl, alkylcarbonyl, alkoxyalkyl, cycloalkylcarbonyl, cycloalkylalkylcarbonyl, aralkylcarbonyl, alkenyl and alkynyl;

where each of R^0 and R^2 is independently selected from hydrido, alkyl, aminoalkyl, hydroxyalkyl, formyl, halo, haloalkyl, cycloalkyl, cycloalkylalkyl, cycloalkylhaloalkyl, cycloalkylcarbonyl, alkoxy, thienylalkyl, phenylalkyl, polycycloalkyl, polycycloalkylalkyl, phenyl, halophenyl, alkylphenyl, alkoxyphenyl, cycloalkenyl, cycloalkenylalkyl, aralkylhaloalkyl, benzoyl, phenoxy, phenoxyalkyl, phenalkyloxy, phenylthio, phenalkylthio, aralkoxy, alkoxyalkyl, acetyl, alkoxyalkyl, alkenyl, cycloalkenyl, alkynyl, cyano, nitro, carboxyl, carboxyalkyl, alkylcarbonyl, alkylcarbonyloxy, mercaptoalkyl, mercaptocarbonyl, alkoxyalkyl, alkylcarbonyloxyalkyl, alkoxyalkyl, aralkoxyalkyl, aralkylcarbonyloxyalkyl, phthalimido, phthalimidoalkyl, imidazoalkyl, tetrazole, tetrazolealkyl, alkylthio, cycloalkylthio, cycloalkylalkylthio, and amino and amido radicals of the formula




wherein X is selected from oxygen atom and sulfur atom;

wherein each n is a number independently selected from zero to six, inclusive;

wherein each of R^{14} through R^{26} is independently selected from hydrido, alkyl, cycloalkyl, cyano, amino, hydroxyalkyl, alkoxyalkyl, phenalkyl and phenyl;

wherein each of R^3 through R^{11} is independently selected from hydrido, hydroxy, alkyl, hydroxyalkyl, halo, haloalkyl, alkoxy, phenyl, benzoyl, phenoxy, alkoxyalkyl, acetyl, alkoxyalkyl, alkenyl, cyano, nitro, carboxyl, alkylthio and mercapto;

and wherein each of R^0 and R^3 through R^{11} may be further independently selected from acidic moieties consisting of CO_2H , CO_2CH_3 , SH , CH_2SH , $\text{C}_2\text{H}_4\text{SH}$, PO_3H_2 , NHSO_2CF_3 , $\text{NHSO}_2\text{C}_6\text{F}_5$, SO_3H , CONHNH_2 , $\text{CONHNHSO}_2\text{CF}_3$, CONHOCH_3 , $\text{CONHOC}_2\text{H}_5$, CONHCF_3 , OH , CH_2OH , $\text{C}_2\text{H}_4\text{OH}$, OPO_3H_2 , OSO_3H ,



35

40

45

50

55

5


10

15

20

25

30



and the esters, amides and salts of said acidic moieties; or a tautomer thereof or a pharmaceutically-acceptable salt thereof.

69. The method of Claim 68 wherein m is one; wherein R¹ is selected from methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, n-pentyl, isopentyl, neopentyl, phenyl, benzyl, phenethyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cyclopropylmethyl, cyclopropylethyl, cyclobutylmethyl, cyclobutylethyl, cyclopentylmethyl, cyclopentylethyl, cyclohexylmethyl, cyclohexylethyl, cyclohexanoyl, methylbutyl, ethylbutyl, dimethylbutyl, thienylmethyl, thienylethyl, thienylpropyl, cyclopentenylmethyl, cyclopentenylethyl, cyclopentenylpropyl, cyclohexenyl, cyclohexenylmethyl, cyclohexenylethyl, adamantyl, adamantylmethyl, adamantylethyl, phenyl, chlorophenyl, dichlorophenyl, fluorophenyl, difluorophenyl, methoxyphenyl, ethoxyphenyl, methylphenyl, ethylphenyl, propylphenyl, isopropylphenyl, dimethylphenyl, diethylphenyl, benzyl, 1-oxo-2-cyclohexylethyl, benzoyl, 1-oxo-2-phenethyl, 1-oxoethyl, 1-oxopropyl, 1-oxobutyl, 1-oxopentyl, 2-butenyl, 3-butenyl, 2-butyryl, 3-butyryl and 2-hydroxybutyl; wherein R⁰ is selected from hydrido, C₄H₉(n), CH₃CH₂CH=CH, C₃H₇(n), SC₃H₇,

45




50



55


C₂H₅, C₅H₁₁(n), C₆H₁₃(n), SC₄H₉,



5 $\text{CH}_3\text{CH}=\text{CH}$, $\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}=\text{CH}$ -, amino, aminomethyl, aminoethyl, aminopropyl, CH_2OH , $\text{CH}_2\text{OCOCH}_3$, CH_2Cl , CH_2OCH_3 , $\text{CH}_2\text{OCH}(\text{CH}_3)_2$, CHO ,


$\text{CH}_2\text{CO}_2\text{H}$, $\text{CH}(\text{CH}_3)\text{CO}_2\text{H}$, NO_2 , Cl ,

10




15

20



$-\text{CO}_2\text{CH}_3$, $-\text{CONH}_2$, $-\text{CONHCH}_3$, $\text{CON}(\text{CH}_3)_2$, $-\text{CH}_2-\text{NHCO}_2\text{C}_2\text{H}_5$,


25



30


$-\text{CH}_2\text{NHCO}_2\text{CH}_3$, $-\text{CH}_2\text{NHCO}_2\text{C}_3\text{H}_7$, $-\text{CH}_2\text{NHCO}_2\text{CH}_2(\text{CH}_3)_2$, $-\text{CH}_2\text{NHCO}_2\text{C}_4\text{H}_9$, CH_2NHCO_2 -adamantyl, $-\text{CH}_2\text{NHCO}_2$ -(1-naphthyl), $-\text{CH}_2\text{NHCONHCH}_3$, $-\text{CH}_2\text{NHCONHC}_2\text{H}_5$, $-\text{CH}_2\text{NHCONHC}_3\text{H}_7$, $-\text{CH}_2\text{NHCONHC}_4\text{H}_9$, $-\text{CH}_2\text{NHCONHCH}(\text{CH}_3)_2$, $-\text{CH}_2\text{NHCONH}$ (1-naphthyl), $-\text{CH}_2\text{NHCONH}$ (1-adamantyl),

35



$-\text{CH}_2\text{CH}_2\text{CH}_2\text{CO}_2\text{H}$, $-\text{CH}_2\text{CH}_2\text{F}$, $-\text{CH}_2\text{OCONHCH}_3$, $-\text{CH}_2\text{OCSNHCH}_3$, $-\text{CH}_2\text{NHCOC}_3\text{H}_7$, $-\text{CH}_2\text{CH}_2\text{CH}_2\text{F}$, $-\text{CH}_2\text{ONO}_2$,


40



45

$-\text{CH}_2\text{SH}$,

50




55

Cl , NO_2 , CF_3 , CH_2OH , Br , F , I , methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, n-pentyl, isopentyl, neopentyl, phenyl, cyclohexyl, cyclohexylmethyl, carboxyl, formyl, 1-oxoethyl, 1-oxopropyl, 1-oxobutyl, 1-oxopentyl, dimethoxymethyl, 1,1-dimethoxypropyl, 1,1-dimethoxypentyl, hydroxyalkyl, halo, 1-oxo-2-phenylethyl, 1-oxo-2-cyclohexylethyl, 1,1-difluoro-2-phenylethyl, monofluoromethyl, 1,1-difluoro-2-cyclohexylethyl, 2-cyclohexylethyl, 1,1-difluoro-3-cyclohexylpropyl, 1,1-

dimethoxybutyl, 1,1-difluoroethyl, 1,1-difluoropropyl, 1,1-difluorobutyl, 1,1-difluoropentyl, benzyl, 2-phenylethyl, 1,1-difluoro-3-phenylpropyl, cyclohexylmethyl, cyclohexylethyl, cyclohexanoyl, 1-butenyl, 2-butenyl, 3-butenyl, 1-butyryl, 2-butyryl, 3-butyryl, difluoromethyl, CO₂H, SH, PO₃H₂, SO₃H, CONHNH₂, CONHNHSO₂CF₃, OH,


5

10



wherein each of R⁴² and R⁴³ is independently selected from chloro, cyano, nitro, trifluoromethyl, methoxycarbonyl and trifluoromethylsulfonyl; wherein R² is selected from methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, n-pentyl, isopentyl, neopentyl, phenyl, benzyl, phenethyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cyclopropylmethyl, cyclopropylethyl, cyclobutylmethyl, cyclobutylethyl, cyclopentylmethyl, cyclopentylethyl, cyclohexylmethyl, cyclohexylethyl, methylbutyl, ethylbutyl, dimethylbutyl, thienylmethyl, thienylethyl, thienylpropyl, cyclopentenylmethyl, cyclopentenylethyl, cyclopentenylpropyl, cyclohexenyl, cyclohexenylmethyl, cyclohexenylethyl, adamantyl, adamantylmethyl, adamantylethyl, phenyl, chlorophenyl, dichlorophenyl, fluorophenyl, difluorophenyl, methoxyphenyl, ethoxyphenyl, methylphenyl, ethylphenyl, propylphenyl, isopropylphenyl, dimethylphenyl, diethylphenyl, benzyl, propylthio, butylthio and hydroxyalkyl; wherein each of R³ through R¹¹ is hydrido with the proviso that at least one of R⁵, R⁶, R⁸ and R⁹ is an acidic group selected from CO₂H, SH, PO₃H₂, SO₃H, CONHNH₂, CONHNHSO₂CF₃, OH,

30



35

wherein each of R⁴² and R⁴³ is independently selected from chloro, cyano, nitro, trifluoromethyl, methoxycarbonyl and trifluoromethylsulfonyl;


or a tautomer thereof or a pharmaceutically-acceptable salt thereof.

40


70. The method of Claim 69 wherein m is one; wherein R¹ is selected from methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, 4-methylbutyl, n-pentyl, isopentyl, neopentyl, phenyl, benzyl, phenethyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cyclopropylmethyl, cyclopropylethyl, cyclobutylmethyl, cyclobutylethyl, cyclopentylmethyl, cyclopentylethyl, cyclohexylmethyl, cyclohexylethyl, cyclohexanoyl, methylbutyl, ethylbutyl, dimethylbutyl, thienylmethyl, thienylethyl, thienylpropyl, cyclopentenylmethyl, cyclopentenylethyl, cyclopentenylpropyl, cyclohexenyl, cyclohexenylmethyl, cyclohexenylethyl, adamantyl, adamantylmethyl, adamantylethyl, phenyl, chlorophenyl, dichlorophenyl, fluorophenyl, difluorophenyl, methoxyphenyl, ethoxyphenyl, methylphenyl, ethylphenyl, propylphenyl, isopropylphenyl, dimethylphenyl, diethylphenyl, benzyl, 1-oxo-2-cyclohexylethyl, benzoyl, 1-oxo-2-phenethyl, 1-oxoethyl, 1-oxopropyl, 1-oxobutyl, 1-oxopentyl, 2-butenyl, 3-butenyl, 2-butyryl, 3-butyryl and 2-hydroxybutyl; wherein R⁰ is selected from hydrido, neopentyl, 1-oxoethyl, 1-oxopropyl, 1-oxobutyl, 1-oxopentyl, 1-oxo-2-phenylethyl, 1-oxo-2-cyclohexylethyl, 1,1-difluoro-2-phenylethyl, 1,1-difluoro-2-cyclohexylethyl, 2-cyclohexylethyl, 1,1-difluoro-3-cyclohexylpropyl, fluoro, chloro, monofluoromethyl, difluoromethyl, trifluoromethyl, formyl, carboxyl, dimethoxymethyl, 1,1-dimethoxybutyl, 1,1-difluoroethyl, 1,1-difluoropropyl, 1,1-difluorobutyl, 1,1-difluoropentyl, benzyl, 2-phenylethyl, 1,1-difluoro-3-phenylpropyl, cyclohexylmethyl, cyclohexylethyl, cyclohexanoyl, 1-butenyl, 2-butenyl, 3-butenyl, 1-butyryl, 2-butyryl, 3-butyryl, propylthio, butylthio,

55

CO₂H, SH, PO₃H₂, SO₃H, CONHNH₂, CONHNHSO₂CF₃, OH,



wherein each of R⁴² and R⁴³ is independently selected from chloro, cyano, nitro, trifluoromethyl, methoxycarbonyl and trifluoromethylsulfonyl; wherein R² is selected from methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, n-pentyl, isopentyl, neopentyl, phenyl, benzyl, phenethyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cyclopropylmethyl, cyclopropylethyl, cyclobutylmethyl, cyclobutylethyl, cyclopentylmethyl, cyclopentylethyl, cyclohexylmethyl, cyclohexylethyl, methylbutyl, ethylbutyl, dimethylbutyl, thienylmethyl, thienylethyl, thienylpropyl, cyclopentenylmethyl, cyclopentenylethyl, cyclopentenylpropyl, cyclohexenyl, cyclohexenylmethyl, cyclohexenylethyl, adamantyl, adamantylmethyl, adamantylethyl, phenyl, chlorophenyl, dichlorophenyl, fluorophenyl, difluorophenyl, methoxyphenyl, ethoxyphenyl, methylphenyl, ethylphenyl, propylphenyl, isopropylphenyl, dimethylphenyl, diethylphenyl, benzyl, propylthio, butylthio and hydroxyalkyl; wherein each of R³ through R¹¹ is hydrido with the proviso that at least one of R⁵, R⁶, R⁸ and R⁹ is an acidic group selected from CO₂H, SH, PO₃H₂, SO₃H, CONHNH₂, CONHNHSO₂CF₃, OH,




wherein each of R⁴² and R⁴³ is independently selected from chloro, cyano, nitro, trifluoromethyl, methoxycarbonyl and trifluoromethylsulfonyl;

or a tautomer thereof or a pharmaceutically-acceptable salt thereof.

71. The method of Claim 70 wherein m is one; wherein R¹ is selected from methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tertbutyl, 4-methylbutyl, n-pentyl, isopentyl, neopentyl, phenyl, benzyl, phenethyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cyclopropylmethyl, cyclopropylethyl, cyclobutylmethyl, cyclobutylethyl, cyclopentylmethyl, cyclopentylethyl, cyclohexylmethyl, cyclohexylethyl, cyclohexanoyl, methylbutyl, ethylbutyl, dimethylbutyl, thienylmethyl, thienylethyl, thienylpropyl, cyclopentenylmethyl, cyclopentenylethyl, cyclopentenylpropyl, cyclohexenyl, cyclohexenylmethyl, cyclohexenylethyl, adamantyl, adamantylmethyl, adamantylethyl, phenyl, chlorophenyl, dichlorophenyl, fluorophenyl, difluorophenyl, methoxyphenyl, ethoxyphenyl, methylphenyl, ethylphenyl, propylphenyl, isopropylphenyl, dimethylphenyl, diethylphenyl, benzyl, 1-oxo-2-cyclohexylethyl, benzoyl, 1-oxo-2-phenethyl, 1-oxopropyl, 1-oxobutyl, 1-oxopentyl and 2-hydroxybutyl; wherein R⁰ is selected from hydrido, methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, 4-methylbutyl, n-pentyl, 1-oxo-2-phenylethyl, 1-oxo-2-cyclohexylethyl, fluoro, chloro, monofluoromethyl, difluoromethyl, trifluoromethyl, formyl, carboxyl, dimethoxymethyl, 1,1-difluoro-2-phenylethyl, 1,1-difluoro-2-cyclohexylethyl, 2-cyclohexylethyl, 1,1-difluoro-3-cyclohexylpropyl, dimethoxymethyl, 1,1-dimethoxybutyl, 1,1-difluoroethyl, 1,1-difluoropropyl, 1,1-difluorobutyl, 1,1-difluoropentyl, benzyl, 2-phenylethyl, 1,1-difluoro-3-phenylpropyl, cyclohexylmethyl, cyclohexylethyl, cyclohexanoyl, 1-butenyl, 2-butenyl, 3-butenyl, 1-butylnyl, 2-butylnyl, 3-butylnyl, propylthio and butylthio; wherein R² is selected from methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, n-pentyl, isopentyl, neopentyl, phenyl, benzyl, phenethyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cyclopropylmethyl, cyclopropylethyl, cyclobutylmethyl, cyclobutylethyl, cyclopentylmethyl, cyclopentylethyl,


cyclohexylmethyl, cyclohexylethyl, methylbutyl, ethylbutyl, dimethylbutyl, thienylmethyl, thienylethyl, thienylpropyl, cyclopentenylmethyl, cyclopentenylethyl, cyclopentenylpropyl, cyclohexenyl, cyclohexenylmethyl, cyclohexenylethyl, adamantyl, adamantylmethyl, adamantylethyl, phenyl, chlorophenyl, dichlorophenyl, fluorophenyl, difluorophenyl, methoxyphenyl, ethoxyphenyl, methylphenyl, ethylphenyl, propylphenyl, isopropylphenyl, dimethylphenyl, diethylphenyl, benzyl, propylthio, butylthio and hydroxyalkyl; wherein each of R³, R⁴, R⁵, R⁷, R⁸, R¹⁰ and R¹¹ is hydrido; wherein one of R⁵ and R⁹ is hydrido and the other of R⁵ and R⁹ is an acidic group selected from COOH, SH, PO₃H₂, SO₃H, CONHNH₂, CONHNHSO₂CF₃, OH,



wherein each of R⁴² and R⁴³ is independently selected from chloro, cyano, nitro, trifluoromethyl, methoxycarbonyl and trifluoromethylsulfonyl;

or a tautomer thereof or a pharmaceutically-acceptable salt thereof.

72. The method of Claim 71 wherein m is one; wherein R¹ is selected from methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, 4-methylbutyl, n-pentyl, isopentyl, neopentyl, phenyl, benzyl, phenethyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cyclopropylmethyl, cyclopropylethyl, cyclobutylmethyl, cyclobutylethyl, cyclopentylmethyl, cyclopentylethyl, cyclohexylmethyl, cyclohexylethyl, cyclohexanoyl, methylbutyl, ethylbutyl, dimethylbutyl, thienylmethyl, thienylethyl, thienylpropyl, cyclopentenylmethyl, cyclopentenylethyl, cyclopentenylpropyl, cyclohexenyl, cyclohexenylmethyl, cyclohexenylethyl, adamantyl, adamantylmethyl, adamantylethyl, phenyl, chlorophenyl, dichlorophenyl, fluorophenyl, difluorophenyl, methoxyphenyl, ethoxyphenyl, methylphenyl, ethylphenyl, propylphenyl, isopropylphenyl, dimethylphenyl, diethylphenyl, benzyl, 1-oxo-2-phenethyl, 1-oxopropyl, 1-oxobutyl, 1-oxopentyl and 2-hydroxybutyl; wherein R⁰ is selected from hydrido, methyl, fluoro, chloro, monofluoromethyl, difluoromethyl, trifluoromethyl, formyl, carboxyl and dimethoxymethyl; wherein R² is selected from methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, n-pentyl, isopentyl, neopentyl, phenyl, benzyl, phenethyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cyclopropylmethyl, cyclopropylethyl, cyclobutylmethyl, cyclobutylethyl, cyclopentylmethyl, cyclopentylethyl, cyclohexylmethyl, cyclohexylethyl, methylbutyl, ethylbutyl, dimethylbutyl, thienylmethyl, thienylethyl, thienylpropyl, cyclopentenylmethyl, cyclopentenylethyl, cyclopentenylpropyl, cyclohexenyl, cyclohexenylmethyl, cyclohexenylethyl, adamantyl, adamantylmethyl, adamantylethyl, phenyl, chlorophenyl, dichlorophenyl, fluorophenyl, difluorophenyl, methoxyphenyl, ethoxyphenyl, methylphenyl, ethylphenyl, propylphenyl, isopropylphenyl, dimethylphenyl, diethylphenyl, benzyl, propylthio, butylthio and hydroxyalkyl; wherein each of R³, R⁴, R⁵, R⁷, R⁸, R¹⁰ and R¹¹ is hydrido; with the proviso that at least one of R⁵ and R⁹ must be selected from COOH, SH, PO₃H₂, SO₃H, CONHNH₂, CONHNHSO₂CF₃, OH,




wherein each of R⁴² and R⁴³ is independently selected from chloro, cyano, nitro, trifluoromethyl, methoxycarbonyl and trifluoromethylsulfonyl;

or a tautomer thereof or a pharmaceutically-acceptable salt thereof.


73. The method of Claim 72 wherein m is one; wherein R¹ is selected from methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, 4-methylbutyl, n-pentyl, isopentyl, neopentyl, phenyl, benzyl, phenethyl, cyclopropyl, cyclobutyl,

cyclopentyl, cyclohexyl, cyclopropylmethyl, cyclopropylethyl, cyclobutylmethyl, cyclobutylethyl, cyclopentylmethyl, cyclopentylethyl, cyclohexylmethyl, cyclohexylethyl, cyclohexanoyl, methylbutyl, ethylbutyl, dimethylbutyl, thienylmethyl, thienylethyl, thienylpropyl, cyclopentenylmethyl, cyclopentenylethyl, cyclopentenylpropyl, cyclohexenyl, cyclohexenylmethyl, cyclohexenylethyl, adamantyl, adamantylmethyl, adamantylethyl, phenyl, chlorophenyl, dichlorophenyl, fluorophenyl, difluorophenyl, methoxyphenyl, ethoxyphenyl, methylphenyl, ethylphenyl, propylphenyl, isopropylphenyl, dimethylphenyl, diethylphenyl, benzyl, 1-oxo-2-cyclohexylethyl, benzoyl, 1-oxo-2-phenethyl, 1-oxopropyl, 1-oxobutyl, 1-oxopentyl and 2-hydroxybutyl; wherein R^0 is selected from hydrido, methyl, fluoro, chloro, monofluoromethyl, difluoromethyl, trifluoromethyl, formyl, carboxyl and dimethoxymethyl; wherein R^2 is selected from methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, n-pentyl, isopentyl, neopentyl, phenyl, benzyl, phenethyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cyclopropylmethyl, cyclopropylethyl, cyclobutylmethyl, cyclobutylethyl, cyclopentylmethyl, cyclopentylethyl, cyclohexylmethyl, cyclohexylethyl, methylbutyl, ethylbutyl, dimethylbutyl, thienylmethyl, thienylethyl, thienylpropyl, cyclopentenylmethyl, cyclopentenylethyl, cyclopentenylpropyl, cyclohexenyl, cyclohexenylmethyl, cyclohexenylethyl, adamantyl, adamantylmethyl, adamantylethyl, phenyl, chlorophenyl, dichlorophenyl, fluorophenyl, difluorophenyl, methoxyphenyl, ethoxyphenyl, methylphenyl, ethylphenyl, propylphenyl, isopropylphenyl, dimethylphenyl, diethylphenyl, benzyl, propylthio, butylthio and hydroxyalkyl; wherein each of R^3 , R^4 , R^6 , R^7 , R^8 , R^{10} and R^{11} is hydrido; wherein one of R^5 and R^9 is hydrido and the other of R^5 and R^9 is an acidic group selected from CO_2H and




or a tautomer thereof or a pharmaceutically-acceptable salt thereof.

74. The method of Claim 73 wherein said active compound is selected from a family of compounds of Formula II




wherein R^1 is selected from methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, 4-methylbutyl, n-pentyl, isopentyl, neopentyl, phenyl, benzyl, phenethyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cyclopropylmethyl, cyclopropylethyl, cyclobutylmethyl, cyclobutylethyl, cyclopentylmethyl, cyclopentylethyl, cyclohexylmethyl, cyclohexylethyl, cyclohexanoyl, methylbutyl, ethylbutyl, dimethylbutyl, thienylmethyl, thienylethyl, thienylpropyl, cyclopentenylmethyl, cyclopentenylethyl, cyclopentenylpropyl, cyclohexenyl, cyclohexenylmethyl, cyclohexenylethyl, adamantyl, adamantylmethyl, adamantylethyl, phenyl, chlorophenyl, dichlorophenyl, fluorophenyl, difluorophenyl, methoxyphenyl, ethoxyphenyl, methylphenyl, ethylphenyl, propylphenyl, isopropylphenyl, dimethylphenyl, diethylphenyl, benzyl, 1-oxo-2-cyclohexylethyl, benzoyl, 1-oxo-2-phenethyl, 1-oxopropyl, 1-oxobutyl, 1-oxopentyl and 2-hydroxybutyl; wherein R^0 is selected from hydrido, methyl, fluoro, chloro, monofluoromethyl, difluoromethyl, trifluoromethyl, formyl, carboxyl and dimethoxymethyl; wherein R^2 is selected from methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, n-pentyl, isopentyl, neopentyl, phenyl, benzyl, phenethyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cyclopropylmethyl, cyclopropylethyl, cyclobutylmethyl, cyclobutylethyl, cyclopentylmethyl, cyclopentylethyl, cyclohexylmethyl, cyclohexylethyl, methylbutyl, ethylbutyl, dimethylbutyl, thienylmethyl, thienylethyl, thienylpropyl, cyclopentenylmethyl, cyclopentenylethyl, cyclopentenylpropyl, cyclohexenyl, cyclohexenylmethyl, cyclohexenylethyl, adamantyl, adamantylmethyl, adamantylethyl, phenyl, chlorophenyl,

dichlorophenyl, fluorophenyl, difluorophenyl, methoxyphenyl, ethoxyphenyl, methylphenyl, ethylphenyl, propylphenyl, isopropylphenyl, dimethylphenyl, diethylphenyl and benzyl; wherein R⁵ is an acidic group selected from CO₂H and



or a tautomer thereof or a pharmaceutically-acceptable salt thereof.

75. The method of Claim 74 wherein R¹ is selected from methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, 4-methylbutyl, n-pentyl, isopentyl, neopentyl, phenyl, benzyl, phenethyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cyclopropylmethyl, cyclopropylethyl, cyclobutylmethyl, cyclobutylethyl, cyclopentylmethyl, cyclopentylethyl, cyclohexylmethyl, cyclohexylethyl, cyclohexanoyl, methylbutyl, ethylbutyl, dimethylbutyl, thienylmethyl, thienylethyl, thienylpropyl, cyclopentenylmethyl, cyclopentenylethyl, cyclopentenylpropyl, cyclohexenyl, cyclohexenylmethyl, cyclohexenylethyl, adamantyl, adamantylmethyl, adamantylethyl, phenyl, chlorophenyl, dichlorophenyl, fluorophenyl, difluorophenyl, methoxyphenyl, ethoxyphenyl, methylphenyl, ethylphenyl, propylphenyl, isopropylphenyl, dimethylphenyl, diethylphenyl, benzyl, 1-oxo-2-cyclohexylethyl, benzoyl, 1-oxo-2-phenethyl, 1-oxopropyl, 1-oxobutyl, 1-oxopentyl and 2-hydroxybutyl; wherein R⁰ is hydrido; wherein R² is selected from methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, n-pentyl, isopentyl, neopentyl, phenyl, benzyl, phenethyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cyclopropylmethyl, cyclopropylethyl, cyclobutylmethyl, cyclobutylethyl, cyclopentylmethyl, cyclopentylethyl, cyclohexylmethyl, cyclohexylethyl, methylbutyl, ethylbutyl, dimethylbutyl, thienylmethyl, thienylethyl, thienylpropyl, cyclopentenylmethyl, cyclopentenylethyl, cyclopentenylpropyl, cyclohexenyl, cyclohexenylmethyl, cyclohexenylethyl, adamantyl, adamantylmethyl, adamantylethyl, phenyl, chlorophenyl, dichlorophenyl, fluorophenyl, difluorophenyl, methoxyphenyl, ethoxyphenyl, methylphenyl, ethylphenyl, propylphenyl, isopropylphenyl, dimethylphenyl, diethylphenyl and benzyl; wherein R⁵ is an acidic group selected from CO₂H and



or a tautomer thereof or a pharmaceutically-acceptable salt thereof.

76. The method of Claim 75 wherein said active compound is selected from compounds and their pharmaceutically-acceptable salts of the group of compounds consisting of
- 1-propyl-4-butyl-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
 - 1,4-dibutyl-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
 - 1-pentyl-4-butyl-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
 - 1-isopentyl-4-butyl-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
 - 1-cyclohexylmethyl-4-butyl-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
 - 1-phenylmethyl-4-butyl-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
 - 1-(2-cyclohexylethyl)-4-butyl-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
 - 1-(2-phenylethyl)-4-butyl-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
 - 1-propyl-4-butyl-5-methyl-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-

2-one;

1,4-dibutyl-5-methyl-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
1-pentyl-4-butyl-5-methyl-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-

2-one;

1-isopentyl-4-butyl-5-methyl-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;

1-cyclohexylmethyl-4-butyl-5-methyl-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;

1-phenylmethyl-4-butyl-5-methyl-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;

1-(2-cyclohexylethyl)-4-butyl-5-methyl-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;

1-(2-phenylethyl)-4-butyl-5-methyl-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;

1-propyl-4-butyl-5-chloro-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;
1,4-dibutyl-5-chloro-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;

1-pentyl-4-butyl-5-chloro-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;

1-isopentyl-4-butyl-5-chloro-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;


1-cyclohexylmethyl-4-butyl-5-chloro-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;

1-phenylmethyl-4-butyl-5-chloro-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one;

1-(2-cyclohexylethyl)-4-butyl-5-chloro-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one; and

1-(2-phenylethyl)-4-butyl-5-chloro-1,3-dihydro-3-[[5-[2-(1H-tetrazol-5-yl)phenyl]-2-pyridinyl]methyl]-2H-imidazol-2-one.


77. The method of Claim 73 wherein said active compound is selected from a family of compounds of Formula III



(III)


wherein R¹ is selected from methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, 4-methylbutyl, n-pentyl, isopentyl, neopentyl, phenyl, benzyl, phenethyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cyclopropylmethyl, cyclopropylethyl, cyclobutylmethyl, cyclobutylethyl, cyclopentylmethyl, cyclopentylethyl, cyclohexylmethyl, cyclohexylethyl, cyclohexanoyl, methylbutyl, ethylbutyl, dimethylbutyl, thienylmethyl, thienylethyl, thienylpropyl, cyclopentenylmethyl, cyclopentenylethyl, cyclopentenylpropyl, cyclohexenyl, cyclohexenylmethyl, cyclohexenylethyl, adamantyl, adamantylmethyl, adamantylethyl, phenyl, chlorophenyl, dichlorophenyl, fluorophenyl, difluorophenyl, methoxyphenyl, ethoxyphenyl, methylphenyl, ethylphenyl, propylphenyl, isopropylphenyl, dimethylphenyl, diethylphenyl, benzyl, 1-oxo-2-cyclohexylethyl, benzoyl, 1-oxo-2-phenethyl, 1-oxopropyl, 1-oxobutyl, 1-oxopentyl and 2-hydroxybutyl; wherein R⁰ is selected from hydrido, methyl, fluoro, chloro, monofluoromethyl, difluoromethyl, trifluoromethyl, formyl, carboxyl and dimethoxymethyl; wherein R² is selected from methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, n-pentyl, isopentyl, neopentyl, phenyl, benzyl, phenethyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cyclopropylmethyl, cyclopropylethyl, cyclobutylmethyl, cyclobutylethyl, cyclopentylmethyl, cyclopentylethyl, cyclohexylmethyl, cyclohexylethyl, methylbutyl, ethylbutyl, dimethylbutyl, thienylmethyl, thienylethyl, thienylpropyl, cyclopentenylmethyl, cyclopentenylethyl, cyclopentenylpropyl, cyclohexenyl, cyclohexenylmethyl, cyclohexenylethyl, adamantyl, adamantylmethyl, adamantylethyl, phenyl, chlorophenyl,

dichlorophenyl, fluorophenyl, difluorophenyl, methoxyphenyl, ethoxyphenyl, methylphenyl, ethylphenyl, propylphenyl, isopropylphenyl, dimethylphenyl, diethylphenyl and benzyl; wherein R⁵ is an acidic group selected from CO₂H and



or a tautomer thereof or a pharmaceutically-acceptable salt thereof.

78. The method of Claim 77 wherein R¹ is selected from methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, 4-methylbutyl, n-pentyl, isopentyl, neopentyl, phenyl, benzyl, phenethyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cyclopropylmethyl, cyclopropylethyl, cyclobutylmethyl, cyclobutylethyl, cyclopentylmethyl, cyclopentylethyl, cyclohexylmethyl, cyclohexylethyl, cyclohexanoyl, methylbutyl, ethylbutyl, dimethylbutyl, thienylmethyl, thienylethyl, thienylpropyl, cyclopentenylmethyl, cyclopentenylethyl, cyclopentenylpropyl, cyclohexenyl, cyclohexenylmethyl, cyclohexenylethyl, adamantyl, adamantylmethyl, adamantylethyl, phenyl, chlorophenyl, dichlorophenyl, fluorophenyl, difluorophenyl, methoxyphenyl, ethoxyphenyl, methylphenyl, ethylphenyl, propylphenyl, isopropylphenyl, dimethylphenyl, diethylphenyl, benzyl, 1-oxo-2-cyclohexylethyl, benzoyl, 1-oxo-2-phenethyl, 1-oxopropyl, 1-oxobutyl, 1-oxopentyl and 2-hydroxybutyl; wherein R⁰ is hydrido; wherein R² is selected from methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, n-pentyl, isopentyl, neopentyl, phenyl, benzyl, phenethyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cyclopropylmethyl, cyclopropylethyl, cyclobutylmethyl, cyclobutylethyl, cyclopentylmethyl, cyclopentylethyl, cyclohexylmethyl, cyclohexylethyl, methylbutyl, ethylbutyl, dimethylbutyl, thienylmethyl, thienylethyl, thienylpropyl, cyclopentenylmethyl, cyclopentenylethyl, cyclopentenylpropyl, cyclohexenyl, cyclohexenylmethyl, cyclohexenylethyl, adamantyl, adamantylmethyl, adamantylethyl, phenyl, chlorophenyl, dichlorophenyl, fluorophenyl, difluorophenyl, methoxyphenyl, ethoxyphenyl, methylphenyl, ethylphenyl, propylphenyl, isopropylphenyl, dimethylphenyl, diethylphenyl and benzyl; wherein R⁵ is an acidic group selected from CO₂H and



or a tautomer thereof or a pharmaceutically-acceptable salt thereof.

79. The method of Claim 78 wherein said active compound is selected from compounds and their pharmaceutically-acceptable salts of the group of compounds consisting of
- 1,4-dibutyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
 - 1,4-dipropyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
 - 1-propyl-4-isobutyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
 - 1-butyl-4-isobutyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
 - 1-butyl-4-isopentyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
 - 1-butyl-4-pentyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;

- 1-butyl-4-cyclohexyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
- 1-butyl-4-cyclopentyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
- 5 1-butyl-4-(2-cyclopropylethyl)-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
- 1-methyl-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
- 10 1-ethyl-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
- 1-propyl-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
- 15 1-isobutyl-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
- 1-isopentyl-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
- 1-pentyl-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
- 20 1-isopropyl-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
- 1-tertbutyl-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
- 25 1-neopentyl-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
- 1-(3,3-dimethylbutyl)-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
- 30 1-(2-ethylbutyl)-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
- 1-cyclohexyl-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
- 35 1-(2-cyclopropylethyl)-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
- 1-cyclopentyl-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
- 40 1-cyclopentylmethyl-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
- 1-[2-(2-thienyl)ethyl]-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
- 45 1-(2-phenylethyl)-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
- 50 1-(2-cyclopentylethyl)-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
- 1-[2-(cyclopenten-1-yl)ethyl]-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;
- 55 1-(2-adamantyl)-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;

1-(1-adamantyl)-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;

1-phenyl-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;

1-(2-chlorophenyl)-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;

1-(2-methylphenyl)-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;

1-(2-methoxyphenyl)-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;


1-(2-isopropylphenyl)-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;

1-(2,6-difluorophenyl)-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one;


1-(2,6-dichlorophenyl)-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one; and

1-(2,6-dimethylphenyl)-4-butyl-1,3-dihydro-3-[[6-[2-(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one.

80. The method of Claim 73 wherein said active compound is selected from a family of compounds of Formula IV




wherein R¹ is selected from methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, 4-methylbutyl, n-pentyl, isopentyl, neopentyl, phenyl, benzyl, phenethyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cyclopropylmethyl, cyclopropylethyl, cyclobutylmethyl, cyclobutylethyl, cyclopentylmethyl, cyclopentylethyl, cyclohexylmethyl, cyclohexylethyl, cyclohexanoyl, methylbutyl, ethylbutyl, dimethylbutyl, thienylmethyl, thienylethyl, thienylpropyl, cyclopentenylmethyl, cyclopentenylethyl, cyclopentenylpropyl, cyclohexenyl, cyclohexenylmethyl, cyclohexenylethyl, adamantyl, adamantylmethyl, adamantylethyl, phenyl, chlorophenyl, dichlorophenyl, fluorophenyl, difluorophenyl, methoxyphenyl, ethoxyphenyl, methylphenyl, ethylphenyl, propylphenyl, isopropylphenyl, dimethylphenyl, diethylphenyl, benzyl, 1-oxo-2-cyclohexylethyl, benzoyl, 1-oxo-2-phenethyl, 1-oxopropyl, 1-oxobutyl, 1-oxopentyl and 2-hydroxybutyl; wherein R⁰ is selected from hydrido, methyl, fluoro, chloro, monofluoromethyl, difluoromethyl, trifluoromethyl, formyl, carboxyl and dimethoxymethyl; wherein R² is selected from methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, n-pentyl, isopentyl, neopentyl, phenyl, benzyl, phenethyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cyclopropylmethyl, cyclopropylethyl, cyclobutylmethyl, cyclobutylethyl, cyclopentylmethyl, cyclopentylethyl, cyclohexylmethyl, cyclohexylethyl, methylbutyl, ethylbutyl, dimethylbutyl, thienylmethyl, thienylethyl, thienylpropyl, cyclopentenylmethyl, cyclopentenylethyl, cyclopentenylpropyl, cyclohexenyl, cyclohexenylmethyl, cyclohexenylethyl, adamantyl, adamantylmethyl, adamantylethyl, phenyl, chlorophenyl, dichlorophenyl, fluorophenyl, difluorophenyl, methoxyphenyl, ethoxyphenyl, methylphenyl, ethylphenyl, propylphenyl, isopropylphenyl, dimethylphenyl, diethylphenyl and benzyl; wherein R⁵ is an acidic group selected from CO₂H and



or a tautomer thereof or a pharmaceutically-acceptable salt thereof.

81. The method of Claim 80 wherein R¹ is selected from methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, 4-methylbutyl, n-pentyl, isopentyl, neopentyl, phenyl, benzyl, phenethyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cyclopropylmethyl, cyclopropylethyl, cyclobutylmethyl, cyclobutylethyl, cyclopentylmethyl, cyclopentylethyl, cyclohexylmethyl, cyclohexylethyl, cyclohexanoyl, methylbutyl, ethylbutyl, dimethylbutyl, thienylmethyl, thienylethyl, thienylpropyl, cyclopentenylmethyl, cyclopentenylethyl, cyclopentenylpropyl, cyclohexenyl, cyclohexenylmethyl, cyclohexenylethyl, adamantyl, adamantylmethyl, adamantylethyl, phenyl, chlorophenyl, dichlorophenyl, fluorophenyl, difluorophenyl, methoxyphenyl, ethoxyphenyl, methylphenyl, ethylphenyl, propylphenyl, isopropylphenyl, dimethylphenyl, diethylphenyl, benzyl, 1-oxo-2-cyclohexylethyl, benzoyl, 1-oxo-2-phenethyl, 1-oxopropyl, 1-oxobutyl, 1-oxopentyl and 2-hydroxybutyl; wherein R⁰ is hydrido; wherein R² is selected from methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, n-pentyl, isopentyl, neopentyl, phenyl, benzyl, phenethyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cyclopropylmethyl, cyclopropylethyl, cyclobutylmethyl, cyclobutylethyl, cyclopentylmethyl, cyclopentylethyl, cyclohexylmethyl, cyclohexylethyl, methylbutyl, ethylbutyl, dimethylbutyl, thienylmethyl, thienylethyl, thienylpropyl, cyclopentenylmethyl, cyclopentenylethyl, cyclopentenylpropyl, cyclohexenyl, cyclohexenylmethyl, cyclohexenylethyl, adamantyl, adamantylmethyl, adamantylethyl, phenyl, chlorophenyl, dichlorophenyl, fluorophenyl, difluorophenyl, methoxyphenyl, ethoxyphenyl, methylphenyl, ethylphenyl, propylphenyl, isopropylphenyl, dimethylphenyl, diethylphenyl and benzyl; wherein R⁵ is an acidic group selected from CO₂H and



or a tautomer thereof or a pharmaceutically-acceptable salt thereof.

82. The method of Claim 81 wherein said active compound is selected from compounds and their pharmaceutically-acceptable salts of the group of compounds consisting of
- 1-propyl-4-butyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 - 1,4-dibutyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 - 1-pentyl-4-butyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 - 1-isopentyl-4-butyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 - 1-cyclohexylmethyl-4-butyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 - 1-phenylmethyl-4-butyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 - 1-(2-cyclohexylethyl)-4-butyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 - 1-(2-phenylethyl)-4-butyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 - 1-propyl-4-butyl-5-methyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 - 1,4-dibutyl-5-methyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 - 1-pentyl-4-butyl-5-methyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

2-one;

1-isopentyl-4-butyl-5-methyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

1-cyclohexylmethyl-4-butyl-5-methyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

1-phenylmethyl-4-butyl-5-methyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

1-(2-cyclohexylethyl)-4-butyl-5-methyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

1-(2-phenylethyl)-4-butyl-5-methyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

1-propyl-4-butyl-5-chloro-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

1,4-dibutyl-5-chloro-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

1-pentyl-4-butyl-5-chloro-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

1-isopentyl-4-butyl-5-chloro-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;


1-cyclohexylmethyl-4-butyl-5-chloro-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

1-phenylmethyl-4-butyl-5-chloro-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;


1-(2-cyclohexylethyl)-4-butyl-5-chloro-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one; and

1-(2-phenylethyl)-4-butyl-5-chloro-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-2-pyridinyl]phenyl]methyl]-2H-imidazol-2-one.

83. The method of Claim 73 wherein said active compound is selected from a family of compounds of Formula V




wherein R¹ is selected from methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, 4-methylbutyl, n-pentyl, isopentyl, neopentyl, phenyl, benzyl, phenethyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cyclopropylmethyl, cyclopropylethyl, cyclobutylmethyl, cyclobutylethyl, cyclopentylmethyl, cyclopentylethyl, cyclohexylmethyl, cyclohexylethyl, cyclohexanoyl, methylbutyl, ethylbutyl, dimethylbutyl, thienylmethyl, thienylethyl, thienylpropyl, cyclopentenylmethyl, cyclopentenylethyl, cyclopentenylpropyl, cyclohexenyl, cyclohexenylmethyl, cyclohexenylethyl, adamantyl, adamantylmethyl, adamantylethyl, phenyl, chlorophenyl, dichlorophenyl, fluorophenyl, difluorophenyl, methoxyphenyl, ethoxyphenyl, methylphenyl, ethylphenyl, propylphenyl, isopropylphenyl, dimethylphenyl, diethylphenyl, benzyl, 1-oxo-2-cyclohexylethyl, benzoyl, 1-oxo-2-phenethyl, 1-oxopropyl, 1-oxobutyl, 1-oxopentyl and 2-hydroxybutyl; wherein R⁰ is selected from hydrido, methyl, fluoro, chloro, monofluoromethyl, difluoromethyl, trifluoromethyl, formyl, carboxyl and dimethoxymethyl; wherein R² is selected from methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, n-pentyl, isopentyl, neopentyl, phenyl, benzyl, phenethyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cyclopropylmethyl, cyclopropylethyl, cyclobutylmethyl, cyclobutylethyl, cyclopentylmethyl, cyclopentylethyl, cyclohexylmethyl, cyclohexylethyl, methylbutyl, ethylbutyl, dimethylbutyl, thienylmethyl, thienylethyl, thienylpropyl, cyclopentenylmethyl, cyclopentenylethyl, cyclopentenylpropyl, cyclohexenyl, cyclohexenylmethyl, cyclohexenylethyl, adamantyl, adamantylmethyl, adamantylethyl, phenyl, chlorophenyl, dichlorophenyl, fluorophenyl, difluorophenyl, methoxyphenyl, ethoxyphenyl, methylphenyl, ethylphenyl, propylphenyl, isopropylphenyl, dimethylphenyl, diethylphenyl and benzyl; wherein R⁵ is an acidic group selected from CO₂H and



or a tautomer thereof or a pharmaceutically-acceptable salt thereof.

84. The method of Claim 83 wherein R¹ is selected from methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, 4-methylbutyl, n-pentyl, isopentyl, neopentyl, phenyl, benzyl, phenethyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cyclopropylmethyl, cyclopropylethyl, cyclobutylmethyl, cyclobutylethyl, cyclopentylmethyl, cyclopentylethyl, cyclohexylmethyl, cyclohexylethyl, cyclohexanoyl, methylbutyl, ethylbutyl, dimethylbutyl, thienylmethyl, thienylethyl, thienylpropyl, cyclopentenylmethyl, cyclopentenylethyl, cyclopentenylpropyl, cyclohexenyl, cyclohexenylmethyl, cyclohexenylethyl, adamantyl, adamantylmethyl, adamantylethyl, phenyl, chlorophenyl, dichlorophenyl, fluorophenyl, difluorophenyl, methoxyphenyl, ethoxyphenyl, methylphenyl, ethylphenyl, propylphenyl, isopropylphenyl, dimethylphenyl, diethylphenyl, benzyl, 1-oxo-2-cyclohexylethyl, benzoyl, 1-oxo-2-phenethyl, 1-oxopropyl, 1-oxobutyl, 1-oxopentyl and 2-hydroxybutyl; wherein R⁰ is hydrido; wherein R² is selected from methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, n-pentyl, isopentyl, neopentyl, phenyl, benzyl, phenethyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cyclopropylmethyl, cyclopropylethyl, cyclobutylmethyl, cyclobutylethyl, cyclopentylmethyl, cyclopentylethyl, cyclohexylmethyl, cyclohexylethyl, methylbutyl, ethylbutyl, dimethylbutyl, thienylmethyl, thienylethyl, thienylpropyl, cyclopentenylmethyl, cyclopentenylethyl, cyclopentenylpropyl, cyclohexenyl, cyclohexenylmethyl, cyclohexenylethyl, adamantyl, adamantylmethyl, adamantylethyl, phenyl, chlorophenyl, dichlorophenyl, fluorophenyl, difluorophenyl, methoxyphenyl, ethoxyphenyl, methylphenyl, ethylphenyl, propylphenyl, isopropylphenyl, dimethylphenyl, diethylphenyl and benzyl; wherein R⁵ is an acidic group selected from CO₂H and



or a tautomer thereof or a pharmaceutically-acceptable salt thereof.

85. The method of Claim 84 wherein said active compound is selected from compounds and their pharmaceutically-acceptable salts of the group of compounds consisting of
- 1-propyl-4-butyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 - 1,4-dibutyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 - 1-pentyl-4-butyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 - 1-isopentyl-4-butyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 - 1-cyclohexylmethyl-4-butyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 - 1-phenylmethyl-4-butyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 - 1-(2-cyclohexylethyl)-4-butyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 - 1-(2-phenylethyl)-4-butyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 - 1-propyl-4-butyl-5-methyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 - 1,4-dibutyl-5-methyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 - 1-pentyl-4-butyl-5-methyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

2-one;

1-isopentyl-4-butyl-5-methyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

1-cyclohexylmethyl-4-butyl-5-methyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

1-phenylmethyl-4-butyl-5-methyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

1-(2-cyclohexylethyl)-4-butyl-5-methyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

1-(2-phenylethyl)-4-butyl-5-methyl-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

1-propyl-4-butyl-5-chloro-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

1,4-dibutyl-5-chloro-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

1-pentyl-4-butyl-5-chloro-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

1-isopentyl-4-butyl-5-chloro-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;


1-cyclohexylmethyl-4-butyl-5-chloro-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

1-phenylmethyl-4-butyl-5-chloro-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;


1-(2-cyclohexylethyl)-4-butyl-5-chloro-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one; and

1-(2-phenylethyl)-4-butyl-5-chloro-1,3-dihydro-3-[[4-[4-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one.

86. The method of Claim 73 wherein said active compound is selected from a family of compounds of Formula VI




wherein R¹ is selected from methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, 4-methylbutyl, n-pentyl, isopentyl, neopentyl, phenyl, benzyl, phenethyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cyclopropylmethyl, cyclopropylethyl, cyclobutylmethyl, cyclobutylethyl, cyclopentylmethyl, cyclopentylethyl, cyclohexylmethyl, cyclohexylethyl, cyclohexanoyl, methylbutyl, ethylbutyl, dimethylbutyl, thienylmethyl, thienylethyl, thienylpropyl, cyclopentenylmethyl, cyclopentenylethyl, cyclopentenylpropyl, cyclohexenyl, cyclohexenylmethyl, cyclohexenylethyl, adamantyl, adamantylmethyl, adamantylethyl, phenyl, chlorophenyl, dichlorophenyl, fluorophenyl, difluorophenyl, methoxyphenyl, ethoxyphenyl, methylphenyl, ethylphenyl, propylphenyl, isopropylphenyl, dimethylphenyl, diethylphenyl, benzyl, 1-oxo-2-cyclohexylethyl, benzoyl, 1-oxo-2-phenethyl, 1-oxopropyl, 1-oxobutyl, 1-oxopentyl and 2-hydroxybutyl; wherein R⁰ is selected from hydrido, methyl, fluoro, chloro, monofluoromethyl, difluoromethyl, trifluoromethyl, formyl, carboxyl and dimethoxymethyl; wherein R² is selected from methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, n-pentyl, isopentyl, neopentyl, phenyl, benzyl, phenethyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cyclopropylmethyl, cyclopropylethyl, cyclobutylmethyl, cyclobutylethyl, cyclopentylmethyl, cyclopentylethyl, cyclohexylmethyl, cyclohexylethyl, methylbutyl, ethylbutyl, dimethylbutyl, thienylmethyl, thienylethyl, thienylpropyl, cyclopentenylmethyl, cyclopentenylethyl, cyclopentenylpropyl, cyclohexenyl, cyclohexenylmethyl, cyclohexenylethyl, adamantyl, adamantylmethyl, adamantylethyl, phenyl, chlorophenyl, dichlorophenyl, fluorophenyl, difluorophenyl, methoxyphenyl, ethoxyphenyl, methylphenyl, ethylphenyl, propylphenyl, isopropylphenyl, dimethylphenyl, diethylphenyl and benzyl; wherein R⁵ is an acidic group selected from CO₂H and



or a tautomer thereof or a pharmaceutically-acceptable salt thereof.

87. The method of Claim 86 wherein R¹ is selected from methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, 4-methylbutyl, n-pentyl, isopentyl, neopentyl, phenyl, benzyl, phenethyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cyclopropylmethyl, cyclopropylethyl, cyclobutylmethyl, cyclobutylethyl, cyclopentylmethyl, cyclopentylethyl, cyclohexylmethyl, cyclohexylethyl, cyclohexanoyl, methylbutyl, ethylbutyl, dimethylbutyl, thienylmethyl, thienylethyl, thienylpropyl, cyclopentenylmethyl, cyclopentenylethyl, cyclopentenylpropyl, cyclohexenyl, cyclohexenylmethyl, cyclohexenylethyl, adamantyl, adamantylmethyl, adamantylethyl, phenyl, chlorophenyl, dichlorophenyl, fluorophenyl, difluorophenyl, methoxyphenyl, ethoxyphenyl, methylphenyl, ethylphenyl, propylphenyl, isopropylphenyl, dimethylphenyl, diethylphenyl, benzyl, 1-oxo-2-cyclohexylethyl, benzoyl, 1-oxo-2-phenethyl, 1-oxopropyl, 1-oxobutyl, 1-oxopentyl and 2-hydroxybutyl; wherein R⁰ is hydrido; wherein R² is selected from methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, n-pentyl, isopentyl, neopentyl, phenyl, benzyl, phenethyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cyclopropylmethyl, cyclopropylethyl, cyclobutylmethyl, cyclobutylethyl, cyclopentylmethyl, cyclopentylethyl, cyclohexylmethyl, cyclohexylethyl, methylbutyl, ethylbutyl, dimethylbutyl, thienylmethyl, thienylethyl, thienylpropyl, cyclopentenylmethyl, cyclopentenylethyl, cyclopentenylpropyl, cyclohexenyl, cyclohexenylmethyl, cyclohexenylethyl, adamantyl, adamantylmethyl, adamantylethyl, phenyl, chlorophenyl, dichlorophenyl, fluorophenyl, difluorophenyl, methoxyphenyl, ethoxyphenyl, methylphenyl, ethylphenyl, propylphenyl, isopropylphenyl, dimethylphenyl, diethylphenyl and benzyl; wherein R⁵ is an acidic group selected from CO₂H and




or a tautomer thereof or a pharmaceutically-acceptable salt thereof.


88. The method of Claim 87 wherein said active compound is selected from compounds and their pharmaceutically-acceptable salts of the group of compounds consisting of
- 1-propyl-4-butyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 - 1,4-dibutyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 - 1-pentyl-4-butyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 - 1-isopentyl-4-butyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 - 1-cyclohexylmethyl-4-butyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 - 1-phenylmethyl-4-butyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 - 1-(2-cyclohexylethyl)-4-butyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 - 1-(2-phenylethyl)-4-butyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 - 1-propyl-4-butyl-5-methyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 - 1,4-dibutyl-5-methyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 - 1-pentyl-4-butyl-5-methyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

- 1-isopentyl-4-butyl-5-methyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-cyclohexylmethyl-4-butyl-5-methyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-phenylmethyl-4-butyl-5-methyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-(2-cyclohexylethyl)-4-butyl-5-methyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-(2-phenylethyl)-4-butyl-5-methyl-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-propyl-4-butyl-5-chloro-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1,4-dibutyl-5-chloro-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-pentyl-4-butyl-5-chloro-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-isopentyl-4-butyl-5-chloro-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-cyclohexylmethyl-4-butyl-5-chloro-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-phenylmethyl-4-butyl-5-chloro-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 1-(2-cyclohexylethyl)-4-butyl-5-chloro-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one; and
 1-(2-phenylethyl)-4-butyl-5-chloro-1,3-dihydro-3-[[4-[3-(1H-tetrazol-5-yl)-4-pyridinyl]phenyl]methyl]-2H-imidazol-2-one.

89. The method of Claim 73 wherein said active compound is selected from a family of compounds of Formula VII




wherein R¹ is selected from methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, 4-methylbutyl, n-pentyl, isopentyl, neopentyl, phenyl, benzyl, phenethyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cyclopropylmethyl, cyclopropylethyl, cyclobutylmethyl, cyclobutylethyl, cyclopentylmethyl, cyclopentylethyl, cyclohexylmethyl, cyclohexylethyl, cyclohexanoyl, methylbutyl, ethylbutyl, dimethylbutyl, thienylmethyl, thienylethyl, thienylpropyl, cyclopentenylmethyl, cyclopentenylethyl, cyclopentenylpropyl, cyclohexenyl, cyclohexenylmethyl, cyclohexenylethyl, adamantyl, adamantylmethyl, adamantylethyl, phenyl, chlorophenyl, dichlorophenyl, fluorophenyl, difluorophenyl, methoxyphenyl, ethoxyphenyl, methylphenyl, ethylphenyl, propylphenyl, isopropylphenyl, dimethylphenyl, diethylphenyl, benzyl, 1-oxo-2-cyclohexylethyl, benzoyl, 1-oxo-2-phenethyl, 1-oxopropyl, 1-oxobutyl, 1-oxopentyl and 2-hydroxybutyl; wherein R⁰ is selected from hydrido, methyl, fluoro, chloro, monofluoromethyl, difluoromethyl, trifluoromethyl, formyl, carboxyl and dimethoxymethyl; wherein R² is selected from methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, n-pentyl, isopentyl, neopentyl, phenyl, benzyl, phenethyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cyclopropylmethyl, cyclopropylethyl, cyclobutylmethyl, cyclobutylethyl, cyclopentylmethyl, cyclopentylethyl, cyclohexylmethyl, cyclohexylethyl, methylbutyl, ethylbutyl, dimethylbutyl, thienylmethyl, thienylethyl, thienylpropyl, cyclopentenylmethyl, cyclopentenylethyl, cyclopentenylpropyl, cyclohexenyl, cyclohexenylmethyl, cyclohexenylethyl, adamantyl, adamantylmethyl, adamantylethyl, phenyl, chlorophenyl, dichlorophenyl, fluorophenyl, difluorophenyl, methoxyphenyl, ethoxyphenyl, methylphenyl, ethylphenyl, propylphenyl, isopropylphenyl, dimethylphenyl, diethylphenyl and benzyl; wherein R⁵ is an acidic group selected from CO₂H and



or a tautomer thereof or a pharmaceutically-acceptable salt thereof.

90. The method of Claim 89 wherein R¹ is selected from methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, 4-methylbutyl, n-pentyl, isopentyl, neopentyl, phenyl, benzyl, phenethyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cyclopropylmethyl, cyclopropylethyl, cyclobutylmethyl, cyclobutylethyl, cyclopentylmethyl, cyclopentylethyl, cyclohexylmethyl, cyclohexylethyl, cyclohexanoyl, methylbutyl, ethylbutyl, dimethylbutyl, thienylmethyl, thienylethyl, thienylpropyl, cyclopentenylmethyl, cyclopentenylethyl, cyclopentenylpropyl, cyclohexenyl, cyclohexenylmethyl, cyclohexenylethyl, adamantyl, adamantylmethyl, adamantylethyl, phenyl, chlorophenyl, dichlorophenyl, fluorophenyl, difluorophenyl, methoxyphenyl, ethoxyphenyl, methylphenyl, ethylphenyl, propylphenyl, isopropylphenyl, dimethylphenyl, diethylphenyl, benzyl, 1-oxo-2-cyclohexylethyl, benzoyl, 1-oxo-2-phenethyl, 1-oxopropyl, 1-oxobutyl, 1-oxopentyl and 2-hydroxybutyl; wherein R⁰ is hydrido; wherein R² is selected from methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, n-pentyl, isopentyl, neopentyl, phenyl, benzyl, phenethyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cyclopropylmethyl, cyclopropylethyl, cyclobutylmethyl, cyclobutylethyl, cyclopentylmethyl, cyclopentylethyl, cyclohexylmethyl, cyclohexylethyl, methylbutyl, ethylbutyl, dimethylbutyl, thienylmethyl, thienylethyl, thienylpropyl, cyclopentenylmethyl, cyclopentenylethyl, cyclopentenylpropyl, cyclohexenyl, cyclohexenylmethyl, cyclohexenylethyl, adamantyl, adamantylmethyl, adamantylethyl, phenyl, chlorophenyl, dichlorophenyl, fluorophenyl, difluorophenyl, methoxyphenyl, ethoxyphenyl, methylphenyl, ethylphenyl, propylphenyl, isopropylphenyl, dimethylphenyl, diethylphenyl and benzyl; wherein R⁵ is an acidic group selected from CO₂H and



or a tautomer thereof or a pharmaceutically-acceptable salt thereof.

91. The method of Claim 90 wherein said active compound is selected from compounds and their pharmaceutically-acceptable salts of the group of compounds consisting of
- 1-propyl-4-butyl-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 - 1,4-dibutyl-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 - 1-pentyl-4-butyl-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 - 1-isopentyl-4-butyl-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 - 1-cyclohexylmethyl-4-butyl-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 - 1-phenylmethyl-4-butyl-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 - 1-(2-cyclohexylethyl)-4-butyl-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 - 1-(2-phenylethyl)-4-butyl-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 - 1-propyl-4-butyl-5-methyl-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 - 1,4-dibutyl-5-methyl-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;
 - 1-pentyl-4-butyl-5-methyl-1,3-dihydro-3-[[4-[2-(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

2-one;

1-isopentyl-4-butyl-5-methyl-1,3-dihydro-3-[[4-[-2(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

1-cyclohexylmethyl-4-butyl-5-methyl-1,3-dihydro-3-[[4-[-2(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

1-phenylmethyl-4-butyl-5-methyl-1,3-dihydro-3-[[4-[-2(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

1-(2-cyclohexylethyl)-4-butyl-5-methyl-1,3-dihydro-3-[[4-[-2(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

1-(2-phenylethyl)-4-butyl-5-methyl-1,3-dihydro-3-[[4-[-2(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

1-propyl-4-butyl-5-chloro-1,3-dihydro-3-[[4-[-2(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

1,4-dibutyl-5-chloro-1,3-dihydro-3-[[4-[-2(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

1-pentyl-4-butyl-5-chloro-1,3-dihydro-3-[[4-[-2(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

1-isopentyl-4-butyl-5-chloro-1,3-dihydro-3-[[4-[-2(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

1-cyclohexylmethyl-4-butyl-5-chloro-1,3-dihydro-3-[[4-[-2(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

1-phenylmethyl-4-butyl-5-chloro-1,3-dihydro-3-[[4-[-2(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one;

1-(2-cyclohexylethyl)-4-butyl-5-chloro-1,3-dihydro-3-[[4-[-2(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one; and

1-(2-phenylethyl)-4-butyl-5-chloro-1,3-dihydro-3-[[4-[-2(1H-tetrazol-5-yl)-3-pyridinyl]phenyl]methyl]-2H-imidazol-2-one.

92. The method of Claim 79 wherein said active compound is 1,4-dibutyl-1,3-dihydro-3-[[6-[-2(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one or a pharmaceutically-acceptable salt thereof.

93. The method of Claim 79 wherein said active compound is 1-(2-ethylbutyl)-4-butyl-1,3-dihydro-3-[[6-[-2(1H-tetrazol-5-yl)phenyl]-3-pyridinyl]methyl]-2H-imidazol-2-one or a pharmaceutically-acceptable salt thereof.

94. The method of Claim 63 wherein said circulatory disorder is a cardiovascular disorder.

95. The method of Claim 94 wherein said cardiovascular disorder is hypertension.

96. The method of Claim 94 wherein said cardiovascular disorder is congestive heart failure.



European Patent
Office

PARTIAL EUROPEAN SEARCH REPORT

which under Rule 45 of the European Patent Convention
shall be considered, for the purposes of subsequent
proceedings, as the European search report

Application Number

EP 92 10 6050

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl.5)
A	EP-A-0 102 227 (WARNER-LAMBERT CO.) ---		C 07 D 401/14
A	EP-A-0 324 377 (E.I. DU PONT DE NEMOURS AND CO.) ---		A 61 K 31/415
A	US-A-4 600 430 (ELI LILLY AND CO.) ---		C 07 D 401/10
P,A	WO-A-9 204 331 (SMITH KLINE & FRENCH LABORATORIES LTD) -----		C 07 D 401/06
			C 07 D 409/14
			TECHNICAL FIELDS SEARCHED (Int. Cl.5)
			C 07 D A 61 K
INCOMPLETE SEARCH			
<p>The Search Division considers that the present European patent application does not comply with the provisions of the European Patent Convention to such an extent that it is not possible to carry out a meaningful search into the state of the art on the basis of some of the claims</p> <p>Claims searched completely : Claims searched incompletely : Claims not searched : Reason for the limitation of the search:</p> <p>See sheet -C-</p>			
Place of search THE HAGUE		Date of completion of the search 14-07-1992	Examiner DE BUYSER I.A.F.
CATEGORY OF CITED DOCUMENTS		T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons ----- & : member of the same patent family, corresponding document	
X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document			

EPO FORM 150 03.92 (P0607)



EP 92 10 6050 -C-

Claims searched completely: 14,17,20,23,26,29,45,48,51,
54,57,60-62,76,79,82,85,88,91-93

Claims searched incompletely: 1-13,15-16,18-19,21-22,24-25,
27-28,32-44,46-47,49-50,52-53,55-56,58-59,63-75,77-78,80-81,
83-84,86-87,89-90,94-96

Reason for the limitation of the search:

As the drafting of the claims is not clear and concise (Art.83-84 EPO) and encompasses such an enormous amount of products, a complete search is not possible on economic grounds [See Guidelines for Examination in the EPO Part B, Chapter III,2]. Guided by the spirit of the application and the inventive concept as disclosed in the descriptive part of the present application the search has been based on the examples.